

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Burr, SE; Hart, JD; Edwards, T; Baldeh, I; Bojang, E; Harding-Esch, EM; Holland, MJ; Lietman, TM; West, SK; Mabey, DC; Sillah, A; Bailey, RL (2013) Association between Ocular Bacterial Carriage and Follicular Trachoma Following Mass Azithromycin Distribution in The Gambia. *PLoS neglected tropical diseases*, 7 (7). e2347. ISSN 1935-2727 DOI: 10.1371/journal.pntd.0002347

Downloaded from: <http://researchonline.lshtm.ac.uk/1130083/>

DOI: [10.1371/journal.pntd.0002347](https://doi.org/10.1371/journal.pntd.0002347)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

Association between Ocular Bacterial Carriage and Follicular Trachoma Following Mass Azithromycin Distribution in The Gambia

Sarah E. Burr^{1,2*}, John D. Hart¹, Tansy Edwards³, Ignatius Baldeh², Ebrima Bojang², Emma M. Harding-Esch¹, Martin J. Holland¹, Thomas M. Lietman⁴, Sheila K. West⁵, David C. W. Mabey¹, Ansumana Sillah⁶, Robin L. Bailey¹

1 Department of Clinical Research, Faculty of Infectious and Tropical Disease, London School of Hygiene and Tropical Medicine, London, United Kingdom, **2** Medical Research Council Unit, Fajara, Banjul, The Gambia, **3** MRC Tropical Epidemiology Group, London School of Hygiene and Tropical Medicine, London, United Kingdom, **4** Francis I. Proctor Foundation for Research in Ophthalmology, University of California San Francisco, San Francisco, California, United States of America, **5** Dana Center for Preventive Ophthalmology, Johns Hopkins University, Baltimore, Maryland, United States of America, **6** National Eye Health Programme, Ministry of Health and Social Welfare, Kanifing, The Gambia

Abstract

Background: Trachoma, caused by ocular *Chlamydia trachomatis* infection, is the leading infectious cause of blindness, but its prevalence is now falling in many countries. As the prevalence falls, an increasing proportion of individuals with clinical signs of follicular trachoma (TF) is not infected with *C. trachomatis*. A recent study in Tanzania suggested that other bacteria may play a role in the persistence of these clinical signs.

Methodology/Principal Findings: We examined associations between clinical signs of TF and ocular colonization with four pathogens commonly found in the nasopharynx, three years after the initiation of mass azithromycin distribution. Children aged 0 to 5 years were randomly selected from 16 Gambian communities. Both eyes of each child were examined and graded for trachoma according to the World Health Organization (WHO) simplified system. Two swabs were taken from the right eye: one swab was processed for polymerase chain reaction (PCR) using the Amplicor test for detection of *C. trachomatis* DNA and the second swab was processed by routine bacteriology to assay for the presence of viable *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Staphylococcus aureus* and *Moraxella catarrhalis*. Prevalence of TF was 6.2% (96/1538) while prevalence of ocular *C. trachomatis* infection was 1.0% (16/1538). After adjustment, increased odds of TF were observed in the presence of *C. trachomatis* (OR = 10.4, 95%CI 1.32–81.2, p = 0.03), *S. pneumoniae* (OR = 2.14, 95%CI 1.03–4.44, p = 0.04) and *H. influenzae* (OR = 4.72, 95% CI 1.53–14.5, p = 0.01).

Conclusions/Significance: Clinical signs of TF can persist in communities even when ocular *C. trachomatis* infection has been controlled through mass azithromycin distribution. In these settings, TF may be associated with ocular colonization with bacteria commonly carried in the nasopharynx. This may affect the interpretation of impact surveys and the determinations of thresholds for discontinuing mass drug administration.

Citation: Burr SE, Hart JD, Edwards T, Baldeh I, Bojang E, et al. (2013) Association between Ocular Bacterial Carriage and Follicular Trachoma Following Mass Azithromycin Distribution in The Gambia. PLoS Negl Trop Dis 7(7): e2347. doi:10.1371/journal.pntd.0002347

Editor: Jeremiah M. Ngondi, University of Cambridge, United Kingdom

Received: March 13, 2013; **Accepted:** June 19, 2013; **Published:** July 25, 2013

Copyright: © 2013 Burr et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by the Bill and Melinda Gates Foundation (<http://www.gatesfoundation.org/>), grant number 48027. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: sburr@mrc.gm

Introduction

Trachoma, caused by ocular infection with the intracellular bacterium *Chlamydia trachomatis*, remains the leading infectious cause of blindness world-wide. In 1986, The Gambia's first national blindness survey indicated that trachoma was a significant cause of blindness in the country with active trachoma affecting 17% of children aged 0–14 years and very high rates of trichiasis in those 15 years and older [1,2]. In response to these findings, the National Eye Health Programme (formerly the National Eye Care Programme), supported by SightSavers International, established a network of Community Ophthalmic Nurses (CONs) trained and

equipped to screen communities for active trachoma and to conduct trichiasis surgery. Since the 1980s, many trachoma endemic communities may have also benefited from water and sanitation interventions such as well digging and latrine construction programmes via the Departments of Water Resources and Community Development. A Healthy Eyes curriculum, focused on preventive eye care, was introduced into primary schools and a network of individuals (*Nyateros* or 'Friends of the Eye') trained in eye health and ocular first aid was established at the community level.

A decade later, the 1996 national survey found a substantially reduced prevalence of active trachoma (to just over 5% in 0–14

Author Summary

Trachoma, the world's leading infectious cause of blindness, is caused by ocular infection with the bacterium *Chlamydia trachomatis*. In low-prevalence settings and following mass treatment campaigns, clinically active follicular trachoma (TF) can be found in the absence of *C. trachomatis* infection. We carried out this study to investigate associations between ocular carriage of non-chlamydial pathogens and a clinical diagnosis of TF following a mass treatment campaign in The Gambia. We found that children who carried *Streptococcus pneumoniae* or *Haemophilus influenzae* in their eyes were more likely to have been diagnosed with TF than children who did not carry these pathogens. In The Gambia, non-chlamydial pathogens may be inducing or exacerbating TF in the absence of *C. trachomatis* infection.

year olds nationally), but which remained above 10% in four regions of the country [2,3]. In 2006, in the absence of a national survey, trachoma surveys were conducted in two regions (North Bank and Lower River Regions) and showed little change in overall trachomatous inflammation, follicular (TF) prevalence from 1996 [4]. On the basis of these data, The Gambia's plan for trachoma control using mass drug administration (MDA) with azithromycin was approved by the Director of Medical Services in 2006 and The Gambia received over 400,000 doses of azithromycin from the International Trachoma Initiative (ITI) to be distributed in 23 priority health districts. The Partnership for the Rapid Elimination of Trachoma trial (PRET) [5,6] was embedded in this plan, with communities in four priority districts randomized to receive treatment annually for three years [as currently recommended by the World Health Organization (WHO)] or to stop treatment if the prevalence of TF or detected ocular *C. trachomatis* infection fell below 5%. The PRET study areas now have very low rates of both disease and *C. trachomatis* infection [7].

As the prevalence of TF has fallen in The Gambia, a disparity between clinical signs of disease and ocular *C. trachomatis* infection has been documented [4,6,8]. This finding has been mirrored in other communities following implementation of mass treatment [9–12] and in communities where prevalence of TF is low [13–16]. It has been suggested that at least some of the TF now observed may be due to organisms other than *C. trachomatis* [4,16]. In the present study, we aimed to investigate possible associations between ocular colonization with four bacterial pathogens commonly found in the nasopharynx and a clinical diagnosis of TF following the implementation of the MDA campaign in The Gambia.

Methods

Ethics statement

Ethical approval was obtained from the London School of Hygiene and Tropical Medicine Ethics Committee and The Gambia Government/Medical Research Council Unit, The Gambia Joint Ethics Committee under study numbers SCC 1107v2 and L2011.25. Oral consent was obtained from the community leaders and informed written (thumbprint or signature) consent was obtained from each child's guardian at the time of examination. The use of thumbprints in the consent procedure was specifically approved by the ethics committees. The consent form was further signed by an independent witness.

Study design

The PRET trial (ClinicalTrials.gov NCT00792922) was a cluster randomized trial (CRT) carried out in The Gambia, the design of which has been described elsewhere [5]. Briefly, clusters, or enumerations areas (EAs, population 600–800 individuals), were randomly sampled from four districts (two districts in the North Bank Region, two districts from the Western Region; 12 EAs per district). Twenty-four EAs received three annual, community-wide azithromycin treatments and 24 EAs received just one mass treatment at baseline. The 48 EAs were further randomized in a 2×2 factorial design to receive standard treatment coverage (one day visit to EAs by treatment team) versus enhanced coverage (two visits to each EA) during the mass treatments that took place. Randomization of EAs was stratified by district (six EAs per district allocated to either three mass treatments or one). The study described here is ancillary to the PRET trial; 16 of the 24 EAs from North Bank Region were included in this study (8 from each frequency allocation arm, Fig. 1).

Field methods

A cross-sectional survey took place during May and June 2011, coinciding with the 36 month follow-up survey of the PRET trial. For EAs randomized to receive three annual rounds of mass treatment, sample collection took place 12 months following the last round; for EAs randomized to receive a single treatment round, sample collection took place 36 months following treatment (Fig. 1). One hundred children aged 0–5 years were randomly selected from each EA for participation. Eligible children were: resident in the EA, did not have an ocular condition precluding examination or specimen collection, willing to have a sample taken and had a guardian willing to provide consent. Children's faces were initially observed for the presence of nasal and ocular discharge and flies on the eyes. The upper eyelids of each child were everted and the tarsal conjunctiva graded for signs of clinical trachoma using the WHO simplified grading system [17]. Two photographs were then taken of the right everted upper tarsal conjunctiva for use in quality assurance as detailed elsewhere [7]. Samples were collected from the upper tarsal conjunctiva and lower fornix of the right eye using a Dacron swab. Upper tarsal conjunctival swabs were placed into dry vials while lower fornix swabs were placed into vials containing 1 ml skimmed-milk-tryptone-glucose glycerol transport medium (STGG) [18]. Samples were kept on wet ice in the field and frozen at –20°C within 10 hours. Samples were transferred to –70°C upon arrival at the MRC laboratory, within 2 weeks' time.

Laboratory methods

Upper tarsal conjunctival samples were processed for *C. trachomatis* infection using the Amplicor CT/NG kit (Roche Molecular Systems, Branchburg, NJ, USA) as previously described [5]. Lower fornix swabs were thawed at room temperature and cultured on blood and chocolate agar plates for the detection of *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Staphylococcus aureus* and *Moraxella catarrhalis*. Blood agar plates were incubated at 37°C in either 5% CO₂ or under anaerobic conditions. Chocolate agar plates were incubated at 37°C in 5% CO₂. Isolates were identified using conventional bacteriology techniques. All laboratory personnel were masked to the clinical trachoma grade.

Sample size

A sample size of 1568 subjects is required to detect an odds ratio (OR) of 2 for a univariate association between TF (anticipated to




Figure 1. Study allocation.

doi:10.1371/journal.pntd.0002347.g001

be 7%) and a bacterial pathogen (assumed to be present in 15% of participants without TF), assuming 80% power to detect univariate associations with 95% confidence. It was expected that a sample size of approximately 1600 children would be achieved from 16 EAs in which a random sample of 100 children was to be taken from each.

Analysis methods

Overall unadjusted prevalence is presented with corresponding exact binomial 95% confidence intervals (CI). Prevalence is also presented adjusted for age, sex and district using random effects logistic regression models to account for any underlying variation detected between households and between EAs.

The prevalence of TF was tabulated by each pathogen, age, sex, district, *C. trachomatis* infection and time since mass treatment (12 months for those in the annual treatment arm versus 36 months for those in the baseline treatment arm). Clustering of TF and bacterial pathogens at household and EA level was investigated using random effects logistic regression models with random intercepts for household and EAs. Likelihood ratio tests (LRTs) were used to compare models with and without the additional random effects terms.

Univariate associations with TF were determined using random effects logistic regression models with and without the exposure of

interest. To assess confounding of the association between each pathogen and TF, unadjusted Mantel-Haenszel ORs for the effect of each pathogen on TF were compared to the ORs adjusted for each exposure. A 10% change in the OR was considered to indicate confounding, if the exposure was associated with both the pathogen of interest and TF. Modification of the effect of each pathogen on TF by other exposures was assessed in exploratory analyses using Mantel-Haenszel methods.

Following identification of univariate associations with TF and confounders of the effect of each pathogen on TF, a multivariate model was built adjusting for each pathogen associated with TF and including any confounding exposures. Independent risk factors for TF were also considered for inclusion in the final model on the basis of a LRT p -value < 0.1. The strength of association with TF for each pathogen and exposure in the final model was estimated from a LRT comparing adjusted models with and without the pathogen of exposure.

Predictors of discharge (nasal and ocular) were identified using random effects logistic regression with discharge as the outcome, accounting for between household and/or between EA variation where appropriate and building a final multivariable model using the same approach as for TF.

Results

In the 16 EAs under study, 1569 children aged 0–5 years were randomly sampled for measurement of trachoma outcomes as part of the PRET CRT. Of these, laboratory results for both *C. trachomatis* PCR and conventional bacteriology were obtained for 1538 children (98.0%). For the 31 children with missing laboratory results, tabulations showed no evidence that children were not missing at random. Analysis was restricted to the results obtained for those 1538 children with complete data.

Prevalence of bacterial colonization

Bacteriology culture detected at least one non-chlamydial pathogen in the eyes of 475 (30.9%) children. *S. aureus* and *S. pneumoniae* were most commonly detected occurring in more than 14% of children (Table 1). Colonization with *H. influenzae* and *M. catarrhalis* was less common. Sixteen children (1.0%) had evidence of *C. trachomatis* infection.

Dual colonization was detected in 53 (3.4%) children; most commonly with *S. pneumoniae* and *S. aureus* (37 children, 2.4%). Dual colonization with *S. pneumoniae* and *H. influenzae* was found in 10 children (0.7%). Six children positive for *C. trachomatis* infection carried one non-chlamydial pathogen concurrently (0.4%); 2 children had *C. trachomatis* and *H. influenzae*, 2 children had *C. trachomatis* with *S. aureus* and 2 children had *C. trachomatis* and *M. catarrhalis*. No children were found to have *C. trachomatis* infection concurrent with *S. pneumoniae*.

In random effects logistic regression adjusting for each pathogen alone (null models), there was evidence that children carrying *S. pneumoniae* were clustered at the household level as demonstrated by evidence of between-household variation ($p < 0.001$) but not at EA level, after accounting for clustering by household ($p = 0.11$). *H. influenzae* carriage also clustered at household level ($p = 0.01$) but not at EA level ($p = 0.46$). The reverse was found for *M. catarrhalis* carriage, which clustered at EA level ($p = 0.01$) but not household ($p = 1.00$). *S. aureus* carriage clustered at both household and EA levels ($p < 0.001$). *C. trachomatis* infection cases were clustered at EA level ($p < 0.001$).

Associations with TF

Ninety-six children (6.2%) were found to have clinical signs of TF; no children were diagnosed as having trachomatous inflammation – intense (TI). TF cases were clustered at both EA and household levels ($p < 0.001$). After accounting for between-household and between-EA variation, age ($p = 0.004$), *C. trachomatis* infection ($p = 0.04$) and district ($p = 0.08$) were associated with TF in univariate analyses (Table 2). Sex ($p = 0.64$), time since mass treatment was carried out in the EA (whether 12 or 36 months

previously; $p = 0.76$) and flies on the eyes ($p = 0.317$) were not associated with TF.

In the univariate analysis accounting for between-household and between-EA variation of TF but no other covariates, increased odds of TF were observed in the presence of *S. pneumoniae* (OR = 1.69, 95%CI 0.85–3.33, $p = 0.14$), *H. influenzae* (OR = 3.40, 95%CI 1.20–9.66, $p = 0.027$) and *S. aureus* (OR = 1.30, 95%CI 0.63–2.67, $p = 0.48$) but a significant effect was observed only for *H. influenzae* in this analysis (Table 2). There were no signs of TF amongst any of the children with ocular *M. catarrhalis* carriage. Dual carriage with more than one non-chlamydial pathogen was not associated with TF ($p = 0.072$).

The presence of *S. pneumoniae* was associated with age ($p < 0.001$), with fewer older children having the organism. A similar association was found between *H. influenzae* and age ($p < 0.001$). After adjustment for age, the association between *S. pneumoniae* and TF was significant at the 5% level (OR = 2.08, 95%CI 1.01–4.30, $p = 0.05$). The effect of *H. influenzae* also appeared stronger after adjustment for age (OR = 4.45, 95%CI 1.44–13.7, $p = 0.01$). *C. trachomatis* infection was not associated with age ($p = 0.54$).

In building a final random effects regression model accounting for between-household and between-EA variation, *H. influenzae* and *S. pneumoniae* were added in turn, followed by age since this was identified as a confounder of the effect of both *H. influenzae* and *S. pneumoniae* on TF. *C. trachomatis* infection and district were also added to the model in turn as independent risk factors for TF.

Multivariable results suggest a two-fold increase in the odds of TF for children with ocular *S. pneumoniae* carriage and a four-fold increase for carriers of *H. influenzae* (Table 2). *C. trachomatis* infection was significantly associated with TF, but estimates of the magnitude of effect were imprecise due to low prevalence of *C. trachomatis* infection and TF in the sample.

Predictors of nasal and ocular discharge

Overall, 839 (54.6%) children were observed to have nasal discharge on the day of screening. In the final regression model, after adjustment for underlying variation at household level, *S. pneumoniae* ($p = 0.02$), *M. catarrhalis* ($p = 0.01$) and TF ($p = 0.05$) were associated with increased odds of nasal discharge (Table 3). Age was also associated with nasal discharge ($p < 0.001$) with increased odds observed for 2–3 year olds versus 0–1 year olds and decreased odds for 4–5 year olds versus 0–1 year olds. In the univariate analysis, flies on the eyes were associated with nasal discharge ($p < 0.001$) but we have not included this variable in the multivariate model as it is not clear whether nasal discharge is the causal effect for flies on the eyes.

Ocular discharge was observed for 190 (12.4%) children. A final model adjusted for household and EA level clustering of cases.

Table 1. Prevalence of ocular bacterial colonization, *C. trachomatis* infection and TF.

	N	Colonization, n	Prevalence (exact binomial 95% CI, percent scale)	Adjusted Prevalence* (95% CI, percent scale)
<i>S. pneumoniae</i>	1538	219	14.2 (12.5–16.1)	12.5 (11.0–13.8)
<i>H. influenzae</i>	1538	57	3.7 (2.8–4.8)	2.6 (1.6–3.6)
<i>S. aureus</i>	1538	233	15.2 (13.4–17.0)	14.7 (13.9–15.6)
<i>M. catarrhalis</i>	1538	19	1.2 (0.7–1.9)	1.1 (0.8–1.5)
<i>C. trachomatis</i>	1538	16	1.0 (0.6–1.7)	1.0 (0.8–1.1)
TF	1538	96	6.2 (5.1–7.6)	6.1 (5.4–6.9)

*Adjusted for age (0–1, 2–3, 4–5 years), sex, district and any underlying variation between households and/or EAs as appropriate.

doi:10.1371/journal.pntd.0002347.t001

Table 2. Associations between bacterial pathogens and TF in 0–5 year olds.

Characteristic	N	TF, n (%)	Univariate OR (95% CI)*	p-value	Multivariate OR (95% CI)*	p-value	
All children	1538	96 (6.2)	-	-	-	-	
<i>S. pneumoniae</i>	No	1319	78 (5.9)	1	1		
	Yes	219	18 (8.2)	1.69 (0.85–3.33)	0.140	2.14 (1.03–4.44)	0.044
<i>H. influenzae</i>	No	1481	88 (5.9)	1	1		
	Yes	57	8 (14.0)	3.40 (1.20–9.66)	0.027	4.72 (1.53–14.5)	0.009
<i>S. aureus</i>	No	1305	81 (6.2)	1	-		
	Yes	233	15 (6.4)	1.30 (0.63–2.67)	0.480	-	
<i>M. catarrhalis</i>	No	1519	96 (6.3)	-	-		
	Yes	19	0 (0)	-	-		
At least one non-chlamydial pathogen	No	1063	60 (5.6%)	1	-		
	Yes	475	36 (7.6%)	1.93 (1.09–3.43)	0.022	-	
Number of non-chlamydial pathogens	0	1036	60 (5.6%)	1	0.072	-	
	1	422	31 (7.4%)	1.96 (1.07–3.57)		-	
	2	53	5 (9.4%)	1.79 (0.51–6.31)		-	
Age (years)	0–1	464	17 (3.7)	1	0.012	1	0.004
	2–3	588	47 (8.0)	2.72 (1.36–5.43)		3.24 (1.55–6.75)	
	4–5	486	32 (6.6)	1.89 (0.93–3.87)		2.46 (1.14–5.32)	
Sex	Male	820	54 (6.6)	1	-		
	Female	718	42 (5.9)	0.89 (0.53–1.47)	0.637	-	
<i>C. trachomatis</i>	No	1522	93 (6.1)	1	1		
	Yes	16	3 (18.8)	8.54 (1.21–60.4)	0.041	10.4 (1.32–81.2)	0.033
District	Lower Baddibu	726	30 (4.1)	1	1		
	Central Baddibu	812	66 (8.1)	2.25 (0.95–5.34)	0.075	2.32 (0.94–5.70)	0.076
Time since mass treatment	12 months	737	44 (6.0)	1	-		
	36 months	801	52 (6.5)	1.16 (0.45–2.99)	0.757	-	
Flies on the eyes†	No	227	16 (7.1)	1	-		
	Yes	1309	80 (6.1)	1.44 (0.71–2.90)	0.317	-	

*Univariate and multivariate ORs are estimated from random effects logistic regression models accounting for between-household and between-EA variation.

†2 missing values for flies on the eyes.

doi:10.1371/journal.pntd.0002347.t002

Results suggest increased odds of ocular discharge in children carrying *H. influenzae* ($p=0.06$) and possibly in male children ($p=0.07$) (Table 4). Age was again associated with ocular discharge ($p=0.04$) with the same pattern as for nasal discharge; increased odds in 2–3 year olds versus 0–1 year olds and decreased odds for 4–5 years versus 0–1 year olds.

Discussion

The PRET trial documented a near disappearance of ocular *C. trachomatis* infection from a catchment area of greater than 65,000 people in the two years following the initiation of MDA; biannual surveys carried out in the four study districts showed no evidence of ocular *C. trachomatis* infection 12 and 18 months post-MDA initiation and just 0.1% prevalence at 6 and 24 months [7]. Yet prevalence of TF over the same time period remained relatively high, reaching more than 10% in some EAs in the North Bank Region of the country. While one cannot rule-out episodes of *C. trachomatis* infection in the intervening months, it is difficult to see how *C. trachomatis* infection alone could sustain the clinical signs of disease during that time. Low-level exposure to *C. trachomatis* may

not be detected by the Amplicor PCR method but it is unclear whether such low bacterial loads result in onward transmission of the microorganism.

In the 16 EAs included in this study, prevalence of ocular *C. trachomatis* infection 36 months post-MDA was 1% while prevalence of clinical signs of TF was 6%. While the association between active ocular *C. trachomatis* infection and TF was strong, it does not account for all the TF recorded. Clinical signs of TF were also associated with ocular *S. pneumoniae* and *H. influenzae* carriage. Both *S. pneumoniae* and *H. influenzae* are well known causes of bacterial conjunctivitis although not generally associated with follicular conjunctivitis [19]. It may be that localised inflammation caused by these microorganisms recalls a follicular response in children who have previously been exposed to *C. trachomatis*. Alternatively, ocular carriage could predispose the eye to infection by *C. trachomatis* thereby inducing subsequent TF. In the present study, ocular carriage with *S. pneumoniae* was associated with nasal discharge while *H. influenzae* was associated with ocular discharge. *Musca sorbens*, which in some environments can act as the vector of trachoma, is more often found on the faces of children with ocular

Table 3. Associations with nasal discharge in 0–5 year olds.

Characteristic	N	Nasal discharge, n (%)	Univariable OR (95% CI)*	p-value	Multivariable OR (95% CI)*	p-value	
All children	1538	839 (54.6)	-	-	-	-	
<i>S. pneumoniae</i>	No	1319	700 (53.1)	1	1		
	Yes	219	139 (63.5)	1.59 (1.15–2.20)	0.005	1.52 (1.08–2.13)	0.015
<i>H. influenzae</i>	No	1481	808 (54.6)	1	-		
	Yes	57	31 (54.9)	0.99 (0.55–1.78)	0.965	-	
<i>S. aureus</i>	No	1305	715 (54.8)	1	-		
	Yes	233	124 (53.2)	0.94 (0.69–1.28)	0.698	-	
<i>M. catarrhalis</i>	No	1519	824 (54.3)	1	1		
	Yes	19	15 (79.0)	3.58 (1.09–11.8)	0.022	4.07 (1.22–13.6)	0.013
TF	No	1442	776 (53.8)	1	1		
	Yes	96	63 (65.6)	1.67 (1.03–2.72)	0.034	1.63 (0.99–2.68)	0.051
<i>C. trachomatis</i>	No	1522	827 (54.3)	1	-		
	Yes	16	12 (75.0)	2.67 (0.75–9.49)	0.114	-	
Age (years)	0–1	464	260 (56.1)	1	<0.001	1	<0.001
	2–3	588	368 (62.6)	1.36 (1.03–1.79)		1.40 (1.06–1.85)	
	4–5	486	211 (43.4)	0.57 (0.43–0.75)		0.59 (0.44–0.79)	
Sex	Male	820	455 (55.5)	1	-		
	Female	718	384 (53.5)	0.90 (0.72–1.21)	0.342	-	
District	Lower Baddibu	726	393 (54.1)	1	-		
	Central Baddibu	812	446 (54.9)	1.00 (0.79–1.28)	0.972	-	
Time since mass treatment	12 months	737	395 (53.6)	1	-		
	36 months	801	444 (55.4)	1.10 (0.86–1.41)	0.433	-	
Flies on the eyes†	No	227	65 (28.6)	1	-		
	Yes	1309	634 (48.4)	2.54 (1.78–3.60)	<0.001	-	

*Univariable and multivariable ORs are estimated from random effects logistic regression models accounting for between-household and between-EA variation, as appropriate.

†2 missing values for flies on the eyes.

doi:10.1371/journal.pntd.0002347.t003

and nasal discharge than on children with no discharge [20,21]; in our study flies on the eyes was associated with nasal discharge. If ocular colonization with these pathogens itself causes discharge, this may in turn increase the risk of *C. trachomatis* transmission.

One limitation of this study was that samples were kept frozen at -20°C for approximately two weeks before transport to the laboratory and subsequent storage at -70°C . Published data suggest these handling conditions are sufficient for cultivation of these respiratory pathogens [22,23] however, we cannot discount the possibility that storage conditions negatively impacted the viability of our samples resulting in an underestimate in ocular carriage prevalence. In addition, it is unclear whether the isolation of non-chlamydial, bacterial pathogens from the conjunctiva in a cross-sectional survey such as this is indicative of established ocular infection. A positive bacteriology culture may simply indicate transient carriage in the conjunctiva following introduction of bacteria by fingers or fomites. Indeed, nasopharyngeal (NP) carriage rates of *S. pneumoniae* in The Gambia are high with up to 90% of children under the age of five years being carriers [24,25]. It is likely that children continually self-inoculate their eyes by touching their hands to their faces, possibly causing clinical signs of trachoma in children who have previously been exposed to *C. trachomatis*.

Our study has also shown that prevalence of ocular carriage of *S. aureus* is high, yet no association of this pathogen with TF was found and its carriage was not associated with either nasal or ocular discharge. Unlike the other pathogens detected, *S. aureus* is a part of the normal skin flora and imprecise sampling of the lower fornix could have introduced skin contamination into the swabs. However, the prevalence of ocular *S. aureus* carriage that we have detected (15%) is consistent with an earlier study carried out in Gambian communities where 11% of adults presenting for trichiasis surgery were found to carry the pathogen [15]. Another study detected *S. aureus* in the eyes of 30% of adults with trachomatous scarring and in 30% of healthy controls' however, the sample size of this study was small (13 cases and 7 controls [26]).

Previous studies on the role of non-chlamydial pathogens in trachoma have been conducted in The Gambia but with a focus on trachomatous scarring (TS) and trichiasis (TT) rather than on TF [15,26]. Bacterial pathogens were more frequently isolated from eyes with TS than from control eyes however, this difference was not statistically significant [26]. In contrast, TT was significantly associated with bacterial pathogens in the eye, an association strengthened with increasing disease severity as measured by the number of lashes touching the eye [15,26]. *S.*

Table 4. Associations with ocular discharge in 0–5 year olds.

Characteristic	N	Ocular discharge, n (%)	Univariable OR (95% CI)*	p-value	Multivariable OR (95% CI)*	p-value	
All children	1538	190 (12.4)	-	-	-	-	
<i>S. pneumoniae</i>	No	1319	166 (12.6)	1	-	-	
	Yes	219	24 (11.0)	0.85 (0.52–1.39)	0.502	-	
<i>H. influenzae</i>	No	1481	177 (12.0)	1	1	-	
	Yes	57	13 (22.8)	2.24 (1.09–4.60)	0.035	2.07 (1.00–4.31)	0.059
<i>S. aureus</i>	No	1305	163 (12.5)	1	-	-	
	Yes	233	27 (11.6)	0.97 (0.60–1.56)	0.884	-	
<i>M. catarrhalis</i>	No	1519	188 (12.4)	1	-	-	
	Yes	19	2 (10.5)	0.81 (0.17–4.00)	0.795	-	
TF	No	1442	172 (11.9)	1	-	-	
	Yes	96	18 (18.8)	1.49 (0.81–2.76)	0.213	-	
<i>C. trachomatis</i>	No	1522	187 (12.3)	1	-	-	
	Yes	16	3 (18.8)	1.86 (0.43–8.13)	0.426	-	
Age (years)	0–1	464	64 (13.8)	1	0.030	1	0.038
	2–3	588	82 (14.0)	1.06 (0.72–1.56)	-	1.11 (0.75–1.63)	-
	4–5	486	44 (9.1)	0.63 (0.40–0.97)	-	0.66 (0.42–1.02)	-
Sex	Male	820	113 (13.8)	1	-	1	-
	Female	718	77 (10.7)	0.74 (0.53–1.04)	0.082	0.73 (0.52–1.03)	0.069
District	Lower Baddibu	726	85 (11.7)	1	-	-	-
	Central Baddibu	812	105 (12.9)	1.18 (0.62–2.25)	0.615	-	-
Time since mass treatment	12 months	737	77 (10.5)	1	-	-	-
	36 months	801	113 (14.1)	1.40 (0.76–2.57)	0.288	-	-
Flies on the eyes†	No	227	193 (85.0)	1	-	-	-
	Yes	1309	1155 (88.2)	1.25 (0.80–1.96)	0.338	-	-

*Univariable and multivariable ORs are estimated from random effects logistic regression models accounting for between-household and between-EA variation, as appropriate.

†2 missing values for flies on the eyes.

doi:10.1371/journal.pntd.0002347.t004

pneumoniae was the most common pathogen isolated from TT cases [26].

The role of non-chlamydial pathogens in TF has recently been studied in a trachoma endemic community in Tanzania, in this case prior to rather than post-MDA [16]. Prevalence of TF was higher than that reported here, with 13.7% of children under 10 years of age displaying signs of active disease. Prevalence of ocular *C. trachomatis* infection was also higher (5.3%) yet no association was seen between TF and a positive Amplicor PCR result. Ocular *S. pneumoniae* and *H. influenzae* carriage however, were both found to be associated with TF, consistent with our present findings.

The prevalence of ocular carriage of *S. pneumoniae* in the Tanzanian study was modestly lower than that which we report here (10.2% versus 14.2%). This may reflect the broader age distribution of children sampled in the Tanzanian study (all children under 10 years of age) as NP carriage rates are known to be higher in younger children. However, prevalence of NP carriage also appears to be lower in Tanzania than in The Gambia, although data are limited [27,28]. The heptavalent pneumococcal conjugate vaccine (PCV-7) was introduced into the Expanded Programme of Immunization (EPI) in The Gambia in August 2009; those children who were less than two years of age when recruited into our study were likely vaccinated. Available evidence however, suggests prevalence of NP carriage of *S. pneumoniae* in The Gambia is not affected by PCVs due to

replacement with non-vaccine serotypes [25,29]. Indeed we found younger children were more likely to carry *S. pneumoniae* than older children, consistent with data of NP carriage from this country indicating prevalence of carriage is associated with decreasing age [24].

H. influenzae carriage rates were markedly different between the two studies with *H. influenzae* type B (Hib) being isolated from 14% and non-Hib *H. influenzae* from 9% of children in the Tanzanian study. We found ocular *H. influenzae* colonization in just 3.7% of children sampled. Hib conjugate vaccine has been used in the Gambian EPI since 1997 and coverage rates are believed to be high [30]. Prevalence of NP Hib carriage in Western Gambia fell from 12% pre-vaccination to 0.25% five years following introduction of the vaccine into the EPI [31]. We therefore did not type for Hib in the present study. It is likely that high Hib vaccine coverage has contributed to the low levels of ocular *H. influenzae* carriage reported here; while Hib vaccination does not cross protect against other *H. influenzae* serotypes, there are only limited data to support the assertion that Hib vaccination results in serotype replacement by other capsulated strains in carriage [32]. Although we did not use molecular techniques to characterise the *H. influenzae* isolates, we speculate the majority of them would be unencapsulated (non-typeable) strains, which are unaffected by the Hib vaccine. Such strains are often associated with conjunctivitis [33] and recently were shown to account for most of the *H.*

influenzae carried in the nasopharynx of a cohort of Gambian infants [34]. Despite the lower prevalence of ocular *H. influenzae* carriage seen in The Gambia as compared to Tanzania, we also found an association between TF and this pathogen.

The Gambia is one of many countries incorporating MDA into its national trachoma control strategy. The ITI has distributed more than 250 million azithromycin treatments in 19 trachoma endemic countries and plans to be active in 42 countries by 2015. While MDA with azithromycin is associated with a significant decrease in pharyngeal carriage of *S. pneumoniae*, this effect is short-lived with carriage rates returning to baseline within two months following a single round of treatment [35,36]. Repeated MDA at six or 12 month intervals also fails to affect prevalence of NP carriage of *S. pneumoniae* in the long-term [37–39]. The effect of MDA with azithromycin on carriage of *H. influenzae* is less well characterized, yet available data from The Gambia indicate pharyngeal carriage of the bacterium is unaffected by three doses (20 mg/kg) given over the course of two weeks [40]. It is therefore likely that as ocular *C. trachomatis* infection is controlled through mass azithromycin distribution, the contribution of these non-chlamydial pathogens to clinical signs of TF will become more apparent.

According to WHO policy, the prevalence of TF is currently the principal determinant for discontinuing MDA [41]. If non-chlamydial pathogens are inducing or exacerbating TF in the

absence of *C. trachomatis* infection, communities may end up receiving unnecessary treatment. Furthermore, proposed criteria for trachoma elimination also relate to prevalence of clinical signs [42,43] and as countries strive to reach their control targets, the information obtained from surveillance may be misleading. The role of these pathogens in the persistence of TF post-MDA therefore warrants further study; the key question is whether TF in these circumstances is likely to progress to conjunctival scarring and blindness.

Acknowledgments

We thank Pateh Makalo, Mass Laye, Sarjo Dibba, Muhamed Jallow, Robyn Damary-Homan, Jane Whitton, Anita Wadagni and Omar Camara for their work in the field. We would also like to thank the community leaders and villagers for their participation in the study.

Author Contributions

Conceived and designed the experiments: SEB JDH IB RLB. Performed the experiments: SEB JDH IB EB. Analyzed the data: TE. Contributed reagents/materials/analysis tools: MJH TML SKW DCWM RLB. Wrote the paper: SEB JDH TE EMHE MJH TML SKW DCWM AS RLB. Coordinated the project: SEB JDH AS EMHE. Interpreted the data: SEB TE RLB.

References

- Faal H, Minassian D, Sowa S, Foster A (1989) National survey of blindness and low vision in The Gambia: results. *Br J Ophthalmol* 73: 82–87. Available: <http://bjoo.bmj.com/content/73/2/82.short>. Accessed 11 March 2013.
- Dolin P, Faal H, Johnson G (1998) Trachoma in the Gambia. *Br J Ophthalmol* 82: 930–933. Available: <http://bjoo.bmj.com/content/82/8/930.short>. Accessed 11 March 2013.
- Faal H, Minassian DC, Dolin PJ, Mohamed a a, Ajewole J, et al. (2000) Evaluation of a national eye care programme: re-survey after 10 years. *The British journal of ophthalmology* 84: 948–951. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1723659&tool=pmcentrez&rendertype=abstract>.
- Harding-Esch EM, Edwards T, Sillah A, Sarr I, Roberts CH, et al. (2009) Active trachoma and ocular Chlamydia trachomatis infection in two Gambian regions: on course for elimination by 2020? *PLoS neglected tropical diseases* 3: e573. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2791206&tool=pmcentrez&rendertype=abstract>. Accessed 29 September 2011.
- Stare D, Harding-Esch E, Munoz B, Bailey R, Mabey D, et al. (2011) Design and baseline data of a randomized trial to evaluate coverage and frequency of mass treatment with azithromycin: the Partnership for Rapid Elimination of Trachoma (PRET) in Tanzania and The Gambia. *Ophthalmic epidemiology* 18: 20–29. Available: <http://www.ncbi.nlm.nih.gov/pubmed/21275593>. Accessed 31 August 2012.
- Harding-Esch EM, Edwards T, Mkocho H, Munoz B, Holland MJ, et al. (2010) Trachoma prevalence and associated risk factors in the gambia and Tanzania: baseline results of a cluster randomised controlled trial. *PLoS neglected tropical diseases* 4: e861. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2970530&tool=pmcentrez&rendertype=abstract>. Accessed 5 September 2012.
- Harding-Esch EM, Sillah A, Edwards T, Burr SE, Hart JD, et al. (2013) Mass Treatment with Azithromycin for Trachoma: When Is One Round Enough? Results from the PRET Trial in The Gambia. *PLoS neglected tropical diseases* 7: e2115. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3681669&tool=pmcentrez&rendertype=abstract>. Accessed 24 June 2013.
- Burton MJ, Holland MJ, Makalo P, Aryce EAN, Sillah A, et al. (2010) Profound and sustained reduction in Chlamydia trachomatis in The Gambia: a five-year longitudinal study of trachoma endemic communities. *PLoS neglected tropical diseases* 4: e835. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2950148&tool=pmcentrez&rendertype=abstract>. Accessed 3 October 2012.
- Solomon AW, Holland MJ, Alexander NDE, Massae P a, Aguirre A, et al. (2004) Mass treatment with single-dose azithromycin for trachoma. *The New England journal of medicine* 351: 1962–1971. Available: <http://www.ncbi.nlm.nih.gov/pubmed/15525721>.
- Solomon AW, Harding-Esch E, Alexander NDE, Aguirre A, Holland MJ, et al. (2008) Two doses of azithromycin to eliminate trachoma in a Tanzanian community. *The New England journal of medicine* 358: 1870–1871. Available: <http://www.ncbi.nlm.nih.gov/pubmed/18434662>.
- Cumberland P, Edwards T, Hailu G, Harding-Esch E, Andreasen A, et al. (2008) The impact of community level treatment and preventative interventions on trachoma prevalence in rural Ethiopia. *International journal of epidemiology* 37: 549–558. Available: <http://www.ncbi.nlm.nih.gov/pubmed/18356196>. Accessed 16 November 2012.
- Biebesheimer JB, House J, Hong KC, Lakew T, Alemayehu W, et al. (2009) Complete local elimination of infectious trachoma from severely affected communities after six biannual mass azithromycin distributions. *Ophthalmology* 116: 2047–2050. Available: <http://www.ncbi.nlm.nih.gov/pubmed/19744717>. Accessed 16 November 2012.
- Baral K, Osaki S, Shreshtha B, Panta CR, Boulter A, et al. (1999) Reliability of clinical diagnosis in identifying infectious trachoma in a low-prevalence area of Nepal. *Bulletin of the World Health Organization* 77: 461–466. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2557684&tool=pmcentrez&rendertype=abstract>. Accessed 16 November 2012.
- Thein J, Zhao P, Liu H, Xu J, Jha H, et al. (2002) Does clinical diagnosis indicate ocular chlamydial infection in areas with a low prevalence of trachoma? *Ophthalmic epidemiology* 9: 263–269. Available: <http://www.ncbi.nlm.nih.gov/pubmed/12187424>. Accessed 16 November 2012.
- Burton MJ, Kintch F, Jallow O, Sillah a, Bah M, et al. (2005) A randomised controlled trial of azithromycin following surgery for trachomatous trichiasis in the Gambia. *The British journal of ophthalmology* 89: 1282–1288. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1772881&tool=pmcentrez&rendertype=abstract>. Accessed 8 December 2011.
- Burton MJ, Hu VH, Massae P, Burr SE, Chevallier C, et al. (2011) What is causing active trachoma? The role of nonchlamydial bacterial pathogens in a low prevalence setting. *Investigative ophthalmology & visual science* 52: 6012–6017. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3176035&tool=pmcentrez&rendertype=abstract>. Accessed 13 November 2012.
- Thylefors B, Dawson CR, Jones BR, West SK, Taylor HR (1987) A simple system for the assessment of trachoma and its complications. *Bulletin of the World Health Organization* 65: 477–483. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2491032&tool=pmcentrez&rendertype=abstract>. Accessed 13 November 2012.
- GIBSON LF, KHOURY JT (1986) Storage and survival of bacteria by ultra-freeze. *Letters in Applied Microbiology* 3: 127–129. Available: <http://doi.wiley.com/10.1111/j.1472-765X.1986.tb01565.x>. Accessed 16 November 2012.
- Krachmer JH, Mannis MJ, Holland EJ (1997) *Cornea: cornea and external disease: clinical diagnosis and management*. Volume 2. St. Louis, Missouri: Mosby-Year Book, Inc. p.
- Emerson PM, Bailey RL, Mahdi OS, Walraven GE, Lindsay SW (2000) Transmission ecology of the fly *Musca sorbens*, a putative vector of trachoma. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 94: 28–32. Available: <http://www.ncbi.nlm.nih.gov/pubmed/10748893>. Accessed 2 October 2012.

21. Miller K, Pakpour N, Yi E, Melese M, Alemayehu W, et al. (2004) Pesky trachoma suspect finally caught. *The British journal of ophthalmology* 88: 750–751. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1772198&tool=pmcentrez&rendertype=abstract>. Accessed 2 October 2012.
22. O'Brien KL, Bronsdon MA, Dagan R, Yagupsky P, Janco J, et al. (2001) Evaluation of a medium (STGG) for transport and optimal recovery of *Streptococcus pneumoniae* from nasopharyngeal secretions collected during field studies. *Journal of clinical microbiology* 39: 1021–1024. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=87867&tool=pmcentrez&rendertype=abstract>. Accessed 2 October 2012.
23. Kajjalainen T, Ruokokoski E, Ukkonen P, Herva E (2004) Survival of *Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Moraxella catarrhalis* frozen in skim milk- tryptone-glucose-glycerol medium. *Journal of clinical microbiology* 42: 412–414. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=321668&tool=pmcentrez&rendertype=abstract>. Accessed 2 October 2012.
24. Hill PC, Akisanya A, Sankareh K, Cheung YB, Saaka M, et al. (2006) Nasopharyngeal carriage of *Streptococcus pneumoniae* in Gambian villagers. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America* 43: 673–679. Available: <http://www.ncbi.nlm.nih.gov/pubmed/16912937>. Accessed 16 November 2012.
25. Cheung Y-B, Zaman SM A, Nsekpong ED, Van Beneden C a, Adegbola R a, et al. (2009) Nasopharyngeal carriage of *Streptococcus pneumoniae* in Gambian children who participated in a 9-valent pneumococcal conjugate vaccine trial and in their younger siblings. *The Pediatric infectious disease journal* 28: 990–995. Available: <http://www.ncbi.nlm.nih.gov/pubmed/19536041>. Accessed 5 July 2011.
26. Burton MJ, Adegbola R a, Kinteh F, Ikumapayi UN, Foster A, et al. (2007) Bacterial infection and trachoma in the gambia: a case control study. *Investigative ophthalmology & visual science* 48: 4440–4444. Available: <http://www.ncbi.nlm.nih.gov/pubmed/17898263>. Accessed 24 July 2012.
27. Leung MHY, Oriyo NM, Gillespie SH, Charalambous BM (2011) The adaptive potential during nasopharyngeal colonisation of *Streptococcus pneumoniae*. *Infection, genetics and evolution: journal of molecular epidemiology and evolutionary genetics in infectious diseases* 11: 1989–1995. Available: <http://www.ncbi.nlm.nih.gov/pubmed/21925618>. Accessed 16 November 2012.
28. Moyo SJ, Steinbakk M, Aboud S, Mkopi N, Kasubi M, et al. (2012) Penicillin resistance and serotype distribution of *Streptococcus pneumoniae* in nasopharyngeal carrier children under 5 years of age in Dar es Salaam, Tanzania. *Journal of medical microbiology* 61: 952–959. Available: <http://www.ncbi.nlm.nih.gov/pubmed/22442292>. Accessed 16 November 2012.
29. Egere U, Townend J, Roca A, Akisanya A, Bojang A, et al. (2012) Indirect effect of 7-valent pneumococcal conjugate vaccine on pneumococcal carriage in newborns in rural Gambia: a randomised controlled trial. *PLoS one* 7: e49143. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3504064&tool=pmcentrez&rendertype=abstract>. Accessed 15 May 2013.
30. World Health Organization (2012) WHO UNICEF coverage estimates WHO World Health Organization: Immunization, Vaccines And Biologicals. Vaccine preventable diseases Vaccines monitoring system 2012 Global Summary Reference Time Series: GMB. Available: http://apps.who.int/immunization_monitoring/en/globalsummary/timeseries/TSWUcoverageByCountry.cfm?country=GMB. Accessed 29 January 2013.
31. Adegbola RA, Secka O, Lahai G, Lloyd-Evans N, Njie A, et al. (2005) Elimination of *Haemophilus influenzae* type b (Hib) disease from The Gambia after the introduction of routine immunisation with a Hib conjugate vaccine: a prospective study. *Lancet* 366: 144–150. Available: <http://www.ncbi.nlm.nih.gov/pubmed/16005337>. Accessed 2 October 2012.
32. Dabernat H, Plisson-Sauné M-A, Delmas C, Séguy M, Faucon G, et al. (2003) *Haemophilus influenzae* carriage in children attending French day care centers: a molecular epidemiological study. *Journal of clinical microbiology* 41: 1664–1672. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=153885&tool=pmcentrez&rendertype=abstract>. Accessed 16 November 2012.
33. Murphy TF, Faden H, Bakaletz LO, Kyd JM, Forsgren A, et al. (2009) Nontypeable *Haemophilus influenzae* as a pathogen in children. *The Pediatric infectious disease journal* 28: 43–48. Available: <http://www.ncbi.nlm.nih.gov/pubmed/19057458>. Accessed 2 October 2012.
34. Kwambana BA, Barer MR, Bottomley C, Adegbola RA, Antonio M (2011) Early acquisition and high nasopharyngeal co-colonisation by *Streptococcus pneumoniae* and three respiratory pathogens amongst Gambian new-borns and infants. *BMC infectious diseases* 11: 175. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3129300&tool=pmcentrez&rendertype=abstract>. Accessed 17 July 2012.
35. Leach AJ, Shelby-James TM, Mayo M, Gratten M, Laming AC, et al. (1997) A prospective study of the impact of community-based azithromycin treatment of trachoma on carriage and resistance of *Streptococcus pneumoniae*. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America* 24: 356–362. Available: <http://www.ncbi.nlm.nih.gov/pubmed/9114185>. Accessed 23 August 2012.
36. Batt SL, Charalambous BM, Solomon AW, Knirsch C, Massae PA, et al. (2003) Impact of azithromycin administration for trachoma control on the carriage of antibiotic-resistant *Streptococcus pneumoniae*. *Antimicrobial agents and chemotherapy* 47: 2765–2769. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=182606&tool=pmcentrez&rendertype=abstract>. Accessed 28 March 2013.
37. Fry a M, Jha HC, Lietman TM, Chaudhary JSP, Bhatta RC, et al. (2002) Adverse and beneficial secondary effects of mass treatment with azithromycin to eliminate blindness due to trachoma in Nepal. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America* 35: 395–402. Available: <http://www.ncbi.nlm.nih.gov/pubmed/12145722>.
38. Gaynor BD, Chidambaram JD, Cevallos V, Miao Y, Miller K, et al. (2005) Topical ocular antibiotics induce bacterial resistance at extraocular sites. *The British journal of ophthalmology* 89: 1097–1099. Available: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1772818&tool=pmcentrez&rendertype=abstract>. Accessed 31 January 2013.
39. Haug S, Lakew T, Habtemariam G, Alemayehu W, Cevallos V, et al. (2010) The decline of pneumococcal resistance after cessation of mass antibiotic distributions for trachoma. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America* 51: 571–574. Available: <http://www.ncbi.nlm.nih.gov/pubmed/20649409>. Accessed 23 August 2012.
40. Adegbola RA, Mulholland EK, Bailey R, Secka O, Sadiq T, et al. (1995) Effect of azithromycin on pharyngeal microflora. *The Pediatric infectious disease journal* 14: 335–337. Available: <http://www.ncbi.nlm.nih.gov/pubmed/7603826>. Accessed 23 August 2012.
41. World Health Organization (2003) World Health Organization Report of the 2nd Global Scientific Meeting on Trachoma. Available: <http://www.who.int/blindness/2nd GLOBAL SCIENTIFIC MEETING.pdf>.
42. Organization WH (2011) Report of the fifteenth meeting of the WHO alliance for the elimination of blinding trachoma by 2020. Available: <http://www.who.int/blindness/publications/GET15REPORT4.pdf>. Accessed 1 February 2013.
43. World Health Organization (2012) Report of the sixteenth meeting of the WHO alliance for the Elimination of Blinding Trachoma by 2020 pp. 14–16. Available: <http://www.who.int/blindness/publications/GET16Report.pdf>.