

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Rudge, JW; Coker, R; (2013) Human to human transmission of H7N9. *BMJ* (Clinical research ed), 347. f4730. ISSN 0959-8138 DOI: <https://doi.org/10.1136/bmj.f4730>

Downloaded from: <http://researchonline.lshtm.ac.uk/1229385/>

DOI: <https://doi.org/10.1136/bmj.f4730>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: Creative Commons Attribution Non-commercial
<http://creativecommons.org/licenses/by-nc/3.0/>

<https://researchonline.lshtm.ac.uk>

EDITORIALS

Human to human transmission of H7N9

Limited transmission between humans is not surprising

James W Rudge *lecturer*¹, Richard Coker *professor*^{1 2}

¹Communicable Diseases Policy Research Group, London School of Hygiene and Tropical Medicine, Mahidol University, Bangkok 10400, Thailand;

²Saw Swee Hock School of Public Health, National University of Singapore, Singapore

Since the new avian influenza virus, H7N9, first emerged in China, a primary concern has been whether it might spread between humans. The vast majority of the 133 confirmed cases reported so far seem to be epidemiologically unconnected, with many patients reporting a recent history of exposure to live poultry, which are suspected to be a main reservoir for the virus. Although an earlier study did report two family clusters of H7N9 cases, it was unclear whether these clusters resulted from person to person transmission or simply from exposure to a common animal source of infection.¹

In the linked paper (doi:10.1136/bmj.f4752) by Qi and colleagues, a detailed investigation into one of these clusters provides the strongest evidence yet of H7N9 transmission between humans.² The index case, a 60 year old man, was likely to have been infected at a nearby live poultry market, and subsequently developed a severe and ultimately fatal respiratory illness. His 32 year old daughter, who provided prolonged bedside care for her father before his admission to intensive care, later also became fatally infected. With no indication that the daughter was exposed to live poultry within the days before becoming sick, along with almost 100% genetic similarity between the viruses isolated from each patient, the evidence points to transmission from father to daughter.

As the authors acknowledge, there are some limitations to the study but, on balance, human to human transmission looks probable. So does this imply that H7N9 has come one step closer towards adapting fully to humans? Probably not. Crucially, there is still no evidence of sustained transmission among humans—all 43 close contacts of these two patients, including a son in law who also helped care for the father, tested negative for infection. In addition, the receptor binding sites of the viruses from the two patients are no more adapted towards humans than those of other available H7N9 isolates. In many ways, the evidence corroborates, rather than challenges, previous assertions that the transmissibility of H7N9 between humans is currently low.

Indeed, the occasional transmission event from human to human appears to be the norm rather than the exception for influenza viruses that sporadically cross the species barrier into humans.

Limited human to human transmission has been reported for highly pathogenic avian influenza H5N1,^{3 4} which continues to cause (usually fatal) infections in humans, as well as another bird flu subtype, H7N7, which caused an outbreak of mostly mild infections in the Netherlands in 2003.⁵ To observe some transmission of H7N9 from human to human is therefore not surprising, and does not necessarily indicate that the virus is on course to develop sustained transmission among humans.

Nevertheless, several traits of H7N9 are of particular concern. The linked paper² comes close on the heels of studies showing airborne transmissibility of H7N9 between ferrets in the laboratory, a mammalian model.^{6 7} Also, it is now well documented that owing to its non-lethality in birds, H7N9 can spread undetected through avian populations. In addition, Chinese surveillance data suggest that the number of confirmed human cases is just the tip of the iceberg—many mild cases are likely to have passed undetected.⁸ The upside of this is that the actual fatality rate among H7N9 cases is likely to be substantially lower than that observed among confirmed cases.⁹ The flipside is that the incidence of human infections, and therefore opportunities for H7N9 to adapt to humans or to re-assort through mixed influenza infections, could be much greater than for other bird flu viruses such as H5N1.

Although the number of H7N9 cases has fallen abruptly since April 2013, with no new cases reported for several weeks, we have been warned to expect a resurgence later in the year owing to seasonal effects on transmission.¹⁰ Thus, while the paper by Qi and colleagues² might not suggest that H7N9 is any closer to delivering the next pandemic, it does provide a timely reminder of the need to remain extremely vigilant: the threat posed by H7N9 has by no means passed.

Competing interests: We have read and understood the BMJ Group policy on declaration of interests and declare that RC has received funding via research grants from Roche to the London School of Hygiene and Tropical Medicine of \$1.8m over five years, to conduct surveillance on flu in Indonesia (on which JWR has also been involved), and \$1m over four years to support a clinical registry of human cases of H5N1 in South East Asia. No other conflicts of interest are declared.

Provenance and peer review: Commissioned; not externally peer reviewed.

- 1 Li Q, Zhou L, Zhou M, Chen Z, Li F, Wu H, et al. Preliminary report: epidemiology of the avian influenza A (H7N9) outbreak in China. *N Engl J Med* 2013, doi:10.1056/NEJMoa1304617.
- 2 Qi X, Qian Y-H, Bao C-J, Guo X-L, Cui L-B, Tang F-Y, et al. Probable person to person transmission of novel avian influenza A (H7N9) virus in Eastern China, 2013: epidemiological investigation. *BMJ* 2013;347:f4752.
- 3 Wang H, Feng Z, Shu Y, Yu H, Zhou L, Zu R, et al. Probable limited person-to-person transmission of highly pathogenic avian influenza A (H5N1) virus in China. *Lancet* 2008;371:1427-34.
- 4 Ungchusak K, Auewarakul P, Dowell SF, Kitphati R, Auwanit W, Puthavathana P, et al. Probable person-to-person transmission of avian influenza A (H5N1). *N Engl J Med* 2005;352:333-40.
- 5 Du Ry van Beest Holle M, Meijer A, Koopmans M, de Jager CM. Human-to-human transmission of avian influenza A/H7N7, The Netherlands, 2003. *Euro Surveill* 2005;10:264-8.
- 6 Zhang Q, Shi J, Deng G, Guo J, Zeng X, He X, et al. H7N9 influenza viruses are transmissible in ferrets by respiratory droplet. *Science* 2013, doi:10.1126/science.1240532.
- 7 Belsler JA, Gustin KM, Pearce MB, Maines TR, Zeng H, Pappas C, et al. Pathogenesis and transmission of avian influenza A (H7N9) virus in ferrets and mice. *Nature* 2013, doi:10.1038/nature12391.
- 8 Ip DK, Liao Q, Wu P, Gao Z, Cao B, Feng L, et al. Detection of mild to moderate influenza A/H7N9 infection by China's national sentinel surveillance system for influenza-like illness: case series. *BMJ* 2013;346:f3693.
- 9 Yu H, Cowling BJ, Feng L, Lau EH, Liao Q, Tsang TK, et al. Human infection with avian influenza A H7N9 virus: an assessment of clinical severity. *Lancet* 2013;382:138-45.
- 10 Centers for Disease Control and Prevention. Avian influenza A (H7N9) virus. 2013. www.cdc.gov/flu/avianflu/h7n9-virus.htm.

Cite this as: *BMJ* 2013;347:f4730

© BMJ Publishing Group Ltd 2013