News Roundup [abridged Versions Appear In The Paper Journal]

UK cancer survival rates are no worse than rest of Europe

BMJ 2005; 330 doi: http://dx.doi.org/10.1136/bmj.330.7495.808-b (Published 07 April 2005) Cite this as: BMJ 2005;330:808

- Article
- · Related content
- Article metrics
- Rapid responses
- Response

Survival is not the same as mortality!

It is unfortunate that the BMJ chose to comment on the analysis by Sir Richard Doll and Jill Boreham in the British Journal of Cancer(1) with a headline that bears little relation to the text of their news item.(2) As Doll and Boreham acknowledged, mortality is a function of both incidence and survival. It is not possible to draw conclusions about survival trends from an analysis restricted to mortality trends. For cancers for which survival has been low for many years (e.g. lung, stomach), mortality trends are largely determined by trends in incidence, and are a poor indicator of the quality of treatment. Survival trends also reflect more than just health service performance: for example, detailed studies by the EUROCARE Working Group suggest that patients in the UK are more likely to have advanced disease at the time of diagnosis. (3,4) If lower survival indicates delays in presentation, referral, diagnosis or treatment, rather than biologically more aggressive disease, there may be public health opportunities to improve outcome, alongside those from advances in therapy.

The online, unabridged version of the news item also states that survival comparisons may be unreliable unless carried out within controlled trials. Unfortunately, less than 5% of adult cancer patients in the UK are treated on trial protocols, and comparisons of mortality statistics are also subject to bias, which is why, for example, mass screening interventions are best tested in randomised controlled trials. Furthermore, the assignment and coding of the underlying cause of death are by no means free from error, and this can affect international comparisons of mortality.(5) Genuine concerns also exist about the validity of international comparisons of population-based survival,(6,7) but regardless of the UK's position in any European "league table" of cancer survival, the fact that it has been possible to demonstrate a benefit of specialisation and multidisciplinary team working in the UK,(8) coupled with evidence of under-capacity,(9) suggest to us that the current programme of investment in, and re-organisation of, cancer services in the

UK is justified. It would be unfortunate if policy-makers were to conclude, from mortality comparisons alone, that this investment may be unnecessary. Some of us have argued previously that cancer control strategies are best informed by examining trends in incidence, survival and mortality alongside each other.(10) We continue to hold that view.

David H Brewster,

Director, Scottish Cancer Registry, Edinburgh, UK

Michel P Coleman,

Professor of Epidemiology and Vital Statistics, London School of Hygiene and Tropical Medicine, London, UK

David Forman,

Director of Information and Research, Northern and Yorkshire Cancer Registry and Information Service, Leeds, UK

Henrik Møller,

Director, Thames Cancer Registry, London, UK

Mike J Quinn,

Director, National Cancer Intelligence Centre, Office for National Statistics, London, UK

References

- 1. Doll R, Boreham J. Recent trends in cancer mortality in the UK. Br J Cancer 2005; 92: 1329-1335
- 2. Dobson R. News roundup: UK cancer survival rates are no worse than rest of Europe. BMJ 2005; 330: 808.
- 3. Gatta G, Capocaccia R, Sant M, Bell CMJ, Coebergh JWW, Damhuis RAM, et al. Understanding variations in survival for colorectal cancer in Europe: a EUROCARE high resolution study. Gut 2000; 47: 533-538.
- 4. Sant M, Allemani C, Capocaccia R, Hakulinen T, Aareleid T, Coebergh JW, et al. Stage at diagnosis is a key explanation of differences in breast cancer survival across Europe. Int J Cancer 2003; 106:416-422.
- 5. Boyle P. Relative value of incidence and mortality data in cancer research. Recent Results in Cancer Research 1989; 114: 41-63.
- 6. Cookson JB. Cancer survival. Lancet 2000; 356: 1611.
- 7. Forman D, Gatta G, Capocaccia R, Janssen-Heijnen ML, Coebergh JW. Cancer survival. Lancet 2001; 357: 555.
- 8. Selby P, Gillis C, Haward R. Benefits from specialised cancer

care. Lancet 1996; 348: 313-318.

- 9. O'Rourke N, Edwards R. Lung cancer treatment waiting times and tumour growth. Clin Oncol 2000; 12: 141-144.
- 10. Coleman MP, Babb P, Stockton D, Forman D, Møller H. Trends in breast cancer incidence, survival, and mortality. Lancet 2000; 356: 590-591.

Competing interests:

None declared

Competing interests: No competing interests

19 April 2005

David H Brewster
Director, Scottish Cancer Registry
Michel P Coleman, David Forman, Henrik Moller, and Mike J Quinn.
Information Services, NHS National Services Scotland, Gyle Square, Edinburgh, EH12 9EB
Click to like:

