

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Clark, TG; Mallard, K; Coll, F; Preston, M; Assefa, S; Harris, D; Ogwang, S; Mumbowa, F; Kirenga, B; O'Sullivan, DM; Okwera, A; Eisenach, KD; Joloba, M; Bentley, SD; Ellner, JJ; Parkhill, J; Jones-Lpez, EC; McNerney, R (2013) Elucidating emergence and transmission of multidrug-resistant tuberculosis in treatment experienced patients by whole genome sequencing. *PloS one*, 8 (12). e83012. ISSN 1932-6203 DOI: <https://doi.org/10.1371/journal.pone.0083012>

Downloaded from: <http://researchonline.lshtm.ac.uk/1440290/>

DOI: [10.1371/journal.pone.0083012](https://doi.org/10.1371/journal.pone.0083012)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

Table S4 Genes with SNP densities greater than 10 per kilobase.

In addition to *pncA* and *gid*, two other genes were found to have SNP densities greater than 20, a hypothetical protein *Rv0095c* and *esxO*, which encodes a secretory protein of the Esat-6 family believed to be of importance for virulence and growth.

Gene	Size	Numbers SNPs	Density/kb
<i>Rv0095c</i>	411	13	31.63
<i>pncA</i>	561	15	26.73
<i>esxO</i>	285	7	24.56
<i>gid</i>	675	15	22.22
<i>Rv0277A</i>	258	5	19.38
<i>esxL</i>	285	5	17.54
<i>lppB</i>	663	11	16.59
<i>ubiA</i>	909	13	14.30
<i>TB7.3</i>	216	3	13.89
<i>Rv2561</i>	294	4	13.61
<i>esxD</i>	324	4	12.35
<i>Rv0061</i>	426	5	11.74
<i>mazE1</i>	174	2	11.49
<i>Rv2628</i>	363	4	11.02
<i>esxN</i>	285	3	10.53
<i>vapC6</i>	384	4	10.42

Distribution of all densities: Min. 0.3178; 1st Qu. 1.2720; Median 1.2720; Mean 2.3920; 3rd Qu. 2.9630; Max. 31.6