

LONDON  
SCHOOL of  
HYGIENE  
& TROPICAL  
MEDICINE



LSHTM Research Online

N'guessan, R; Ngufor, C; Kudom, AA; Boko, P; Odjo, A; Malone, D; Rowland, M; (2014) Mosquito Nets Treated with a Mixture of Chlorfenapyr and Alphacypermethrin Control Pyrethroid Resistant *Anopheles gambiae* and *Culex quinquefasciatus* Mosquitoes in West Africa. PloS one, 9 (2). e87710. ISSN 1932-6203 DOI: <https://doi.org/10.1371/journal.pone.0087710>

Downloaded from: <http://researchonline.lshtm.ac.uk/1520198/>

DOI: <https://doi.org/10.1371/journal.pone.0087710>

**Usage Guidelines:**

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact [researchonline@lshtm.ac.uk](mailto:researchonline@lshtm.ac.uk).

Available under license: <http://creativecommons.org/licenses/by/2.5/>

<https://researchonline.lshtm.ac.uk>

# Mosquito Nets Treated with a Mixture of Chlorfenapyr and Alphacypermethrin Control Pyrethroid Resistant *Anopheles gambiae* and *Culex quinquefasciatus* Mosquitoes in West Africa

Raphael N'Guessan<sup>1,2,3\*</sup>, Corine Ngufor<sup>1,2,3</sup>, Andreas A. Kudom<sup>4</sup>, Pelagie Boko<sup>1</sup>, Abibathou Odjo<sup>1</sup>, David Malone<sup>5</sup>, Mark Rowland<sup>2,3</sup>

**1** Centre de Recherche Entomologique de Cotonou, Cotonou, Benin, **2** London School of Hygiene and Tropical Medicine, London, United Kingdom, **3** Pan-African Malaria Vector Research Consortium (PAMVERC), London, United Kingdom, **4** Department of Entomology and Wildlife, School of Biological Sciences, University of Cape Coast, Cape Coast, Ghana, **5** Innovative Vector Control Consortium, Liverpool, United Kingdom

## Abstract

**Background:** The effectiveness of insecticide treated nets is under threat across Africa south of the Sahara from the selection of pyrethroid resistance in *Anopheles gambiae* mosquitoes. To maintain progress against malaria it is necessary to identify alternative residual insecticides for mosquito nets. Mixtures of pyrethroid and insecticides with novel mode of action provide scope for both improved control and management of resistance through concurrent exposure to unrelated insecticides.

**Methods:** The pyrrole chlorfenapyr and the pyrethroid alphacypermethrin were tested individually and as a mixture on mosquito nets in an experimental hut trial in southern Benin against pyrethroid resistant *An. gambiae* and *Culex quinquefasciatus* mosquitoes. The nets were deliberately holed to simulate the effect of wear and tear.

**Results:** The nets treated with the mixture of chlorfenapyr 200 mg/m<sup>2</sup> and alphacypermethrin 25 mg/m<sup>2</sup> killed a proportion of *An. gambiae* (77%, 95%CI: 66–86%) significantly greater than nets treated with alphacypermethrin 25 mg/m<sup>2</sup> (30%, 95%CI: 21–41%) but not significantly different from nets treated with chlorfenapyr 200 mg/m<sup>2</sup> (69%, 95%CI: 57–78%). The nets treated with the mixtures procured personal protection against *An. gambiae* biting (58–62%) by a greater margin than the alphacypermethrin treated net (39%), whereas the chlorfenapyr treated net was not protective. A similar trend in mortality and blood feeding inhibition between treatments was observed in *Cx. quinquefasciatus* to that seen in *An. gambiae*, although the effects were lower. A mixture of alphacypermethrin with chlorfenapyr applied at 100 mg/m<sup>2</sup> had an effect similar to the mixture with chlorfenapyr at 200 mg/m<sup>2</sup>.

**Conclusion:** The effectiveness of ITNs against pyrethroid resistant mosquitoes was restored by the mixture: the alphacypermethrin component reduced human-vector contact while the chlorfenapyr controlled pyrethroid-resistant mosquitoes. The complementary action of these unrelated insecticides demonstrates that the combination on nets has potential for preventing malaria transmission in areas compromised by the spread of pyrethroid resistance.

**Citation:** N'Guessan R, Ngufor C, Kudom AA, Boko P, Odjo A, et al. (2014) Mosquito Nets Treated with a Mixture of Chlorfenapyr and Alphacypermethrin Control Pyrethroid Resistant *Anopheles gambiae* and *Culex quinquefasciatus* Mosquitoes in West Africa. PLoS ONE 9(2): e87710. doi:10.1371/journal.pone.0087710

**Editor:** David J. Sullivan Jr, Johns Hopkins Bloomberg School of Public Health, United States of America

**Received:** September 4, 2013; **Accepted:** January 2, 2014; **Published:** February 3, 2014

**Copyright:** © 2014 N'Guessan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** Project grant was obtained from Innovative Vector Control Consortium (IVCC), Liverpool, UK from a programme grant of the Bill & Melinda Gates Foundation (<http://www.ivcc.com/>). The Bill & Melinda Gates Foundation had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. However the IVCC did advise on the study design via their Expert Scientific Committee.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: Raphael.N'Guessan@lshtm.ac.uk

## Introduction

Long lasting insecticidal nets (LLINs) are considered best practice for malaria vector control because they are effective, reliable, robust and relatively simple to deliver even in remote regions [1]. The recent reductions in malaria-associated morbidity and mortality across sub-Saharan Africa is largely attributed to the massive roll out of LLINs during the last decade [1–2]. Owing to their low cost, longer residual activity and safety, the pyrethroids remain the ideal insecticides for treating LLINs [1,3]. Unfortu-

nately, resistance to pyrethroids is spreading fast across Africa south of the Sahara and has now been reported in malaria vectors in 27 countries [4].

While a negative epidemiological impact of pyrethroid resistance on malaria control has yet to be demonstrated unequivocally, an increasing number of reports have indicated that pyrethroid resistance is capable of undermining the effectiveness of LLINs [5–6]. The situation in southern Benin is particularly grave since the premier brands of LLIN give only limited personal protection and kill fewer mosquitoes in that region than in regions

of susceptibility [5]. It is now recognised that if nothing is done and further selection of resistance to pyrethroids leads to failure of LLINs, the progress achieved so far in reducing the burden of malaria could be reversed [7].

The need to identify alternative insecticides that can circumvent resistance and preserve the effectiveness of LLINs has become critical. Some alternative insecticides (such as the organophosphates and carbamates) have been tested on nets [8–9]. These candidates show toxicity but generally lack the irritancy of pyrethroids, an important property for reducing mosquito biting rates and providing personal protection to users of insecticide treated nets (ITNs). Mosquito nets can be treated with a mixture of pyrethroid and non-pyrethroid insecticides to maximise insecticidal efficacy against pyrethroid resistant mosquitoes while providing a degree of excito-repellency. Such mixtures may also provide opportunity for insecticide resistance management because insect phenotypes which are not killed by one component due to resistance will be controlled by the other provided they are not resistant to both insecticides [10–11].

An alternative insecticide to pyrethroids that is presently under development is chlorfenapyr, a pyrrole which owing to its novel mode of action is active against both pyrethroid resistant and susceptible anophelines and culicines [12–13]. When evaluated on mosquito nets against wild mosquitoes in experimental huts, chlorfenapyr was found to be more effective than pyrethroids at killing resistant *Anopheles* and *Culex* [14–15]. The use of this pyrrole by itself on mosquito nets neither deters nor repels and so needs to be combined with an excito-repellent insecticide [15]. Chlorfenapyr has already shown great promise when applied as IRS in conjunction with LLINs for improved transmission control of pyrethroid resistant mosquitoes [16]. Preliminary studies with a mixture of chlorfenapyr and alphacypermethrin on ITNs has shown good potential for control of *An arabiensis* [17].

The objective of the trial reported here was to determine in experimental huts the potential of mosquito nets treated with a mixture of chlorfenapyr and the pyrethroid alphacypermethrin to protect against and control host-seeking *An gambiae* and *Cx quinquefasciatus* that are strongly resistant to pyrethroids.

## Materials and Methods

### Study site and experimental huts

The study was conducted on a private land at Akron, a village on the outskirts of Porto Novo, Benin. The owner of the land gave permission to conduct the study on this site. The site supported breeding of *An gambiae* M form that are pyrethroid-resistant due to high frequency of *kdr* (>90%) and increased activity of cytochrome P450s [18]. The nuisance mosquito *Cx quinquefasciatus* is present year round and shows resistance to pyrethroids, carbamates and organophosphates [19]. Five experimental huts were selected for the trial. The hut design was the West African type recommended by WHO [20].

### Mosquito net treatments and trial procedure

The nets were made of white 100-denier polyester (SiamDutch Mosquito Netting Co., Bangkok, Thailand). To simulate damaged nets, six standardized holes (each measuring 4 cm×4 cm) were cut into the sides and ends of each net as recommended by WHO [20]. The nets were treated with SC formulations of alphacypermethrin (Fendona 60SC, BASF, Germany) and chlorfenapyr (Phantom 240SC, BASF, Germany) in aqueous solution either separately or mixed together. The treatment arms were: (i) untreated control net, (ii) chlorfenapyr 200 mg/m<sup>2</sup> treated net, (iii) alphacypermethrin 25 mg/m<sup>2</sup> treated net, (iv) chlorfenapyr

100 mg/m<sup>2</sup> and alphacypermethrin 25 mg/m<sup>2</sup> mixture treated net, and (v) chlorfenapyr 200 mg/m<sup>2</sup> and alphacypermethrin 25 mg/m<sup>2</sup> mixture treated net.

The 5 treatments were randomly allocated to the 5 experimental huts and rotated weekly between huts. Adult volunteers slept in the huts from 20.00 - 05.00 and were rotated between huts on consecutive nights. Informed consent to participate was given before recruitment and daily chemoprophylaxis was provided until 4 weeks after the trial. Each morning the volunteers helped to collect mosquitoes from inside the rooms, nets and verandah traps. Mosquito collections were made on 36 nights over six weeks between April and June 2010. The identification to species, mortality counts and determination of feeding status and gonotrophic condition were made in the laboratory. Live mosquitoes were held in netted plastic cups and provided with 10% honey solution; mortality was recorded after 72 h. Climatic information for each day of the trial was collected from the weather station based in Porto Novo.

The principal aim was to assess the efficacy of treatments relative to untreated control nets in terms of (i) deterrence: the proportional reduction in the number of mosquitoes entering huts with treated nets, (ii) insecticide induced exiting rates: estimated from the proportions of mosquitoes collected from the verandahs of treatment and control huts, (iii) blood-feeding inhibition rate: the reduction in the proportion of blood fed mosquitoes in huts with treated nets compared to huts with untreated nets =  $(1 - \text{proportion bloodfed in treatment} / \text{proportion bloodfed in control}) \times 100$ ; (iv) mortality rate: the proportion of mosquitoes dying within 72 h, corrected for control mortality; (v) personal protection.

### Data analysis

The analysis of numbers of mosquitoes collected within huts (overall total, blood-fed and dead totals) was carried out using negative binomial regression after adjusting for the effects of hut and sleeper. The analysis of treatment effects on the proportions of mosquitoes blood-feeding, exiting or killed was carried out using logistic regression, with treatments as fixed effects and huts, sleepers as random effects. All statistical analysis was conducted by using STATA 9 software (STATA Corp., College Station, USA).

### Species and resistance characterisation

Samples of *An. gambiae* s.l. reared from larval collections near the trial site were identified to species using PCR [21] and to molecular form using PCR RFLP [22].

WHO test kits, lined with test papers treated with alphacypermethrin in silicon oil were used to determine susceptibility of *An. gambiae* and *Cx. quinquefasciatus* females reared from larval collections.

PCR diagnostic test for detection of *kdr* mutations was carried out on *An. gambiae* and *Cx. quinquefasciatus* mosquitoes as described by Martinez-Torres et al. [23].

### Toxicology

Chlorfenapyr has a WHO toxicological classification III, an LD50 oral toxicity in rats of >400 mg/kg body weight and acute dermal toxicity >2000 mg/kg; a category of similar to many insecticides used in public health [24]. A risk assessment of the use of chlorfenapyr on nets was undertaken by BASF toxicologists using assumptions, parameters and default values defined in the WHO generic risk assessment model [25]. The calculated exposure levels to chlorfenapyr were all below the relevant dermal and systemic acceptable exposure levels for repeated exposure. Exposure was deemed acceptable based on conservative scenarios

from the WHO model, indicating safe use of the chlorfenapyr-treated nets for the intended use.

### Ethical clearance

Written informed consent was obtained from all volunteers recruited to the experimental hut studies. The study was approved by ethics committees of the London School of Hygiene and Tropical Medicine and the Ministry of Health in Benin.

## Results

### Experimental Hut Trial

Over the six week trial, 515 *Anopheles gambiae* s.l., 3764 *Culex quinquefasciatus* and 453 *Mansonia* females were caught in the huts. Only the data on *An. gambiae* s.l. and *Cx. quinquefasciatus* were analysed further (Table 1&2).

Compared to huts containing untreated nets the entry rates of both species into huts with ITNs were up to 40% less ( $p < 0.001$ ) (Table 1&2). Hut entry rates did not differ significantly between huts with alphacypermethrin, chlorfenapyr or mixture treated nets.

In huts where nets were treated with alphacypermethrin or alphacypermethrin-chlorfenapyr mixtures the proportions of *An. gambiae* and *Cx. quinquefasciatus* that exited into verandahs were significantly greater than huts where nets were untreated or treated with chlorfenapyr ( $p < 0.0001$ ) (Table 1&2). For *An. gambiae* the proportions that exited into verandahs did not differ between huts with chlorfenapyr treated or untreated nets (Table 1), whereas for *Cx. quinquefasciatus* the proportion that exited into verandahs was greater for chlorfenapyr treated nets ( $p < 0.001$ ) (Table 2).

Blood-feeding inhibition of *An. gambiae* was not evident where nets were treated with chlorfenapyr alone (figure 1A). There was some blood-feeding inhibition of *An. gambiae* where nets were treated with alphacypermethrin alone (22%,  $P = 0.042$ ), and further inhibition where nets were treated with the mixtures (35%, 51%,  $P < 0.01$ ). The difference in bloodfeeding inhibition between the mixture with chlorfenapyr 200 mg/m<sup>2</sup> and alphacypermethrin alone was not significant ( $P = 0.061$ ).

Blood-feeding inhibition rates were generally higher for *Cx. quinquefasciatus* than for *An. gambiae*, reaching 28% inhibition for the chlorfenapyr net, 81% for the alphacypermethrin net and 78% and 82% for the mixtures (Figure 1). For *Cx. quinquefasciatus* the inhibition of blood-feeding was attributable more to the pyrethroid component than to the chlorfenapyr component.

Personal protection against the biting of *An. gambiae* ranged from 58% to 62% with the insecticide mixtures, higher than with either insecticide when applied to nets alone (Table 1). Personal protection from the mixtures was higher still against *Cx. quinquefasciatus* (86% to 90%) but was not significantly different from alphacypermethrin net (Table 2).

Mortality rates of *An. gambiae* and *Cx. quinquefasciatus* where nets were treated with alphacypermethrin were less than 30%, and were presumably due to the high level of pyrethroid resistance in the two species (Figure 2). Nets treated with the insecticide mixtures induced three times higher mortality of *An. gambiae* (75%) and *Cx. quinquefasciatus* (47–50%) than the alphacypermethrin treated nets ( $P < 0.01$ ). Mortalities of *An. gambiae* with the mixtures were not significantly different to that induced by chlorfenapyr alone ( $P = 0.082$  for the mixture with chlorfenapyr 100 mg/m<sup>2</sup> and  $P = 0.065$  for chlorfenapyr 200 mg/m<sup>2</sup>) and hence the mixture's toxicity is attributable mainly to the chlorfenapyr component (Figure 2). The difference in mortality of *An. gambiae* between mixtures with low and high dosages of chlorfenapyr was not significant ( $P = 0.917$ ).

A similar trend in mortality to that seen in *An. gambiae* was observed in *Cx. quinquefasciatus*, although the effect of each treatment on mortality was 20–50% lower.

The average daily temperature (minimum, minimum) during the first week of the trial was 31.5 degree (27.7, 34.3). Temperature gradually decreased each week and by the 12<sup>th</sup> week was 27.8 (25.4, 30.2).

### Residual activity

Cone bioassays conducted each month with the *An. gambiae* (Kisumu) susceptible strain on nets treated with the mixture or alphacypermethrin induced 87–100% mortality ( $N = 50$  per test) throughout the 3 months. Bioassays on nets treated with chlorfenapyr killed 77% initially and 65% by the end of the trial.

### Species characterization and resistance status

Only *An. gambiae* s.s. M form was found in the trial area. In WHO susceptibility tests with pyrethroid test papers percentage mortality was 20% for *An. gambiae* and 17% for *Cx. quinquefasciatus*. In molecular assays on *An. gambiae* the frequency of *kdr* was 0.86. No molecular assays were conducted on contemporary samples of *Cx. quinquefasciatus* but in an earlier characterization in the area the frequency of *kdr* was 0.63 [19].

**Table 1.** Experimental hut trial results against pyrethroid resistant *Anopheles gambiae*.

	Untreated net	Chlorfenapyr 200 mg/m <sup>2</sup>	Alphacypermethrin 25 mg/m <sup>2</sup>	Mixture 1 (Chlorfenapyr 100+Alpha 25)	Mixture 2 (Chlorfenapyr 200+Alpha 25)
Total females caught	102 <sup>a</sup>	76 <sup>b</sup>	81 <sup>b</sup>	80 <sup>b</sup>	66 <sup>b</sup>
Deterrence %	-	26	21	22	35
Total females in verandah trap	36	31	56	59	50
Exiting % (95% CI)	35 (26–45) <sup>a</sup>	41 (30–52) <sup>a</sup>	69 (59–79) <sup>b</sup>	74 (64–83) <sup>b</sup>	76 (65–86) <sup>b</sup>
Total females blood fed	77	59	47	29	32
Blood-feeding % (95% CI)	75 (66–83) <sup>a</sup>	78 (68–87) <sup>a</sup>	58 (47–69) <sup>b</sup>	36 (26–47) <sup>c</sup>	49 (36–61) <sup>bc</sup>
Personal protection %	-	23	39	62	58

Mixture 1 and 2 applied alphacypermethrin at 25 mg/m<sup>2</sup> and chlorfenapyr at 100 and 200 mg/m<sup>2</sup> respectively. Numbers in the same row sharing a letter superscript do not differ significantly ( $P > 0.05$ ).

doi:10.1371/journal.pone.0087710.t001

**Table 2.** Experimental hut trial results against pyrethroid resistant *Culex quinquefasciatus*.

	Untreated net	Chlorfenapyr 200 mg/m <sup>2</sup>	Alphacypermethrin 25 mg/m <sup>2</sup>	Mixture 1 (Chlorfenapyr 100+Alpha 25)	Mixture 2 (Chlorfenapyr 200+Alpha 25)
Total females caught	811 <sup>a</sup>	527 <sup>b</sup>	567 <sup>b</sup>	487 <sup>b</sup>	492 <sup>b</sup>
Deterrence %	-	35	30	40	39
Total females in verandah trap	227	279	408	370	354
Exiting % (95% CI)	28 (25–31) <sup>a</sup>	53 (49–57) <sup>b</sup>	72 (68–75) <sup>c</sup>	76 (72–79) <sup>c</sup>	72 (68–76) <sup>c</sup>
Total females blood fed	430	200	57	44	59
Blood-feeding % (95% CI)	53 (49–56) <sup>a</sup>	38(34–42) <sup>b</sup>	10 (7–12) <sup>c</sup>	9 (7–12) <sup>c</sup>	12 (7–15) <sup>c</sup>
Personal protection %	-	53	87	90	86

Mixture 1 and 2 applied alphacypermethrin at 25 mg/m<sup>2</sup> and chlorfenapyr at 100 and 200 mg/m<sup>2</sup> respectively. Numbers in the same row sharing a letter superscript do not differ significantly ( $P>0.05$ ).

doi:10.1371/journal.pone.0087710.t002

## Discussion


The aim of the study was to determine whether ITNs treated with a mixture of alphacypermethrin and chlorfenapyr have the potential to provide individual protection against mosquito biting and control of mosquitoes in regions of West Africa where the development of pyrethroid resistance in *An. gambiae* is undermining the effectiveness of pyrethroid treated nets and threatening malaria control. In experimental huts nets treated with the insecticide mixture reduced survival of *An. gambiae* by 75% and prevented host-vector contact by 40–50%, an effect similar to pyrethroid treated nets and LLINs in areas that have not yet developed pyrethroid resistance [5,26]. The loss of efficacy of pyrethroid treated nets when encountering with pyrethroid resistance at the level that occurs in Southern Benin [6,27] was therefore restored by the mixture.

By combining a pyrethroid and a pyrrole on the same net, it was possible to benefit from the properties of each insecticide: the protective (excito-repellent) effect of the pyrethroid and the killing effect of the pyrrole against pyrethroid resistant Anopheline and Culicine mosquitoes. The low rates of mosquito mortality and blood feeding inhibition shown by the alphacypermethrin ITN being compared were typical of pyrethroid ITN trials in this area

against *An. gambiae* and *Cx. quinquefasciatus* [5,27]. Transmission control through mosquito mortality and personal protection from mosquito biting are important attributes of any vector control tool as these work together to reduce vectorial capacity. By both procuring individual protection and killing mosquitoes the chlorfenapyr-alphacypermethrin mixture shows greater potential for malaria control in areas with high level pyrethroid resistance than could be achieved by pyrethroid LLINs.


Chlorfenapyr toxicity to mosquitoes in bioassay is positively temperature dependent (N'Guessan and Rowland, unpublished data). At the ambient temperatures recorded outdoors during the trial the lowest minimum daily temperature was 25.4 and the highest minimum daily temperature was 28.7 degrees. Chlorfenapyr was very effective within this range.

Previously, mixtures of a carbamate (carbosulfan) and a pyrethroid were evaluated on mosquito nets but were not taken forward due to mammalian toxicity issues associated with the carbamate [28]. Chlorfenapyr has a non neurological mode of action and would be a more appropriate companion insecticide due to its safety [24], effectiveness against mosquitoes, and absence of cross resistance to any existing class of insecticide [13,29]. Reducing the concentration of chlorfenapyr by half (to 100 mg/



**Figure 1.** Blood-feeding inhibition rates of *Anopheles gambiae* and *Culex quinquefasciatus* in experimental huts.

doi:10.1371/journal.pone.0087710.g001



**Figure 2. Control corrected mortality of *Anopheles gambiae* and *Culex quinquefasciatus* in experimental huts.**  
doi:10.1371/journal.pone.0087710.g002

m<sup>2</sup>) did not reduce the effectiveness of the mixture, and has obvious cost benefits.

Quite apart from the restoration of control which mixtures promise, empirical research and modelling predict that mixtures are the most efficient tactic for managing insecticide resistance [7]. The attributes required of mixtures in order to prevent the selection of resistance alleles are rigorous, and include the maintenance of effective concentrations of both insecticide components over the lifetime of the net [10–11]. This is the challenge which the next generation of bi-treated LLINs and the formulators of mixtures on nets must meet. In any event, insecticide mixtures for malaria control are a modern day reality. LLIN coverage is going universal and IRS with non-pyrethroid insecticides is being applied concurrently with LLINs as malaria control policy in many areas of high malaria transmission [30–31]. Chlorfenapyr is already showing potential as an IRS treatment in combination with pyrethroid LLIN [16]. A long lasting mixture of chlorfenapyr and alphacypermethrin on nets, if realised, will make an essential contribution to the next generation of LLINs and to prevention of malaria.

## References

- WHO (2012b) World Malaria Report 2012. Geneva: WHO Global Malaria Programme. 38p.
- Lim S, Fullman N, Stokes A, Ravishankar N, Masiye F, et al. (2011) Net Benefits: A Multicountry Analysis of Observational Data Examining Associations between Insecticide-Treated Mosquito Nets and Health Outcomes. *Plos med* 8:9.
- Zaim M, Aitio A, Nakashima N (2000) Safety of pyrethroid-treated nets. *Med. Vet. Entomol* 14: 1–5.
- Ranson H, N'Guessan R, Lines J, Moiroux N, Nkuni Z, et al. (2011) Pyrethroid resistance in African anopheline mosquitoes: what are the implications for malaria control?. *Trends Parasitol* 27: 91–98.
- N'Guessan R, Corbel V, Akogbeto M, Rowland M (2007a) Reduced efficacy of insecticide treated nets and indoor residual spraying for malaria control in pyrethroid resistance area, Benin. *Emerg Infect Dis* 13: 199–206.
- Asidi A, N'Guessan R, Akogbeto M, Curtis C, Rowland M, et al. (2012) Loss of household protection from use of insecticide-treated nets against pyrethroid-resistant mosquitoes, Benin. *Emerg Infect Dis* 18:1101–1106.
- WHO (2012) Global Plan for Insecticide Resistance Management in vectors. WHO Global Malaria Programme. Roll Back Malaria. ISBN 978 92 4 156447 2
- Asidi A N, N'Guessan R, Hutchinson R A, Traore-Lamizana M, Carnevale P, et al. (2004) Experimental hut comparisons of nets treated with carbamate or pyrethroid insecticides, washed or unwashed, against pyrethroid-resistant mosquitoes. *Med Vet Entomol*, 18: 134–140
- Malima R C, Oxborough R M, Tungu P K, Maxwell C, Lyimo I, et al. (2009) Behavioural and insecticidal effects of organophosphate-, carbamate- and pyrethroid-treated mosquito nets against African malaria vectors. *Med Vet Entomol* 23: 317–325
- Curtis CF, Otoo LN (1986). A simple model of the build up of resistance to mixtures of anti malaria drugs. *Trans R Soc Trop Med Hyg* 80:889–892
- Denholm I, Rowland MW (1992) Tactics for managing pesticide resistance in arthropods: theory and practice. *Annual Review of Entomology* 37: 91–112.
- Lovell JB, Wright DE, Gard IE, Miller TE, Treacy ME, et al. (1990) An insecticide/acaricide from a novel class of chemistry. *Brighton Crop Conf* 2/3, 37–42.
- N'Guessan R, Boko P, Odjo A, Akogbeto M, Yates A, et al. (2007b) Chlorfenapyr: a pyrrole insecticide for the control of pyrethroid or DDT resistant *Anopheles gambiae* (Diptera: Culicidae) mosquitoes. *Acta Trop* 102:69–78
- Mosha FW, Lyimo IN, Oxborough RM, Malima R, Tenu F, et al. (2008) Experimental hut evaluation of the pyrrole insecticide chlorfenapyr on bed nets for the control of *Anopheles arabiensis* and *Culex quinquefasciatus*. *Trop Med Int Health* 13: 644–652
- N'Guessan R, Boko P, Odjo A, Knols B, Akogbeto M, et al. (2009) Control of pyrethroid-resistant *Anopheles gambiae* and *Culex quinquefasciatus* mosquitoes with chlorfenapyr in Benin. *Trop Med Int Health* 14: 389–395

## Supporting Information

**Table S1** Checklist of information when reporting an experimental hut trial. (DOC)

## Acknowledgments

The authors thank Drs Susanne Stutz, Wolfgang Paulus (BASF Corp., Ludwigshafen, Germany), James Austin (BASF Corp., NC, USA) and Robert Farlow for supplying samples and for technical input; Robert Sloss for technical programme support (IVCC, Liverpool, UK). We are grateful to E. Vigninou and A. Founnikin for field and laboratory assistance. LSHTM and KCMC are members of the Pan African Malaria Vector Research Consortium (<http://www.pamvec.or.tz>). RN, CN and MR are members of the Malaria Centre of the London School of Hygiene & Tropical Medicine (<http://malaria.lshtm.ac.uk>).

## Author Contributions

Conceived and designed the experiments: MR RN. Performed the experiments: AAK PB AO. Analyzed the data: RN. Contributed reagents/materials/analysis tools: DM. Wrote the paper: RN CN AAK MR. Revised the manuscript: DM.

16. Ngufor C, N'Guessan R, Boko P, Odjo A, Vigninou E, et al. (2011) Combining indoor residual spraying with chlorfenapyr and long-lasting insecticidal bed nets for improved control of pyrethroid-resistant *Anopheles gambiae*: an experimental hut trial in Benin. *Malar J* 10: 343
17. Oxborough R, Kitau J, Matowo J, Feston E, Mndeme R, et al. (2013). ITN mixtures of chlorfenapyr (pyrrole) and alphacypermethrin (pyrethroid) for improved control of pyrethroid resistant *Anopheles arabiensis* and *Culex quinquefasciatus*. *PLoS ONE* 8: e55781.
18. Djouaka RF, Bakare AA, Coulibaly ON, Akogbeto MC, Ranson H, et al. (2008) Expression of the cytochrome P450s, CYP6P3 and CYP6M2 are significantly elevated in multiple pyrethroid resistant populations of *Anopheles gambiae* s.s. from Southern Benin and Nigeria. *BMC Genomics* 9:53
19. Corbel V, N'Guessan R, Brengues C, Chandre F, Djogbenou L, et al. (2007) Multiple insecticide resistance mechanisms in *Anopheles gambiae* and *Culex quinquefasciatus* from Benin, West Africa. *Acta Trop* 101: 207–216
20. WHO (2006) Guidelines for testing mosquito adulticides for indoor residual spraying and treatment of mosquito nets. Geneva: World Health Organization; WHO/CDS/NTD/WHOPES/GCDPP/2006.3.
21. Scott JA, Brogdon WG, Collins FH, (1993) Identification of single specimens of the *Anopheles gambiae* complex by the polymerase chain reaction. *Am J Trop Med Hyg* 49: 520–529.
22. Favia G, Della Torre A, Bagayoko M, Lanfrancotti T, Sagnon NF, et al. (1997) Molecular identification of sympatric chromosomal forms of *Anopheles gambiae* and further evidence of their reproductive isolation. *Insect Mol Biol* 6: 377–383.
23. Martínez-Torres D, Chandre F, Williamson MS, Darriet F, Berge JB, et al. (1998) Molecular characterization of pyrethroid knockdown resistance (kdr) in the major malaria vector *Anopheles gambiae* s.s. *Insect Mol Biol* 7: 179–184.
24. Tomlin C (2000) *The Pesticide Manual, A World Compendium*, 12th edn. British Crop Protection Council, London.
25. WHO (2004) A Generic Risk Assessment Model for Insecticide Treatment and Subsequent Use Of Mosquito Nets. WHO/CDS/WHOPES/GCDPP/2004.6/WHO/PCS/04.1. WHO, Geneva.
26. Tungu P, Magesa S, Maxwell C, Malima R, Masue D, et al. (2010) Evaluation of PermaNet3.0 a deltamethrin-PBO combination net against *Anopheles gambiae* and pyrethroid resistant *Culex quinquefasciatus* mosquitoes: an experimental hut trial in Tanzania. *Malar J* 9:21
27. N'Guessan R, Asidi A, Boko P, Odjo A, Akogbeto M, et al. (2010) An experimental hut evaluation of PermaNet 3.0, a deltamethrin-piperonyl butoxide combination net, against pyrethroid-resistant *Anopheles gambiae* and *Culex quinquefasciatus* mosquitoes in southern Benin. *Trans R Soc Trop Med Hyg* 104: 758–765
28. Hougard JM, Corbel V, N'Guessan R, Darriet F, Chandre F, et al. (2003) Efficacy of mosquito nets treated with insecticide mixtures or mosaics against insecticide resistant *Anopheles gambiae* and *Culex quinquefasciatus* (Diptera: Culicidae) in Côte d'Ivoire. *Bull Entomol Res* 93:491–498
29. Oliver S, Kaiser M, Wood O, Coetzee M, Rowland M, et al. (2010) Evaluation of the pyrrole insecticide chlorfenapyr against pyrethroid resistant and susceptible *Anopheles funestus* (Diptera: Culicidae). *Tropical Medicine and International Health*, 15: 127–131.
30. Corbel V, Akogbeto M, Damien G, Djenontin A, Chandre F, et al. (2012) Combination of malaria vector control interventions in pyrethroid resistance area in Benin: a cluster randomised controlled trial. *Lancet Infect Dis* 12: 617–26.
31. West P, Protopopoff N, Wright A, Kivaju Z, Tigererwa R, et al. (2013) Added Protection against malaria due to indoor residual house spraying in combination with insecticide treated nets versus insecticide treated nets alone: results of a cluster randomised trial. *Lancet Infect Dis* in press.