

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Prendiville, Walter J P; Elbourne, Diana; McDonald, Susan J (2009)
WITHDRAWN: Active versus expectant management in the third
stage of labour. Cochrane database of systematic reviews (Online)
(3). CD000007. ISSN 1469-493X

Downloaded from: <http://researchonline.lshtm.ac.uk/19906/>

DOI:

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: Copyright the author(s)

Active versus expectant management in the third stage of labour (Review)

Prendiville WJP, Elbourne D, McDonald SJ



**THE COCHRANE
COLLABORATION[®]**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2009, Issue 3

<http://www.thecochranelibrary.com>

WILEY

TABLE OF CONTENTS

HEADER	1
REASON FOR WITHDRAWAL	1
WHAT'S NEW	1
HISTORY	1

[Intervention Review]

Active versus expectant management in the third stage of labour

Walter JP Prendiville¹, Diana Elbourne², Susan J McDonald³

¹Department of Obstetrics and Gynaecology, Coombe Lying-In Hospital, Dublin 8, Ireland. ²Medical Statistics Unit, London School of Hygiene and Tropical Medicine, London, UK. ³Midwifery Professorial Unit, La Trobe University/Mercy Hospital for Women, Heidelberg, Australia

Contact address: Sonja Henderson, Cochrane Pregnancy and Childbirth Group, School of Reproductive and Developmental Medicine, Division of Perinatal and Reproductive Medicine, The University of Liverpool, First Floor, Liverpool Women's NHS Foundation Trust, Crown Street, Liverpool, L8 7SS, UK. sonjah@liverpool.ac.uk. s.l.henderson@liverpool.ac.uk.

Editorial group: Cochrane Pregnancy and Childbirth Group.

Publication status and date: Withdrawn from publication for reasons stated in the review, published in Issue 3, 2009.

Review content assessed as up-to-date: 8 March 2000.

Citation: Prendiville WJP, Elbourne D, McDonald SJ. Active versus expectant management in the third stage of labour. *Cochrane Database of Systematic Reviews* 2009, Issue 3. Art. No.: CD000007. DOI: 10.1002/14651858.CD000007.pub2.

Copyright © 2009 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

The editorial group responsible for this previously published document have withdrawn it from publication.

REASON FOR WITHDRAWAL

This review was withdrawn from *The Cochrane Library* 2009, Issue 3, because it was out of date. It has been updated by a new review ('Active versus expectant management for women in the third stage of labour') first published in *The Cochrane Library* 2010, Issue 7.

WHAT'S NEW

Last assessed as up-to-date: 8 March 2000.

Date	Event	Description
16 June 2010	Amended	Updated published note - see Published notes

HISTORY

Protocol first published: Issue 1, 1997

Review first published: Issue 1, 1997

Date	Event	Description
4 March 2009	Feedback has been incorporated	We have received feedback from Susan van Wyk on behalf of the Masters Clinical Epidemiology class, Stellenbosch University, about the validity of the findings of the current publication pending the completion of the update, as a result of which we have withdrawn the review from publication
3 March 2009	Amended	Review withdrawn from publication in Issue 3, 2009.
5 August 2008	Amended	A note has been added to explain that this review is currently being updated following a new protocol which has been published in Issue 4, 2008 of <i>The Cochrane Library</i> .
22 January 2008	Amended	Converted to new review format. A note has been added to explain that this review is currently being updated following a new protocol which has been submitted for editorial approval to publish
9 March 2000	New citation required and conclusions have changed	Substantive amendment