

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Mabey, DC; Sollis, KA; Kelly, HA; Benzaken, AS; Bitarakwate, E; Chandalucha, J; Chen, XS; Yin, YP; Garcia, PJ; Strasser, S; +5 more... Chintu, N; Pang, T; Terris-Prestholt, F; Sweeney, S; Peeling, RW; (2012) Point-of-Care Tests to Strengthen Health Systems and Save Newborn Lives: The Case of Syphilis. PLoS medicine, 9 (6). e1001233. ISSN 1549-1277 DOI: <https://doi.org/10.1371/journal.pmed.1001233>

Downloaded from: <http://researchonline.lshtm.ac.uk/21098/>

DOI: <https://doi.org/10.1371/journal.pmed.1001233>

Usage Guidelines:

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

<https://researchonline.lshtm.ac.uk>

Point-of-Care Tests to Strengthen Health Systems and Save Newborn Lives: The Case of Syphilis

David C. Mabey¹, Kimberly A. Sollis¹, Helen A. Kelly¹, Adele S. Benzaken², Edward Bitarakwate³, John Changalucha⁴, Xiang-Sheng Chen⁵, Yue-Ping Yin⁵, Patricia J. Garcia⁶, Susan Strasser⁷, Namwinga Chintu⁸, Tikki Pang⁹, Fern Terris-Prestholt¹, Sedona Sweeney¹, Rosanna W. Peeling^{1*}

1 London School of Hygiene and Tropical Medicine, London, United Kingdom, **2** Alfredo da Matta Foundation, Manaus, Brazil, **3** Elizabeth Glaser Pediatric AIDS Foundation, Kampala, Uganda, **4** National Institute of Medical Research, Mwanza, Tanzania, **5** The National Center for Sexually Transmitted Diseases, Center for Disease Control, Nanjing, China, **6** The Universidad Peruana Cayetano Heredia, Lima, Peru, **7** Elizabeth Glaser Pediatric AIDS Foundation, Lusaka, Zambia, **8** Centre for Infectious Disease Research in Zambia, Lusaka, Zambia, **9** Research Policy & Cooperation (RPC/IER), World Health Organization, Geneva, Switzerland

The Challenge

Poor rural populations in low- and middle-income countries (LMICs) do not have access to diagnostic laboratories. Until recently, this has meant that they could not be screened for asymptomatic infections such as HIV or syphilis, and could only be treated syndromically when presenting with symptoms such as fever. This situation has now changed. Simple, rapid, and affordable point-of-care tests (POCTs), which do not require electricity, a laboratory, or highly trained staff, are now available and widely used for several common infections in LMICs [1]. POCTs offer an unprecedented opportunity to reduce inequalities in health, and to help LMICs achieve the health-related Millennium Development Goals (MDGs) [2].

Syphilis in pregnancy causes more than half a million stillbirths or neonatal deaths annually [3,4]. These deaths could be prevented if all pregnant women were screened for syphilis and treated with a single dose of penicillin before the third trimester [5]. Screening of pregnant women for syphilis is recommended policy in almost every country, and even in LMICs the vast majority of women attend antenatal clinic (ANC) at least once during pregnancy; yet in sub-Saharan Africa, it has been estimated that fewer than 40% of those who attend ANC are screened for syphilis [4]. Why is this? The rapid plasma reagin (RPR) test, which is used for screening in many countries, is only available in laboratories, as reagents require refrigeration and the reactions are often batched. Blood is often sent to a central laboratory, or the tests are

performed at the end of the day, requiring patients to return for their results. In one study in Nairobi, fewer than 10% of women who tested positive received treatment [6].

New POCTs for syphilis are now available and some of them meet the ASSURED criteria, being Affordable, Sensitive, Specific, User-friendly, Rapid and robust, Equipment-free, and Deliverable to those who need them [7]. However, for POCTs to maximize their potential, a number of challenges must be addressed. Health care workers (HCWs) must be properly trained in how to use them and how to interpret their results; the quality of the tests and the testing must be maintained through a quality assurance programme; and supply chains must be improved to ensure that POCTs and effective treatment are continuously available. Here we describe our experiences in attempting to overcome these challenges and introduce syphilis POCTs to prevent adverse outcomes of pregnancy in six LMICs. Our goal was to determine the

feasibility of introducing POCTs into different settings in countries with different health systems and cultural and socioeconomic contexts. We measured several health systems indicators to determine whether we met the success criteria as shown in Table 1.

The Project

Project sites were located in countries where national guidelines recommend antenatal screening for syphilis. We convened a workshop with all the principal investigators (PIs) at the start of this project to develop a set of guiding principles and a generic protocol for all countries to use as a template.

Policy and Stakeholder Consultations

Each country project team consulted with national and local health authorities and other stakeholders on the design and location of the project, and sought their endorsement of the strategy of same-day

Citation: Mabey DC, Sollis KA, Kelly HA, Benzaken AS, Bitarakwate E, et al. (2012) Point-of-Care Tests to Strengthen Health Systems and Save Newborn Lives: The Case of Syphilis. *PLoS Med* 9(6): e1001233. doi:10.1371/journal.pmed.1001233

Published: June 12, 2012

Copyright: © 2012 Mabey et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Financial support for this study was provided by the Bill & Melinda Gates Foundation through a grant (#47697) to the corresponding author at the UNICEF/UNDP/World Bank/WHO Special Programme for Research and Training for Tropical Diseases and subsequently at the London School of Hygiene and Tropical Medicine. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abbreviations: ANC, antenatal clinic; ARV, anti-retroviral; HCW, health care workers; LMIC, low- and middle-income country; MCH, maternal and child health; MDG, Millennium Development Goal; MOH, Ministry of Health; PI, principal investigator; PMTCT, prevention of mother-to-child transmission; POCT, point-of-care test; RPR, rapid plasma reagin; STAT, same-day testing and treatment; STI, sexually transmitted infection; WHO, World Health Organization.

* E-mail: Rosanna.Peeling@lshtm.ac.uk

Provenance: Not commissioned; externally peer reviewed.

The Health in Action section is a forum for individuals or organizations to highlight their innovative approaches to a particular health problem.

Summary Points

- The widespread adoption and implementation of new diagnostic technologies in developing countries is a major challenge.
- We used a non-traditional approach to implementation research, engaging policy makers in each country in the design of a prenatal syphilis screening project using point-of-care tests and ensuring relevance to the local health care system.
- As a result of this study, all six countries changed policy to adopt point-of-care syphilis testing into their prenatal screening programmes.
- We showed that it is possible to implement a quality assurance and quality management system for point-of-care testing in all settings.
- Lessons learnt and policies implemented as a result of the project have strengthened health systems by improving access to quality-assured prenatal screening and saving newborn lives.

testing and treatment (STAT) for syphilis. All site protocols were approved by the relevant institutional review boards (IRBs) and the World Health Organization (WHO) Ethics Review Committee.

Project Settings and Study Populations

Syphilis POCTs were introduced in rural antenatal clinics in Tanzania, Uganda, and China; and in both rural and urban clinics in Peru and Zambia. In Brazil, community-based screening was introduced in remote indigenous populations where there were no fixed health facilities or laboratories, and no screening was previously available. The opportunity was taken to screen all sexually active individuals (Table 2).

Country-specific testing and treatment algorithms and the test used for the project were selected by each country in consultation with the Ministry of Health (MOH). Brazil, Peru, Tanzania, Uganda, and Zambia used the SD Biotec Rapid Syphilis test (Standard Diagnostics, Seoul, Korea), which we had previously evaluated and found to have acceptable performance [7]

and was available through the WHO Bulk Procurement Programme at negotiated pricing. In China the Rapid Syphilis test (Wantai Ltd, Beijing, China) was used.

Baseline Survey

In each country, published data and MOH reports on the prevalence of syphilis and the coverage of screening and treatment in ANCs were reviewed. Except in Brazil, where no testing had previously been available for remote indigenous populations, a baseline survey was undertaken, at which project sites were visited, HCWs were interviewed concerning current practice for syphilis screening and barriers to achieving full coverage, and recent records on the number and proportion of women screened for syphilis were reviewed. Where coverage was sub-optimal, reasons for this were elicited from HCWs and their supervisors.

Site Preparation: Training, Quality Assurance, and Supply Chain Management

HCWs were trained in the use of syphilis POCTs, using training materials

developed for the project, translated into the local language, and adapted according to the local cultural and social context. Courses for trainers and supervisors included practical training on how to perform the tests and interpret the results, and how to train HCWs and assess their competence, with additional training in stock management, record keeping, and quality control.

For quality assurance, supervisors were provided with proficiency panels prepared by a reference laboratory, consisting of tubes in which sera characterized as positive or negative had been allowed to dry at room temperature. Dried sera were reconstituted with buffered saline on site and used to assess the performance of POCTs by HCWs. This simple, low-cost method was originally developed to assure the quality of HIV testing [8]. Syphilis POCTs were provided to health facilities through the normal supply chain to allow the PIs to monitor supply chain problems and provide sustainable solutions in case of stock-outs. All individuals who tested positive at each site were treated with the first dose of benzathine penicillin on the same day.

Continuous Quality Improvement, Outcome Measures, and Success Criteria

An internal monitor was appointed in each project team to monitor quality and outcome measures continuously so that problems were brought to the attention of the PIs and regional medical officers, and solutions were tried until a sustainable solution was found. External monitors visited the sites and provided suggestions for further improvement. The opinions of HCWs on the ease of use and acceptability of the POCTs were sought through interviews and focus group discussions. The results of the baseline survey and

Table 1. Health systems indicators and proposed success criteria.

| Health System Indicators | Outcome Measures | Success Criteria |
|---|---|---|
| Coverage (increase in access to screening) | # women pregnant screened pre- and post-POCT introduction | 90% coverage post-POCT introduction |
| Improved health outcome | # infected women treated | 90% of women who tested positive treated |
| Acceptability by clients and health workers | Job satisfaction; client satisfaction with service | Increased job satisfaction and client satisfaction with services |
| Quality assurance | # workers who passed proficiency test | 90% of CHWs passed the proficiency test |
| Integration into existing programmes | Effect on ANC and prevention of mother-to-child transmission (PMTCT) programmes for HIV | Synergy with existing programmes |
| Sustainability | # countries which change policy | Policy change, development of national guidelines for use, and plans for scale up |

doi:10.1371/journal.pmed.1001233.t001

Table 2. Key indicators for syphilis screening before and after POCT introduction.

| Country and Population Screened | Screening Method Pre-POCT | % People Screened Pre-POCT | % People Screened Post-POCT | % Positive Post-POCT | % People Treated Post-POCT |
|--|---------------------------------|----------------------------|-----------------------------|---|-------------------------------|
| Amazonas region, Brazil (sexually active population) | No screening available | No screening available | 45,971/84,038 (54.5%) | Sexually active: 745/45,971 (1.62%); pregnant women ^a : 64/4,695 (1.36%) | 808/808 (100%) |
| Rural ANC in Guangdong province, China | TRUST (RPR) confirmed with TPPA | N/a ^b | 5,272/5,489 (96.0%) | 109/5,272 (1.9%) | 102/109 (93.6%) |
| Maternity hospital in Lima and ANCs in Callao, Peru | RPR | 9,595/18,757 (51%) | 15,985/16,839 (95%) | 146/15,985 (1.0%) | 128/134 ^c (97%) |
| District Hospital and 51 health facilities in Geita District, Tanzania | RPR | 634/3,561 (17.8%) | 58,249/58,249 (100%) | 6,345/58,249 (10.9%) | 5,717/6,345 (90.1%) |
| Kampala Hospital and rural ANCs in Uganda | RPR | 140/8,475 (1.7%) | 13,131/14,540 (90.3%) | 690/13,131 (5.3%) | 715/690 (103.6%) ^d |
| Lusaka Hospital and rural ANCs in Mongu district, Zambia | RPR | 12,761/15,967 (79.9%) | 11,460/11,985 (95.6%) | 1,050/11,460 (9.2%) | 1,000/1,050 (95.2%) |

^aPregnant women were a subset of the sexually active population.

^bN/a: No data available before the introduction of POCTs as testing was done by a laboratory and these services were not widely available in rural China.

^cTwelve POCT-positive pregnant women had a previous history of syphilis and were either RPR negative or had received treatment recently. The treating physicians did not think additional treatment was required.

^dANC attenders whose accompanying partners were found to have syphilis were offered presumptive treatment even if their own result was negative. This resulted in more women being treated than had positive tests.

RPR, rapid plasma reagin test; TPPA, *Treponema pallidum* particle agglutination assay.

doi:10.1371/journal.pmed.1001233.t002

interim data were shared with national and regional policy makers and programme managers through dissemination workshops at each site. Data on coverage of screening and treatment and on HCWs' experiences were collected using standard data collection instruments, double entered, and analysed in country. The results from each country are being analysed in more detail and will be submitted for publication separately.

Results

Impact of POCTs on Coverage of Syphilis Screening and Treatment

Over 100,000 pregnant women were screened for syphilis. The introduction of POCTs increased the proportion of ANC attenders screened for syphilis to >90%, and the proportion of pregnant women with syphilis who were treated the same day exceeded 90% in all countries (Table 2). In Brazil, HCWs in remote communities succeeded in screening 55% of the sexually active population (defined as ≥ 10 years of age) for syphilis, exceeding the 30%–40% target originally set.

Acceptability of POC Syphilis Tests among HCWs and Study Populations

POCTs were well accepted, and considered easy to perform, by HCWs at all

sites. For example, in Uganda, 82% of HCWs described the POCTs as “very easy to perform.” At sites where RPR had been previously performed, HCWs reported that they could test more patients per day with the POCT. Clients at all sites appreciated receiving results and treatment at their first visit, rather than being asked to return for their results as they had in the past. Most clients at all sites preferred a finger prick to venepuncture because of the smaller volume of blood required.

Laboratory staff in the main maternity hospital in Peru were initially opposed to the idea of nurses and midwives doing a task that had traditionally been done by laboratory technicians, and the hospital management did not believe it was necessary to introduce POCTs. Although coverage of screening and treatment for syphilis at the ANC was high before the introduction of POCTs, women did not receive test results for at least 15 days [9]. Moreover, a significant number of women who delivered in the hospital had not attended ANC and, among women admitted with early or mid-term abortions, few were screened. When shown these data, the hospital director welcomed the introduction of POCTs. Laboratory staff trained and supervised the nurses and midwives doing the tests, and set up a

quality assurance programme using a proficiency panel. This programme was enthusiastically approved by both laboratory staff and nurses, each of whom received a certificate if they performed well on the proficiency panel.

The indigenous population in Brazil initially had reservations about the POCT because of the possibility of a positive test result, the pain caused by the finger prick, and concerns about the treatment. In rural areas of China, pregnant women of ethnic minorities were less likely to accept the test due to the low perception of risk. This gradually changed as the programme became more established and expanded into the community.

Experience with Quality Assurance

The proficiency panel using dried tube specimens allowed the study team to identify HCWs in need of refresher training. Figure 1 shows the results obtained in health facilities in Geita District in Tanzania over an 18-month period, using a panel consisting of six tubes containing a mixture of positive and negative samples. Six months after the initial training workshop, the percentage of facilities failing the proficiency panel rose to 65%. In several of these clinics, the staff member performing the tests was not the ones who had been trained to do so. In




Figure 1. Percentage of facilities receiving a passing* or failing score on each of the seven proficiency panels. *Pass defined as $\geq 67\%$ score on the proficiency panel. doi:10.1371/journal.pmed.1001233.g001

response to the poor performance across the district, the laboratory coordinator arranged for one staff member from each of the failing health facilities to attend a refresher training workshop. HCWs were instructed to re-train colleagues upon return to their facility. Following the refresher training, proficiency panel scores improved greatly.

Integration into PMTCT Services for HIV

In Uganda and Zambia, we assessed the impact of introducing syphilis POCTs on existing prevention of mother-to-child transmission (PMTCT) programmes for HIV (Table 3). After POCT introduction, there was a significant increase in ANC attendance, resulting in more pregnant women being tested for HIV in both countries. In Zambia, there was also a significant increase in the number of women receiving anti-retroviral (ARV) prophylaxis and referred to care and treatment following the introduction of syphilis POCTs. We believe that increased awareness of the importance of syphilis and HIV in pregnancy, better supportive supervision, and greater efficiency of integrated services all contributed to this improvement.

Impact on Health Policy

This project has brought to the forefront the power and value of implementation research in bringing about rapid policy change and strengthening health systems. It also has broader implications for health systems strengthening, which is fundamental to the achievement of the

health-related MDGs. The inclusion of policy makers from the beginning of the project, and keeping them informed during POCT implementation, led to policy change even before the research was completed. In all six countries, regular meetings were held between project staff and health authority officials at national, regional, and district levels, focussing on those responsible for both maternal and child health (MCH) and HIV/AIDS programmes. This is important, as previous studies have found that it is often not clear whether antenatal syphilis screening is the responsibility of the MCH, HIV/AIDS, or sexually transmitted infection (STI) control programmes [4,10].

All six countries changed their policy to recommend the use of rapid tests to provide a STAT service. National guidelines have been amended accordingly. Meetings jointly sponsored by our programme and the WHO have been held with policy makers from many countries in Asia, Africa, and South America, at which we shared the toolkit developed for test introduction and presented project results. This has resulted in many more countries in all three regions adopting and rolling out the use of POCTs for syphilis screening.

In Brazil, as a result of this study, the minister of health announced in 2011 that syphilis screening for remote populations would be one of the three main priorities for the national programme for the control of HIV and STIs, and the president announced the “Stork” Initiative to bring healthy babies free of both HIV and syphilis. In China, Premier Wen Jiabao chaired a meeting in November 2010 at

which he announced that a screening policy for syphilis would be introduced in ANCs in China [11]. The WHO has now included the use of POCTs for syphilis in its STI Management Guidelines. Our project leaders have worked with their national MOH to develop national guidelines for the use of POCTs in prenatal screening, and are now involved in scaling up training and quality assurance for neighbouring countries as well.

Discussion

This project has shown that POCTs for syphilis can be effectively introduced in a range of settings, from cities in China and Peru, to remote villages in East Africa, and even more remote indigenous populations in the Amazon rain forest. By working with the existing health care system to integrate testing, the introduction of POCTs resulted in large numbers of women being tested and treated for syphilis, averting many stillbirths and reducing neonatal mortality. Importantly, the outcomes resulted in policy change in all project countries to incorporate POCTs into national guidelines. The project has highlighted the drivers for success as well as the key challenges that need to be overcome to scale up the use of POCTs. Their introduction could have a major impact in reducing inequalities in health care and motivating HCWs, who are happy to be able to provide a diagnostic service to their patients that results in immediate treatment. Unlike the financial motivation provided by many vertical programmes, this type of motivation through increased job satisfaction has

Table 3. Effect of introducing POCTs for syphilis on PMTCT of HIV programmes in Uganda and Zambia.

| Outcome Measure | Uganda | Zambia |
|---|------------------|------------------|
| HIV testing pre-POCT (%) | 95.6% (n=6,479) | 95.5% (n=7,479) |
| HIV testing post-POCT (%) | 96.4% (n=11,192) | 97.7% (n=11,414) |
| p-value | p=0.009 | p<0.0001 |
| ARV prophylaxis pre-POCT (%) | 78.5% (n=570) | 98.3% (n=1,303) |
| ARV prophylaxis post-POCT (%) | 83.6% (n=964) | 100.0% (n=2,036) |
| p-value | p=0.007 | p<0.0001 |
| Refer to care & treatment pre-POCT (%) | 16.9% (n=85) | 73.7% (n=977) |
| Refer to care & treatment post-POCT (%) | 16.1% (n=118) | 84.6% (n=1,721) |

doi:10.1371/journal.pmed.1001233.t003

no adverse impact on other services and is more sustainable.

The introduction of POCTs can serve as a “probe” for the efficient functioning of a health system more broadly, as the intention to introduce POCTs (a “technology”) for syphilis had “knock-on” effects on other important “building blocks” of the health system including service delivery, the health workforce, health information, health financing, and leadership and governance [12]. It represents a real and powerful example of the need for “systems thinking” [13], an approach to problem solving that views “problems” as part of a wider, dynamic system and which demands a deeper understanding of the linkages, relationships, interactions, and behaviours among the elements that characterize the entire system.

This project has highlighted the importance of quality assurance programmes for POC testing. Without them, failures in HCW proficiency would have gone unnoticed. It is deeply concerning that most HIV testing programmes in developing countries now rely on POCTs, but have no quality assurance programme to ensure that the tests are properly performed [14]. The continuing use of proficiency panels to monitor the quality of testing is particularly important where there is a rapid turnover of staff.

In spite of the success of this project in increasing access to syphilis screening in many countries, a number of challenges remain. POCTs cost a little more than the RPR test, but are available through the WHO bulk procurement programme for less than US\$1. The cost per woman screened ranged from US\$1.9 to \$6.1 in Tanzanian health facilities [15], and screening pregnant women for syphilis remains one of the most cost-effective health interventions [16]. HCWs and their supervisors received training in stock

management, but problems remain at the national level, and stock outs of tests and treatments still occurred. One visit to Geita District at the end of the project found that ten of 17 facilities had recorded at least one day of stock out of POCTs over a 4-month period.

Limitations of This Study

We were not able to show impact, as we were funded only to determine feasibility of POCT introduction for one year. We adopted a non-traditional approach to implementation research since we wanted to determine feasibility in a variety of cultural and socioeconomic settings. We decided not to have a rigid protocol to allow countries to improvise as necessary to maximize POCT implementation based on STAT. However, we did ensure a uniform set of outcome measures were collected across all sites, including quality assurance of POC testing. We used policy change as an indicator of success, but strengthening fragile health systems requires sustained effort of which policy is only a start.

Moving Forward

The project has implications for improved health care delivery beyond the diagnosis and management of syphilis in pregnancy. Every case of congenital syphilis represents a failure of health care delivery; attempts to prevent it could act as a “window of knowledge” or “proxy indicator” for the robustness of health service delivery systems more generally. Lessons learnt about appropriate technologies, supply chain management, training and motivating HCWs, quality assurance, effective teamwork, and open communication are valuable not just for syphilis, but for health care delivery, and can ultimately lead to sustained health systems strengthening. There is even a potential for

“reverse innovation” where improved, cost-effective processes discovered in the developing world are adopted by health care delivery organizations in industrialized countries.

We wrote about the tragedy of babies avoiding HIV and dying of syphilis because of vertical disease control programmes and compartmentalized health care infrastructures [17]. Our long-term vision is to facilitate the development and implementation of an essential POCT package for prenatal care that could be broadened to include diagnostics for both infectious diseases and conditions such as anaemia and pre-eclampsia to ensure safe motherhood and healthy babies. Such an integrated approach could result in improved efficiency, a more robust health care system, and lives saved.

Acknowledgments

This work was supported by the Bill & Melinda Gates Foundation (Grant 47697). We thank Ron Ballard, Ned Hook, King Holmes, Myron Cohen, Maurine Murtagh, Renee Ridzon, Christine Rousseau, and Kathy Wilfert for expert advice, and Izabela Suder and Beth Downe for administrative and organizational assistance throughout the project.

Author Contributions

Conceived and designed the experiments: DCM ASB EB JC XSC YPY PJG SStrasser NC FTP RWP. Performed the experiments: ASB EB JC XSC YPY PJG SStrasser NC FTP. Analyzed the data: DCM KAS HAK ASB EB JC XSC YPY PJG SStrasser NC FTP SSweeney TP RWP. Wrote the first draft of the manuscript: DCM KAS RWP. Contributed to the writing of the manuscript: DCM KAS HAK ASB EB JC XSC YPY PJG SStrasser NC FTP SSweeney TP RWP. ICMJE criteria for authorship read and met: DCM KAS HAK ASB EB JC XSC YPY PJG SStrasser NC FTP SSweeney TP RWP. Agree with manuscript results and conclusions: DCM KAS HAK ASB EB JC XSC YPY PJG SStrasser NC FTP SSweeney TP RWP.

References

1. Peeling RW, Mabey D (2010) Point-of-care tests for diagnosing infections in the developing world. *Clin Microbiol Infect* 16: 1062–1069.
2. Lozano R, Wang H, Foreman KJ, Rajaratnam JK, Naghavi M, et al. (2011) Progress towards Millennium Development Goals 4 and 5 on maternal and child mortality: an updated systematic analysis. *Lancet* 378: 1139–1165.
3. Schmid GP, Stoner BP, Hawkes S, Broutet N (2007) The need and plan for global elimination of congenital syphilis. *Sex Transm Dis* 34(suppl): S5–S10.
4. Gloyd S, Chai S, Mercer MA (2001) Antenatal syphilis in sub-Saharan Africa: missed opportunities for mortality reduction. *Health Policy Plan* 16: 29–34.
5. Watson-Jones D, Gumodoka B, Weiss H, Changalucha J, Todd J, et al. (2002) Syphilis in pregnancy in Tanzania. II. The effectiveness of antenatal syphilis screening and single-dose benzathine penicillin treatment for the prevention of adverse pregnancy outcomes. *J Infect Dis* 186: 948–957.
6. Temmerman M, Mohamedali F, Fransen L (1993) Syphilis prevention in pregnancy: an opportunity to improve reproductive and child health in Kenya. *Health Policy Plan* 8: 122–127.
7. Mabey D, Peeling RW, Ballard R, Benzaken AS, Galbán E, et al. (2006) Prospective, multi-centre clinic-based evaluation of four rapid diagnostic tests for syphilis. *Sex Transm Infect* 82 Suppl 5: S13–S16.
8. Parekh BS, Anyanwu J, Patel H, Downer M, Kalou M, et al. (2010) Dried tube specimens: A simple and cost-effective method for preparation of HIV proficiency testing panels and quality control materials for use in resource-limited settings. *J Virol Methods* 163: 295–300.
9. La Rosa Roca S, Garcia P (2011) Can rapid syphilis tests be implemented and improve screening for maternal syphilis in a third level hospital? *Sex Transm Infect* 87, suppl 1: A211.
10. Watson-Jones D, Oliff M, Terris-Prestholt F, Changalucha J, Gumodoka B, et al. (2005) Antenatal syphilis screening in sub-Saharan Africa – lessons learned from Tanzania. *Trop Med Int Health* 9: 934–943.
11. GOV.cn (29 November 2010) Chinese government vows to step up HIV-AIDS control ahead of World AIDS Day. Available: http://www.gov.cn/english/2010-11/29/content_1755901.htm. Accessed 4 May 2012.
12. WHO (2007) Everybody's business-strengthening health systems to improve health outcomes. Geneva: World Health Organization.
13. Alliance for Health Policy and Systems Research (2009) Systems thinking. Geneva: Alliance for Health Policy and Systems Research.
14. Plate DK on behalf of the Rapid HIV Test Evaluation Working Group (2007) Evaluation and implementation of rapid HIV tests: the experience in 11 African countries. *AIDS Research and Human Retroviruses* 23: 1491–1498.
15. Sweeney S, Mosha J, Terris-Prestholt F, Vickerman P, Changalucha J, et al. (July 2011) Efficiency versus equity in screening: considerations in the scale-up of rapid syphilis testing in rural Tanzania. Abstract O2-S4.01 ISSITDR, July 2011.
16. Terris-Prestholt F, Watson-Jones D, Mugeye K, Kumaranayake L, Ndeki L, et al. (2003) Is antenatal syphilis screening still cost-effective in Sub-Saharan Africa? *Sex Transm Inf* 79: 375–381.
17. Peeling RW, Mabey D, Fitzgerald DW, Watson-Jones D (2004) Avoiding HIV and dying of syphilis. *Lancet* 364: 1561–1563.