

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Hamaguchi, S; Cuong, NC; Tra, DT; Doan, YH; Shimizu, K; Tuan, NQ; Yoshida, LM; Mai, LQ; Duc-Anh, D; Ando, S; Arikawa, J; Parry, CM; Ariyoshi, K; Thuy, PT (2015) Clinical and Epidemiological Characteristics of Scrub Typhus and Murine Typhus Among Hospitalized Patients with Acute Undifferentiated Fever in Northern Vietnam. *The American journal of tropical medicine and hygiene*. ISSN 0002-9637 DOI: <https://doi.org/10.4269/ajtmh.14-0806>

Downloaded from: <http://researchonline.lshtm.ac.uk/2131852/>

DOI: [10.4269/ajtmh.14-0806](https://doi.org/10.4269/ajtmh.14-0806)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

SUPPLEMENTAL TABLE 1

Comparison of basic information between 579 included and 52 excluded patients

	Patients		<i>P</i> value
	Included, <i>N</i> = 579	Excluded, <i>N</i> = 52	
Age (mean, SD)	46.2, 15.7	52.2, 17.6	0.009
Sex	358 (61.8)	36 (69.2)	0.3
Hanoi	241 (41.6)	23 (44.2)	0.7
Rainy season (May–October)	344 (59.4)	36 (69.2)	0.2
High-exposure occupation*	271 (47.1)	23 (45.1)	0.8
β-Lactam antibiotics not effective	189 (77.5)	15 (83.3)	0.8
Rash	217 (37.5)	25 (48.1)	0.1
Lymphadenopathy	215 (37.1)	13 (25.0)	0.08
Eschar	161 (27.8)	9 (17.3)	0.1
Hepatomegaly/splenomegaly	253 (43.7)	22 (42.3)	0.8

SD = standard deviation.

*High-exposure occupation: occupation with frequent contacts with natural environment, such as farming, dairy husbandry, and environmental construction engineering.