

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Elliott, AM; Mawa, PA; Webb, EL; Nampijja, M; Lyadda, N; Bukusuba, J; Kizza, M; Namujju, PB; Nabulime, J; Ndibazza, J; Muwanga, M; Whitworth, JA (2010) Effects of maternal and infant co-infections, and of maternal immunisation, on the infant response to BCG and tetanus immunisation. *Vaccine*, 29 (2). pp. 247-55. ISSN 0264-410X
DOI: <https://doi.org/10.1016/j.vaccine.2010.10.047>

Downloaded from: <http://researchonline.lshtm.ac.uk/2195/>

DOI: [10.1016/j.vaccine.2010.10.047](https://doi.org/10.1016/j.vaccine.2010.10.047)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>



Published as: *Vaccine*. 2010 December 16; 29(2-2): 247–255.

Effects of maternal and infant co-infections, and of maternal immunisation, on the infant response to BCG and tetanus immunisation

Alison M. Elliott^{a,b,*}, Patrice A. Mawa^a, Emily L. Webb^b, Margaret Nampijja^a, Nancy Lyadda^c, Joseph Bukusuba^a, Moses Kizza^a, Proscovia B. Namujju^{a,1}, Juliet Nabulime^a, Juliet Ndibazza^a, Moses Muwanga^c, and James A.G. Whitworth^{a,b,2}

^aMRC/UVRI Uganda Research Unit on AIDS, P.O. Box 49, Entebbe, Uganda ^bLondon School of Hygiene & Tropical Medicine, Keppel Street, London WC1E 7HT, UK ^cEntebbe Hospital, P.O. Box 29, Entebbe, Uganda

Abstract

Some vaccines show poor efficacy in tropical countries. Within a birth cohort in Uganda, we investigated factors that might influence responses to BCG and tetanus immunisation. Whole blood assay responses to crude culture filtrate proteins of *Mycobacterium tuberculosis* (cCFP) and tetanus toxoid (TT) were examined among 1506 and 1433 one-year-olds, respectively. Maternal *Mansonella perstans* infection was associated with higher interleukin (IL)-10 responses to both immunogens but no reduction in gamma interferon (IFN- γ), IL-5 and IL-13 responses; other maternal helminth infections showed little effect. Tetanus immunisation during pregnancy was associated with higher infant responses to TT; maternal BCG scar (from past immunisation) with lower infant IL-5 and IL-13 responses to cCFP. IFN- γ , IL-5 and IL-13 to TT were reduced in HIV-exposed-uninfected infants; infant malaria and HIV were associated with lower IFN- γ , IL-5 and IL-13 responses to both immunogens. We conclude that maternal helminth infections are unlikely to explain poor vaccine efficacy in the tropics. Effects of maternal immunisation on infant responses to vaccines should be explored. Prevention of infant malaria and HIV could contribute to effectiveness of immunisation programmes.

Keywords

BCG; Tetanus; Immunisation

1 Introduction

Immunisation is key to the control of infectious diseases but the efficacy of some vaccines is poor in tropical, developing countries, where they are most needed [1]. In particular, Bacille

Â© 2010 Elsevier Ltd.

*Corresponding author at: MRC/UVRI Uganda Research Unit on AIDS, P.O. Box 49, Entebbe, Uganda. Tel.: +256 752 720609; fax: +256 414 321137. alison.tom@infocom.co.ug.

¹Current address: National Institute for Health and Welfare, Aapstie 1, P.O. Box 310, Oulu, Finland.

²Current address: The Wellcome Trust, 215 Euston Road, London NW1 2BE, UK.

This document was posted here by permission of the publisher. At the time of deposit, it included all changes made during peer review, copyediting, and publishing. The U.S. National Library of Medicine is responsible for all links within the document and for incorporating any publisher-supplied amendments or retractions issued subsequently. The published journal article, guaranteed to be such by Elsevier, is available for free, on ScienceDirect.

Calmette-Guérin (BCG) immunisation has over 70% efficacy against tuberculosis in temperate countries, but low efficacy in tropical settings [2,3]. The reasons for this need to be understood. Tuberculosis remains a major cause of global morbidity and mortality [4]; effective immunisation could revolutionise tuberculosis control and new vaccines are being developed [5], but new vaccines may be influenced by the same factors that inhibit the response to BCG in the tropics. Understanding these factors may allow the development of interventions to improve the effectiveness of immunisation programmes. Several hypotheses as to the nature of these factors have been advanced. Genetic differences between populations may be important, but efficacy in migrant populations tends to approach that observed in the native populations of the adoptive country [3,6,7]; differences in BCG strains used have been considered, but trials using the same source of BCG have also shown differences in efficacy by latitude [3]; effects of vaccine exposure to sunlight and breakdown in the cold chain have been considered, but are unlikely to explain low efficacy in carefully conducted trials. Two outstanding hypotheses particularly remain to be considered.

One of these is that exposure to environmental mycobacteria, which is more common in the tropics, masks, or blocks, the response to BCG in this setting. Early evidence for this hypothesis [3] has been supported by subsequent studies showing higher levels of sensitisation to mycobacterial antigens in unvaccinated Malawian compared to British populations, and smaller increases in the gamma interferon (IFN- γ) response following BCG in Malawian than in British adolescents [8]. However, vaccine-induced responses were not directly related to prior sensitisation to environmental mycobacteria [9], suggesting that other factors might play a role. Also, differences in response to BCG immunisation were demonstrated between Malawian and British infants at an age too young for effects to be explained by direct exposure to environmental organisms [10]; thus prenatal exposures are likely to be important.

A second hypothesis is that chronic helminth infections influence responses to BCG and other vaccines [11]. Helminths elicit strong type 2 and regulatory immune responses [12]; these effects can “spill over” to influence responses to unrelated antigens and can inhibit type 1 responses that are a component of the protective response against tuberculosis [13–16]. De-worming prior to BCG immunisation can improve the induced response to purified protein derivative of *Mycobacterium tuberculosis* [17]. Also, sensitisation to helminth antigens in utero may be associated with a switch to a type 2 response profile following BCG immunisation at birth, again emphasising the potential role of exposures very early in life [18].

In response to this last observation, we set up a randomised, controlled trial of anthelmintic treatment during pregnancy to investigate the hypothesis that exposure to, and treatment of, maternal worms during pregnancy would influence the infant response to BCG and other immunisations [19]. At age one year we assessed cytokine responses induced by BCG given at birth and by tetanus immunisation given at 6, 10 and 14 weeks of age. While there is no major concern regarding efficacy of tetanus immunisation, assessment of responses to tetanus toxoid (TT) allowed us to investigate effects on a vaccine that elicits a more type-2 biased response than BCG, and that is given later in infancy. The impact of the anthelmintic intervention on cytokine responses has been reported elsewhere [20]. We here describe planned observational analyses conducted to investigate factors affecting the infant response to immunisation during pre-natal and early post-natal life.

2 Materials and methods

2.1 Study design

The study was a randomised, double-blind, placebo-controlled trial of albendazole or praziquantel treatment during pregnancy, with a 2×2 factorial design, resulting in four arms, albendazole plus praziquantel, albendazole plus placebo for praziquantel, praziquantel plus placebo for albendazole and double placebo [ISRCTN32849447] [19]. Using the trial birth cohort, this observational analysis examined associations between infant cytokine responses to BCG and tetanus immunisation, and pre- and post-natal exposure to helminths, other co-infections and other potentially related factors.

2.2 Setting and participants

The study area comprised Entebbe Municipality and surrounding communities (Fig. 1). Women from the study area, in the second or third trimester of pregnancy, were recruited at Entebbe Hospital antenatal clinic between 2003 and 2005 if planning to deliver in the hospital and willing to know their HIV status; they were excluded for haemoglobin <8 g/dl, clinically apparent severe liver disease, diarrhoea with blood in stool, history of adverse reaction to anthelmintics, abnormal pregnancy, or if already enrolled during an earlier pregnancy.

The study was approved by ethical committees of the Uganda Virus Research Institute and London School of Hygiene & Tropical Medicine, and by the Uganda National Council for Science and Technology. All participants gave written informed consent.

2.3 Procedures

Socio-demographic details were recorded and blood and stool samples obtained prior to treatment of women with the trial intervention (single dose albendazole 400 mg or matching placebo and praziquantel 40 mg/kg or matching placebo). The intervention medication was given during the second or third trimester of pregnancy (according to when the women presented at the clinic and completed screening procedures). Women received standard antenatal care including haematinics and intermittent presumptive treatment for malaria with sulfadoxine–pyrimethamine. Tetanus immunisation, up to a maximum of three doses, was given during pregnancy unless the woman had completed a total of five doses during previous pregnancies. HIV-positive women were offered single dose nevirapine for themselves and their infants for prevention of mother-to-child HIV transmission [21]. Six weeks after delivery all women received treatment with both albendazole and praziquantel. Infants received BCG and polio immunisation at birth, polio, diphtheria, pertussis, tetanus, hepatitis B and *Haemophilus influenzae* immunisation at 6, 10 and 14 weeks of age, and measles immunisation at nine months. Infant illnesses were treated at the study clinic. At age 12 months blood was obtained from infants; weight and height were measured.

Vaccines were those provided by the Ugandan National Medical Stores: during the study period, BCG vaccine was provided from three suppliers: BB-NCIPD Ltd., Bulgaria, Serum Institute of India, India and Statens Seruminstitut, Denmark.

2.4 Diagnostic tests, parasitology and haematology

HIV serology was performed for mothers, and for infants aged 18 months, by rapid test algorithm [22]. HIV DNA PCR was performed [20], and HIV load measured (Bayer Versant branched DNA assay version 3.0; Bayer HealthCare, Leverkusen, Germany), for infants of HIV-positive mothers at age six weeks. Stools were examined for helminth ova by Kato-Katz method [23] and by culture for *Strongyloides* [24]; blood samples were examined by

modified Knott's method for microfilariae [25] and by thick film for malaria parasites, as previously described [22].

Clinical malaria was defined as fever ≥ 37.5 °C plus parasitaemia. Asymptomatic malaria was defined as parasitaemia in the absence of fever or other symptoms of malaria.

2.5 Immunological assays

Primary outcomes were infant immune responses to mycobacterial antigen and to TT, taken to represent the response to BCG and tetanus immunisation, respectively. We examined stimulated cytokine production in a whole blood assay, as described elsewhere: IFN- γ was measured to assess type 1 responses; IL-5 and IL-13 were measured to assess type 2 responses (since IL-4, the hallmark of the type 2 response, is seldom detectable in culture supernatant, particularly following stimulation with mycobacterial antigen) and IL-10 was measured to assess regulatory responses [26]. Briefly, unseparated, heparinised blood was diluted to a final concentration of one-in-four using RPMI supplemented with penicillin, streptomycin and glutamine, plated in 96-well plates, and stimulated with crude culture filtrate protein from *M. tuberculosis* (cCFP; 5 μ g/ml) (kindly provided by John Belisle, University of Colorado, Fort Collins, USA), TT (12 Lf/ml; Statens Seruminstitut, Denmark), phytohaemagglutinin (PHA; 10 μ g/ml; Sigma, UK), or left unstimulated. Supernatants were harvested on day 6 and frozen at -80 °C until analysed.

Cytokine concentrations in supernatants were measured by ELISA (Becton Dickinson, UK). Test responses were regarded as positive if greater than the mean plus two standard deviations of negative control results for all assays: IFN- γ > 73 pg/ml; IL-5 > 34 pg/ml; IL-13 > 18 pg/ml; IL-10 > 48 pg/ml. Values below the cut-off were set to zero. Cytokine production in unstimulated test wells was subtracted from concentrations produced in response to stimulation. Assays were performed after all samples had been collected, in a randomised sequence, to avoid confounding of secular trends with variations in assay performance.

2.6 Study size

The study size was determined for the trial objectives, rather than for this analysis. It was anticipated that, with recruitment of 2500 women, at least 1594 infants would be assessed at one year; this would give 80% power with $p < 0.05$ to detect differences of 0.11 \log_{10} in cytokine responses for exposures with two equal-sized categories [19].

2.7 Statistical analyses

The objective of this observational analysis was to determine socio-demographic, maternal and infant factors associated with cytokine responses following BCG and tetanus immunisation.

Socio-demographic factors were maternal age, maternal education (categories none, primary, secondary or tertiary), household socioeconomic status (a six-level score based on building materials, number of rooms, items owned) and location of residence (by zone, Fig. 1). Maternal factors were the three commonest maternal helminth infections (hookworm, *Mansonella perstans*, *Schistosoma mansoni*), maternal asymptomatic malaria parasitaemia (*Plasmodium falciparum*) and maternal immunisation status (absence or presence of a maternal BCG scar; number of documented doses of tetanus immunisation during pregnancy). Infant factors were gender, birth weight, anthropometric scores at age one year (weight-for-age, height-for-age and weight-for-height [27]), infant malaria (current, asymptomatic malaria on the day of the assay; number of documented clinical malaria

episodes in the preceding year) and HIV status (based on maternal and infant serology, and infant PCR at age six weeks: unexposed, exposed-uninfected, or infected).

Cytokine responses showed skewed distributions, with a disproportionate number of zero values, as has commonly been observed for immunoepidemiological data and, in particular, for the use of whole blood stimulation and cytokine response assays [28–30]. Results were transformed to $\log_{10}(\text{cytokine concentration} + 1)$ and analysed by linear regression using bootstrapping with 10,000 iterations to estimate standard errors and bias-corrected accelerated confidence intervals [29]. Regression coefficients and confidence limits were back-transformed to express results as ratios of geometric means. Crude associations were first examined. The following strategy was then employed to investigate multivariate associations. A simple hierarchical causal diagram was developed (Fig. 2). Socio-demographic factors were considered as potential confounders for the relationship between each exposure and cytokine response, and maternal co-infections (malaria parasitaemia and helminths) were considered as potential confounders for each other and for infant exposures. Treatment with albendazole was considered as a potential effect modifier for maternal hookworm and *M. perstans* infections, and treatment with praziquantel for *S. mansoni* infection. Infant co-infections were considered as potential confounders for infant anthropometric exposures. For all exposures, any remaining variable that showed a crude association with the outcome and that was not on the causal pathway between the exposure of interest and the outcome was included in the model to improve the precision of the estimated effects. BCG supplier (for analyses of response to BCG) and assay characteristics (antigen batch and lymphocyte count) were also considered in all models.

3 Results

3.1 Participants

The flow of participants through the study has been described elsewhere [20] and is summarised in Fig. 3. Of 2507 women enrolled, information was obtained on 2345 live births. Results from 1542 babies (singletons or older twins or triplets) were available at one year. Of these, 36 had not received BCG immunisation at Entebbe Hospital and 109 had incomplete tetanus immunisation: therefore 1506 infants were included in analyses for responses following BCG immunisation, and 1433 for tetanus immunisation.

As previously reported, the median maternal age was 23 years; most women (54%) had either primary or no formal education [31]. The majority (41%) lived in Entebbe Municipality, 28% in Manyago and Kabale, 11% Katabi roadside, 9% Katabi rural and 11% Kiggungu fishing village (Fig. 1). Sixty-eight percent had at least one helminth infection; 44% had hookworm, 21% *M. perstans* and 18% *S. mansoni*; 11% had asymptomatic malaria at enrolment; 12% had HIV infection [31]. Sixty percent had a BCG scar; 22%, 61% and 17% had zero, one and two or more recorded doses of tetanus immunisation during pregnancy, respectively. Women whose infants had cytokine results available at one year were older, of higher socioeconomic status and less likely to live in Katabi, and had lower prevalence of helminths, asymptomatic malaria and HIV infection during pregnancy, than those without results (data not shown).

Among infants with results at one year, 50% were female; the mean birth weight was 3.18 kg; at one year the mean weight-for-age *z* score was -0.33 , mean height-for-age *z* score -0.84 and mean weight-for-height *z* score 0.10 ; 6% had asymptomatic *P. falciparum* malaria; 9% were HIV-exposed-uninfected and 1% were HIV-infected. Only 44 of 1358 infants examined had helminth infections at age one year (most common were *Ascaris* (15 infants), *Trichuris* (12 infants) and *Mansonella* (eight infants)) so effects of infant helminths

were not considered in this analysis. Ninety-nine percent of infants were breast-fed to age six weeks, and 80% were still being breast-fed at age one year.

3.2 Cytokine responses

Type 1 (IFN- γ) and regulatory (IL-10) cytokines were dominant in the response to cCFP; following tetanus immunisation, type 2 cytokines were more prominent (Fig. 4). Crude associations between factors examined and cytokine responses are shown in Tables 1 and 2; multivariate analyses in Tables 3 and 4.

3.3 Socio-demographic factors

The infant IFN- γ and IL-5 response to TT increased with maternal education, with adjusted geometric mean ratios (aGMR) (95% confidence interval (CI)) of 1.25 (1.03, 1.54) and 1.25 (1.04, 1.50) respectively, while the IL-10 response to TT was inversely associated with socio-economic status (aGMR 0.90 (0.82, 0.98)).

3.4 Maternal factors

Maternal *M. perstans* infection showed a positive association with infant IL-10 production in response to both cCFP (aGMR 1.23 (0.99, 1.50) and TT (aGMR 1.36 (1.04, 1.80)). Both associations showed a marked interaction with maternal albendazole treatment (interaction *p*-values 0.02 and 0.001, respectively), being evident only in the albendazole-placebo group (cCFP aGMR 1.57 (1.19, 2.00) and TT aGMR 1.99 (1.35, 2.97)). No consistent associations were observed for other species.

Maternal BCG scar was associated with a markedly lower infant IL-5 and IL-13 responses to cCFP (aGMR 0.76 (0.61, 0.94) and 0.80 (0.64, 1.00)) and a somewhat lower IFN- γ response (aGMR 0.87 (0.70, 1.09)). An increasing number of doses of maternal tetanus immunisation during the pregnancy was associated with increased infant IFN- γ (aGMR 1.44 (1.16, 1.79)) and IL-13 (1.22 (1.01, 1.46)), and a weak increase in IL-5 (aGMR 1.19 (0.97, 1.44)) responses to TT.

3.5 Infant factors

Female infants had broadly lower responses for both cCFP and TT, with aGMRs for each cytokine response ranging from 0.69 to 0.86 (Tables 1–4).

Associations for anthropometric variables were somewhat variable; after adjustment for confounding, associations remained for the IL-13 response for TT and IL-10 response to cCFP, which both showed increased responses for higher scores: IL-13 for TT, birth weight aGMR 1.43 (1.09, 1.89), weight-for-age *z*-score at one year, 1.13 (1.01, 1.28), height-for-age *z*-score at one year 1.13 (1.01, 1.26); IL-10 for CFP, height-for-age *z*-score at one year, 1.08 (1.00, 1.17).

Current malaria parasitaemia was strongly associated with reduced IFN- γ , IL-5 and IL-13 responses for cCFP (aGMR 0.49 (0.28, 0.80), 0.41 (0.30, 0.60) and 0.46 (0.29, 0.75) respectively), and for TT (aGMR 0.47 (0.25, 0.85), 0.32 (0.21, 0.53) and 0.50 (0.26, 0.93) respectively), and with a reciprocal increase in IL-10 responses for TT (aGMR 2.38 (1.48, 3.80)). Previous episodes of malaria during infancy showed weaker effects, but a high number of episodes was associated with a reduced IL-5 response to cCFP (aGMR 0.84 (0.76, 0.95)) and an increased IL-10 response to TT (aGMR 1.18 (1.03, 1.34)).

Associations with infant HIV status differed for cCFP and TT. For cCFP, HIV-exposed-uninfected infants showed no difference in response compared to HIV-unexposed infants, but HIV-positive infants showed markedly lower IFN- γ , IL-5 and IL-13 responses (aGMR

0.06 (0.02, 0.23), 0.37 (0.25, 1.00) and 0.20 (0.09, 0.53) respectively), and higher IL-10 responses (aGMR 2.19 (1.56, 3.15)). For TT, both HIV-exposed-uninfected infants, and HIV-infected infants, showed impaired IFN- γ , IL-5 and IL-13 responses: HIV exposed-uninfected, aGMR 0.57 (0.35, 0.94), 0.51 (0.33, 0.82) and 0.61 (0.39, 0.95); HIV-infected, aGMR 0.35 (0.11, 1.13), 0.16 (0.10, 0.52) and 0.09 (0.04, 0.27); there was no effect on the IL-10 response.

4 Discussion

In this large birth cohort in Uganda, infant responses to BCG and tetanus immunisation were associated with pre-natal and early post-natal exposure to maternal *M. perstans* co-infection, maternal and infant HIV co-infection, and infant malaria. Maternal BCG scar showed associations with the infant response to BCG, and maternal immunisation with tetanus toxoid during pregnancy was associated with higher infant responses to their own tetanus immunisation.

As in any observational analysis, some findings may be explained by unmeasured confounders. However, most key factors identified were biological, rather than social or environmental, and adjustment for measured confounders produced little change in their effect estimates, suggesting that they are closely linked to causal mechanisms. Many statistical tests were conducted, so some apparently “significant” findings could have occurred by chance. Individual results are therefore treated with caution; rather than formally adjust for multiplicity, we focus on patterns and consistency of results, and on biological plausibility with reference to other findings.

Maternal *M. perstans* microfilaraemia was associated with enhanced IL-10 responses to both cCFP and TT in the offspring. This filarial infection is highly prevalent in Africa and central South America, but usually asymptomatic [32,33]. Adult worms inhabit serous cavities and microfilariae circulate in the blood, sometimes in thousands per millilitre, the lack of symptoms testifying to this helminth's potent immunoregulatory properties. Such helminth-induced regulation can influence host responses to unrelated antigens and IL-10 may be one key mediator of such effects [12]; among other filariases, IL-10 responses to tetanus immunisation have been found to be elevated in adults with asymptomatic *Onchocerca volvulus* infection [34,35]. Our key observation is that the non-specific effect of helminths on this regulatory cytokine response can be transmitted from mother to infant. Notably, infant IFN- γ , IL-5 and IL-13 responses were not reduced, suggesting the possibility that protective immune responses may not be impaired, and it is possible that the overall impact of exposure to maternal helminth infection in utero is an enhancement of regulatory immune responses rather than suppression of the ability to mount protective responses to vaccines and pathogens. This might be broadly beneficial, protecting against excessive inflammatory responses, including allergy [36,37].

The lack of observed effects of maternal hookworm or *S. mansoni* on type 1 and type 2 responses to mycobacterial antigens was surprising, given our own earlier findings [38], and those of Malhotra and colleagues [18]. However, in Malhotra's study all women had helminth infection: comparisons were made between infants sensitised and not sensitised to helminth antigens. Our study compared infants of mothers infected or not infected with each species, in a setting where most women had at least one helminth infection; moreover, for logistical reasons, a single stool sample was used for Kato Katz analysis giving limited sensitivity for diagnosis of intestinal helminths [39,40]. Both these factors may have resulted in underestimation of effects of schistosomiasis and other intestinal helminths. However, taken together with the finding (reported elsewhere [20]) that anthelmintics during pregnancy had little effect on infant responses to cCFP and TT in this study, these results

suggest that maternal helminth infection may not be the major explanation for the poor efficacy of BCG immunisation in the tropics. Subsequent acquisition of helminths by the infant may be a different story [17].

Tetanus immunisation during pregnancy was associated with enhanced IFN- γ , IL-13 and (to some extent) IL-5 responses following tetanus immunisation of the offspring. These results accord with the earlier report of Gill and colleagues [41] and show that priming of the infant response to TT can be influenced by immunisation of the mother. This antigen-specific effect may result from transfer of TT across the placenta within an immune complex, utilising the immunoglobulin receptor systems involved in transfer of maternal antibody to the fetus [42–44]. Fetal exposure to antigen can result in tolerisation, but immune complexes are potent activators of the immune system, and this may explain why priming occurred in this case. The lower response to tetanus immunisation in HIV-exposed-uninfected infants may have resulted from reduced transfer of maternal antibody and antigen in this group [45,46].

By contrast, presence of a maternal BCG scar showed a negative association with infant type 2 cytokine response, and (to some extent) IFN- γ response to cCFP following BCG immunisation. This may have been a non-specific effect since maternal BCG scar was also associated with reductions in these cytokine responses to PHA (data not shown). The association was not explained by adjusting for potential confounding factors, and suggests an immunological interaction between mother and infant related to maternal mycobacterial exposure or infection. There is evidence for sensitisation to mycobacterial antigens in utero in mouse models and in humans [47,48], but tolerisation is also a possibility, and would accord with the lower response to mycobacterial antigen observed in Malawian, compared to British, infants following BCG immunisation [10]. It may be important to investigate the role of maternal mycobacterial infection, and maternal immune responses to mycobacteria, in the infant response to BCG.

Current infant malaria and infant HIV infection were associated with broad reductions in IFN- γ , IL-5 and IL-13 responses. These findings were in keeping with the recognised immunosuppressive effects of these pathogens and thus, incidentally, demonstrate the ability of this immuno-epidemiological approach to detect important effects. They contrast with the IL-10-restricted effects of maternal *M. perstans*. For malaria, these effects were mainly short-term, observed only in the presence of current infection; however, decreased IL-5 cCFP and increased IL-10 TT responses were associated with both current and previous malaria. It is possible that the independent association between increased IL-10 TT responses and household socio-economic status might be mediated by repeated, unmeasured, exposures to infection.

Consistently lower responses were seen in girls. This shows that gender differences in immune response are present at an early age, and could be related to reported gender differences in the non-specific effects of immunisation on infant mortality [49].

This study examined factors influencing the cytokine responses induced by BCG and tetanus immunisation, not their efficacy. In the case of BCG, it is likely that IFN- γ is required, although not sufficient for, protective immunity [15], while excessive production of type 2 cytokines may be detrimental [50]. Excess production of IL-10 may also be detrimental, if it is associated with suppression of protective responses, but evidence from the mouse model suggests that adequate production may be required to prevent a pathological, inflammatory response [51]. Follow up of the cohort is in progress to determine how the observed responses are related to rates of *M. tuberculosis* infection and disease. In the case of tetanus immunisation, the induction of neutralising antibody is key to protective immunity [52]; the

relationship between observed effects on cytokine responses and the production of antibody will be the subject of further investigation. From a public health perspective, our results demonstrate strong effects of current, or recent infant infections on the infant response to vaccine antigens, and reinforce the importance of control and treatment of malaria and HIV infection for the immunological health of mothers and their children; but suggest that maternal helminth infection may have little, if any, adverse effect on the outcome of infant immunisation. Immunisation during pregnancy may enhance the infant response to selected vaccines, and this, as well as the role of prior maternal BCG immunisation and mycobacterial infection in determining the infant response to BCG immunisation, needs to be explored in further research.

References

1. Labeaud A.D. Malhotra I. King M.J. King C.L. King C.H. Do antenatal parasite infections devalue childhood vaccination? *PLoS Negl Trop Dis* 352009e44219478847
2. Colditz G.A. Berkey C.S. Mosteller F. Brewer T.F. Wilson M.E. Burdick E. The efficacy of bacillus Calmette-Guerin vaccination of newborns and infants in the prevention of tuberculosis: meta-analyses of the published literature *Pediatrics* 961 Pt 1199529357596718
3. Fine P.E. Variation in protection by BCG: implications of and for heterologous immunity *Lancet* 34689861995133913457475776
4. WHO. Global tuberculosis control – epidemiology, strategy, financing. WHO/HTM/TB/2009411; 2009.
5. Beresford B. Sadoff J.C. Update on research and development pipeline: tuberculosis vaccines *Clin Infect Dis* 50 Suppl. 32010S178S18320397946
6. Packe G.E. Innes J.A. Protective effect of BCG vaccination in infant Asians: a case-control study *Arch Dis Child* 63319882772813258499
7. Rodrigues L.C. Noel Gillo. Smith P.G. BCG vaccination in the first year of life protects children of Indian subcontinent ethnic origin against tuberculosis in England *J Epidemiol Community Health* 451199178802045751
8. Black G.F. Weir R.E. Floyd S. Bliss L. Warndorff D.K. Crampin A.C. BCG-induced increase in interferon-gamma response to mycobacterial antigens and efficacy of BCG vaccination in Malawi and the UK: two randomised controlled studies *Lancet* 359931520021393140111978337
9. Weir R.E. Black G.F. Nazareth B. Floyd S. Stenson S. Stanley C. The influence of previous exposure to environmental mycobacteria on the interferon-gamma response to Bacille Calmette-Guerin vaccination in southern England and northern Malawi *Clin Exp Immunol* 1463200639039917100757
10. Lalor M.K. Ben-Smith A. Gorak-Stolinska P. Weir R.E. Floyd S. Blitz R. Population differences in immune responses to Bacille Calmette-Guerin vaccination in infancy *J Infect Dis* 1996200979580019434928
11. Bentwich Z. Kalinkovich A. Weisman Z. Borkow G. Beyers N. Beyers A.D. Can eradication of helminthic infections change the face of AIDS and tuberculosis? *Immunol Today* 2011199948548710529774
12. van Riet E. Hartgers F.C. Yazdanbakhsh M. Chronic helminth infections induce immunomodulation: consequences and mechanisms *Immunobiology* 2126200747549017544832
13. Elias D. Akuffo H. Pawlowski A. Haile M. Schon T. Britton S. Schistosoma mansoni infection reduces the protective efficacy of BCG vaccination against virulent Mycobacterium tuberculosis *Vaccine* 231120051326133415661380
14. Elias D. Akuffo H. Thors C. Pawlowski A. Britton S. Low dose chronic Schistosoma mansoni infection increases susceptibility to Mycobacterium bovis BCG infection in mice *Clin Exp Immunol* 1393200539840415730384
15. Fletcher H.A. Correlates of immune protection from tuberculosis *Curr Mol Med* 73200731932517504116
16. Pearlman E. Kazura J.W. Hazlett F.E. Jr. Boom W.H. Modulation of murine cytokine responses to mycobacterial antigens by helminth-induced T helper 2 cell responses *J Immunol* 15191993485748648409444

17. Elias D, Wolday D, Akuffo H, Petros B, Bronner U, Britton S. Effect of deworming on human T cell responses to mycobacterial antigens in helminth-exposed individuals before and after Bacille Calmette-Guerin (BCG) vaccination. *Clin Exp Immunol* 123:22001-21922511207651
18. Malhotra I, Mungai P, Wamachi A, Kioko J, Ouma J, H. Kazura J, W. Helminth- and Bacillus Calmette-Guerin-induced immunity in children sensitized in utero to filariasis and schistosomiasis. *J Immunol* 162:1119996843684810352306
19. Elliott A, M. Kizza M, Quigley M, A. Ndibazza J, Nampijja M, Muhangi L. The impact of helminths on the response to immunization and on the incidence of infection and disease in childhood in Uganda: design of a randomized, double-blind, placebo-controlled, factorial trial of deworming interventions delivered in pregnancy and early childhood [ISRCTN32849447]. *Clin Trials* 4:12007425717327245
20. Webb EL, Mawa PA, Ndibazza J, Kizito D, Namatovu A, Kyosiimire-Lugemwa J, et al. The impact of single-dose anthelmintic treatment during the second or third trimester of pregnancy on the infant response to immunisation and on susceptibility to infectious diseases in infancy: results of a randomised double-blinded placebo-controlled trial. *Lancet*; in press.
21. Guay L, A. Musoke P, Fleming T, Bagenda D, Allen M, Nakabiito C. Intrapartum and neonatal single-dose nevirapine compared with zidovudine for prevention of mother-to-child transmission of HIV-1 in Kampala, Uganda: HIVNET 012 randomised trial. *Lancet* 354:9181199979580210485720
22. Muhangi L, Woodburn P, Omara M, Omoding N, Kizito D, Mpairwe H. Associations between mild-to-moderate anaemia in pregnancy and helminth, malaria and HIV infection in Entebbe, Uganda. *Trans R Soc Trop Med Hyg* 101:9200789990717555783
23. Katz N, Chaves A, Pellegrino J. A simple device for quantitative stool thick-smear technique in Schistosomiasis mansoni. *Rev Inst Med Trop Sao Paulo* 14:619723974004675644
24. Helminths F, J. Collee J, G. Fraser A, Marmion B, Simmons A, Mackie & McCartney, practical medical microbiology 1996 Churchill Livingstone Edinburgh
25. Melrose W, D. Turner P, F. Pisters P, Turner B. An improved Knott's concentration test for the detection of microfilariae. *Trans R Soc Trop Med Hyg* 94:2200017610897361
26. Elliott A, M. Hurst T, J. Balyeku M, N. Quigley M, A. Kaleebu P, French N. The immune response to Mycobacterium tuberculosis in HIV-infected and uninfected adults in Uganda: application of a whole blood cytokine assay in an epidemiological study. *Int J Tuberc Lung Dis* 3:3199923924710094326
27. WHO anthro, version 32009
28. Joseph S, Jones F, M. Kimani G, Mwatha J, K. Kamau T, Kazibwe F. Cytokine production in whole blood cultures from a fishing community in an area of high endemicity for Schistosoma mansoni in Uganda: the differential effect of parasite worm and egg antigens. *Infect Immun* 72:2200472873414742514
29. McGuinness D, Bennett S, Riley E. Statistical analysis of highly skewed immune response data. *J Immunol Methods* 2011:1997991149032413
30. Weir R, E. Brennan P, J. Butlin C, R. Dockrell H, M. Use of a whole blood assay to evaluate in vitro T cell responses to new leprosy skin test antigens in leprosy patients and healthy subjects. *Clin Exp Immunol* 116:2199926326910337017
31. Woodburn P, W. Muhangi L, Hillier S, Ndibazza J, Namujju P, B. Kizza M. Risk factors for helminth, malaria, and HIV infection in pregnancy in Entebbe, Uganda. *PLoS Negl Trop Dis* 3:62009e47319564904
32. Asio S, M. Simonsen P, E. Onapa A, W. Mansonella perstans filariasis in Uganda: patterns of microfilaraemia and clinical manifestations in two endemic communities. *Trans R Soc Trop Med Hyg* 103:3200926627318809192
33. Simonsen PE, Onapa AW, Asio SM. Mansonella perstans filariasis in Africa. *Acta Trop*. 2010, Feb 10 [Epub ahead of print], doi:10.1016/j.actatropica.2010.01.014.
34. Cooper P, J. Espinel I, Paredes W, Guderian R, H. Nutman T, B. Impaired tetanus-specific cellular and humoral responses following tetanus vaccination in human onchocerciasis: a possible role for interleukin-10. *J Infect Dis* 178:41998113311389806045

35. Nookala S, Srinivasan S, Kaliraj P, Narayanan R, B, Nutman T, B. Impairment of tetanus-specific cellular and humoral responses following tetanus vaccination in human lymphatic filariasis *Infect Immun* 72:520042598260415102768
36. Elliott A, M, Mpairwe H, Quigley M, A, Nampijja M, Muhangi L, Oweka-Onyee J. Helminth infection during pregnancy and development of infantile eczema *JAMA* 294:1620052032203416249415
37. Mpairwe H, Webb E, Muhangi L, Ndibazza J, Tumusiime J, Rodrigues L. Effects of treatment of helminths with albendazole and praziquantel in pregnancy on the incidence of allergy in the first year of life: trial results *Allergy* 64 Suppl. 90:200992
38. Elliott A, M, Namujju P, B, Mawa P, A, Quigley M, A, Nampijja M, Nkurunziza P, M, A. A randomised controlled trial of the effects of albendazole in pregnancy on maternal responses to mycobacterial antigens and infant responses to Bacille Calmette-Guerin (BCG) immunisation [ISRCTN32849447] *BMC Infect Dis* 5:200511516371154
39. Knopp S, Rinaldi L, Khamis I, S, Stothard J, R, Rollinson D, Maurelli M, P, A. A single FLOTAC is more sensitive than triplicate Kato-Katz for the diagnosis of low-intensity soil-transmitted helminth infections *Trans R Soc Trop Med Hyg* 103:4200934735419168197
40. Utzinger J, Booth M, N', Goran E, K, Muller I, Tanner M, Lengeler C. Relative contribution of day-to-day and intra-specimen variation in faecal egg counts of *Schistosoma mansoni* before and after treatment with praziquantel *Parasitology* 122 Pt 5:200153754411393827
41. Gill T, J, 3rd, Repetti C, F, Metlay L, A, Rabin B, S, Taylor F, H, Thompson D, S. Transplacental immunization of the human fetus to tetanus by immunization of the mother *J Clin Invest* 72:319839879966886013
42. Malek A. Ex vivo human placenta models: transport of immunoglobulin G and its subclasses *Vaccine* 21:2420033362336412850340
43. May K, Grube M, Malhotra I, Long C, A, Singh S, Mandalika K. Antibody-dependent transplacental transfer of malaria blood-stage antigen using a human ex vivo placental perfusion model *PLoS One* 4:112009e798619956710
44. Schneider H, Miller R, K. Receptor-mediated uptake and transport of macromolecules in the human placenta *Int J Dev Biol* 54:2–3201036737519876831
45. de Moraes-Pinto M, I, Verhoeff F, Chimsuku L, Milligan P, J, Wesumperuma L, Broadhead R, L. Placental antibody transfer: influence of maternal HIV infection and placental malaria *Arch Dis Child Fetal Neonatal Ed* 79:31998F202F20510194992
46. Scott S, Cumberland P, Shulman C, E, Cousens S, Cohen B, J, Brown D, W. Neonatal measles immunity in rural Kenya: the influence of HIV and placental malaria infections on placental transfer of antibodies and levels of antibody in maternal and cord serum samples *J Infect Dis* 191:1120051854186015871118
47. Malhotra I, Ouma J, Wamachi A, Kioko J, Mungai P, Omollo A. In utero exposure to helminth and mycobacterial antigens generates cytokine responses similar to that observed in adults *J Clin Invest* 99:71997175917669120021
48. Rahman M, J, Degano I, R, Singh M, Fernandez C. Influence of maternal gestational treatment with mycobacterial antigens on postnatal immunity in an experimental murine model *PLoS One* 5:32010e969920300629
49. Aaby P, Jensen H, Rodrigues A, Garly M, L, Benn C, S, Lisse I, M. Divergent female–male mortality ratios associated with different routine vaccinations among female–male twin pairs *Int J Epidemiol* 33:200436737315082642
50. Rook G, A, Hernandez-Pando R, Zumla A. Tuberculosis due to high-dose challenge in partially immune individuals: a problem for vaccination? *J Infect Dis* 199:5200961361819199549
51. Higgins D, M, Sanchez-Campillo J, Rosas-Taraco A, G, Lee E, J, Orme I, M, Gonzalez-Juarrero M. Lack of IL-10 alters inflammatory and immune responses during pulmonary *Mycobacterium tuberculosis* infection *Tuberculosis (Edinb)* 89:2200914915719213603
52. Dietz V, Galazka A, van Loon F, Cochi S. Factors affecting the immunogenicity and potency of tetanus toxoid: implications for the elimination of neonatal and non-neonatal tetanus as public health problems *Bull World Health Organ* 75:1199781939141753

Acknowledgments

We thank all staff and participants of the Entebbe Mother and Baby Study, the midwives of the Entebbe Hospital Maternity Department, the community field team in Entebbe and Katabi, and the staff of the Clinical Diagnostic Services Laboratory at the MRC/UVRI Uganda Research Unit on AIDS. We thank Dr. Stephen Cose for critical review of the manuscript. The study was funded by Wellcome Trust grant numbers 064693 and 079110; mycobacterial antigens were provided through the National Institutes of Health contract NOI-AI-25147.

Conflict of interest: James Whitworth is now a member of staff with the Wellcome Trust, the funders of the study. His role in the initial design and conduct of the study preceded his appointment at the Wellcome Trust. He has had no role in the study since his appointment. None of the other authors has any conflict of interest in relation to this work.




Fig. 1.
Study area.




Fig. 2.
Causal diagram.




Fig. 3.
Flow of participants through the study.




Fig. 4. Cytokine responses measured in supernatant of a six-day whole blood assay among infants aged one year. (A) Responses to crude culture filtrate proteins of *Mycobacterium tuberculosis* among infants who received BCG immunisation at birth. (B) Responses to tetanus toxoid among infants who received three doses of tetanus immunisation during infancy. The central line represents the median; boxes represent 75th and 25th centiles; whiskers represent upper and lower adjacent values and dots represent outside values.

Table 1

Crude associations with response to crude culture filtrate proteins of *Mycobacterium tuberculosis* in infants who received BCG immunisation at birth.

Factor ^a	IFN- γ	IL-5	IL-13	IL-10
Socio-demographic characteristics				
Household socioeconomic status	1.02 (0.94, 1.11)	1.05 (0.97, 1.15)	1.04 (0.95, 1.14)	0.99 (0.91, 1.07)
Mother's education	1.03 (0.88, 1.19)	0.99 (0.86, 1.15)	1.03 (0.88, 1.21)	0.92 (0.80, 1.05)
Mother's age at enrolment	1.01 (0.99, 1.03)	1.02 (1.00, 1.03)	1.01 (0.99, 1.03)	1.00 (0.98, 1.02)
Location of residence				
Entebbe Town, urban	1	1	1	1
Manyago and Kabale, semi-urban	1.07 (0.82, 1.37)	1.01 (0.79, 1.29)	0.93 (0.72, 1.21)	1.01 (0.79, 1.27)
Kigungu, fishing village	1.10 (0.76, 1.51)	1.00 (0.71, 1.44)	0.98 (0.68, 1.39)	1.19 (0.85, 1.60)
Katabi, roadside	1.15 (0.76, 1.66)	0.87 (0.63, 1.27)	1.34 (0.92, 1.92)	1.36 (0.96, 1.83)
Katabi, rural	0.91 (0.54, 1.40)	0.66 (0.46, 0.99)	0.86 (0.54, 1.35)	1.67 (1.12, 2.30)
Maternal characteristics				
Maternal hookworm infection at enrolment	0.89 (0.71, 1.11) ^b	0.93 (0.76, 1.15)	1.01 (0.81, 1.26)	0.91 (0.75, 1.10)
Maternal <i>S. mansoni</i> infection at enrolment	0.80 (0.58, 1.07)	0.83 (0.64, 1.08)	0.91 (0.68, 1.20)	0.71 (0.53, 0.93)
Maternal <i>M. perstans</i> infection at enrolment	0.95 (0.72, 1.21)	0.82 (0.64, 1.04)	0.95 (0.72, 1.23)	1.31 (1.03, 1.63)^c
Maternal malaria parasitaemia at enrolment	0.88 (0.57, 1.23)	0.61 (0.46, 0.86)	0.81 (0.56, 1.16)	1.33 (0.95, 1.71)
Maternal BCG scar	0.85 (0.68, 1.06)	0.77 (0.62, 0.95)	0.77 (0.62, 0.97)	1.00 (0.82, 1.22)
Infant characteristics				
Infant female	0.77 (0.63, 0.95)	0.74 (0.61, 0.91)	0.79 (0.64, 0.98)	0.85 (0.70, 1.03)
Birth weight	1.21 (0.97, 1.53)	1.17 (0.93, 1.48)	1.27 (0.98, 1.63)	1.14 (0.93, 1.42)
Weight for age z score at 1 year	1.10 (1.01, 1.21)	1.09 (1.00, 1.19)	1.04 (0.95, 1.14)	1.09 (1.01, 1.18)
Height for age z score at 1 year	1.06 (0.97, 1.16)	1.10 (1.01, 1.20)	1.08 (0.99, 1.18)	1.05 (0.97, 1.14)
Weight for height z score at 1 year	1.09 (0.99, 1.19)	1.05 (0.97, 1.14)	1.00 (0.91, 1.09)	1.07 (0.99, 1.16)
Infant malaria (current, asymptomatic)	0.47 (0.27, 0.75)	0.43 (0.32, 0.62)	0.48 (0.30, 0.77)	0.97 (0.63, 1.35)
Number of malaria episodes in infancy	0.97 (0.85, 1.09)	0.84 (0.76, 0.94)	1.00 (0.88, 1.12)	1.09 (0.96, 1.21)
Infant HIV status				
Unexposed	1	1	1	1
Exposed, uninfected	1.00 (0.64, 1.44)	1.00 (0.72, 1.44)	1.12 (0.77, 1.63)	0.87 (0.59, 1.21)
Infected	0.06 (0.02, 0.22)	0.33 (0.26, 0.74)	0.16 (0.09, 0.38)	2.20 (1.49, 3.21)

Data are for 1506 children who received BCG immunisation at Entebbe Hospital. Confidence intervals not including one are highlighted in bold.

^aMissing values: maternal BCG scar, 54; maternal malaria, 26; maternal education, 3; household socioeconomic status, 32; location of residence, 15; birth weight 248; weight for age, 2; height for age, 16; weight for height z scores, 17; infant malaria parasitaemia, 28.

^bEffect of maternal hookworm on IFN- γ response modified by maternal treatment with albendazole: albendazole group, GMR = 0.67 (0.48, 0.90); placebo group GMR = 1.19 (0.88, 1.61).

^cEffect of maternal *M. perstans* on IL-10 response modified by maternal treatment with albendazole: albendazole group, GMR = 0.97 (0.66, 1.36); placebo group GMR = 1.69 (1.27, 2.19).

Table 2

Crude associations for infant response to tetanus toxoid after tetanus immunisation at 6, 10 and 14 weeks of age.

Factor ^a	IFN- γ ^b	IL-5	IL-13	IL-10
Socio-demographic characteristics				
Household socioeconomic status	1.09 (0.97, 1.23)	1.03 (0.92, 1.15)	1.03 (0.93, 1.14)	0.91 (0.82, 1.00)
Mother's education	1.22 (1.00, 1.49)	1.22 (1.01, 1.47)	1.15 (0.97, 1.36)	0.88 (0.75, 1.03)
Mother's age at enrolment	0.98 (0.96, 1.01)	1.00 (0.98, 1.03)	0.99 (0.97, 1.01)	0.99 (0.97, 1.01)
Location of residence				
Entebbe Town, urban	1	1	1	1
Manyago and Kabale, semi-urban	0.88 (0.62, 1.22)	0.82 (0.61, 1.13)	0.77 (0.58, 1.04)	0.94 (0.72, 1.23)
Kigungu, fishing village	1.00 (0.61, 1.64)	1.12 (0.72, 1.75)	1.02 (0.67, 1.50)	1.31 (0.90, 1.94)
Katabi, roadside	1.13 (0.67, 1.83)	1.34 (0.84, 2.11)	1.33 (0.86, 1.94)	1.48 (1.01, 2.18)
Katabi, rural	0.92 (0.50, 1.66)	0.61 (0.36, 1.08)	0.65 (0.37, 1.09)	2.00 (1.23, 3.28)
Maternal characteristics				
Maternal hookworm infection at enrolment	0.85 (0.65, 1.14)	0.95 (0.73, 1.22) ^c	0.87 (0.69, 1.10) ^d	1.17 (0.94, 1.46)
Maternal <i>S. mansoni</i> infection at enrolment	0.88 (0.61, 1.29)	0.97 (0.69, 1.36)	0.96 (0.69, 1.30)	0.81 (0.61, 1.08)
Maternal <i>M. perstans</i> infection at enrolment	0.87 (0.61, 1.24)	1.03 (0.75, 1.44)	0.95 (0.70, 1.27)	1.46 (1.11, 1.95)^e
Maternal malaria parasitaemia at enrolment	0.85 (0.52, 1.38)	0.70 (0.45, 1.09)	0.79 (0.51, 1.20)	1.09 (0.74, 1.61)
Number of doses of tetanus toxoid during pregnancy	1.45 (1.18, 1.79)	1.21 (0.99, 1.47)	1.25 (1.05, 1.50)	0.86 (0.72, 1.02)
Infant characteristics				
Infant female	0.70 (0.53, 0.93)	0.78 (0.61, 1.01)	0.85 (0.67, 1.08)	0.77 (0.62, 0.95)
Birth weight	1.06 (0.77, 1.47)	1.28 (0.95, 1.69)	1.43 (1.08, 1.87)	0.88 (0.69, 1.13)
Weight for age z score at 1 year	1.11 (0.98, 1.25)	1.19 (1.07, 1.32)	1.20 (1.09, 1.33)	1.04 (0.94, 1.15)
Height for age z score at 1 year	1.10 (0.98, 1.23)	1.14 (1.02, 1.27)	1.22 (1.11, 1.35)	0.89 (0.81, 0.98)
Weight for height z score at 1 year	1.08 (0.96, 1.22)	1.15 (1.03, 1.29)	1.12 (1.01, 1.24)	1.14 (1.03, 1.25)
Infant malaria (current, asymptomatic)	0.45 (0.24, 0.83)	0.32 (0.20, 0.54)	0.45 (0.26, 0.74)	2.73 (1.65, 4.43)
Number of malaria episodes in infancy	0.92 (0.78, 1.08)	0.90 (0.79, 1.05)	0.87 (0.75, 0.98)	1.24 (1.09, 1.41)
Infant HIV status				
Unexposed	1	1	1	1
Exposed, uninfected	0.57 (0.34, 0.94)	0.53 (0.35, 0.84)	0.67 (0.43, 1.02)	0.94 (0.64, 1.41)
Infected	0.30 (0.10, 1.00)	0.15 (0.09, 0.59)	0.09 (0.04, 0.25)	0.81 (0.34, 2.64)

Data are for 1433 children who received three doses of tetanus immunisation during infancy. Confidence intervals not including one are highlighted in bold.

^aMissing values: maternal doses of tetanus immunisation, 2; maternal malaria, 24; maternal education, 3; household socioeconomic status, 28; location of residence, 15; birth weight 233; height and weight for height z scores, 12; infant malaria parasitaemia, 26.

^bOne additional missing value for the IFN- γ response to tetanus toxoid.

^cEffect of maternal hookworm on IL-5 response modified by maternal treatment with albendazole: albendazole group, GMR = 0.70 (0.48, 1.01); placebo group GMR = 1.26 (0.87, 1.82).

^dEffect of maternal hookworm on IL-13 response modified by maternal treatment with albendazole: albendazole group, GMR = 0.61 (0.44, 0.86); placebo group GMR = 1.23 (0.88, 1.75).

^eEffect of maternal *M. perstans* on IL-10 response modified by maternal treatment with albendazole: albendazole group, GMR = 0.95 (0.65, 1.43); placebo group GMR = 2.17 (1.46, 3.21).

Table 3

Factors associated with the cytokine response to crude culture filtrate proteins of *Mycobacterium tuberculosis* in infants who received BCG immunisation at birth, after adjustment for potential confounders and other crudely associated factors.

Cytokine/factor	Adjusted GMR (95% CI) ^a
Interferon- γ	
Maternal hookworm infection	
Albendazole treatment group	0.70 (0.50, 0.96)
Albendazole placebo group	1.29 (0.95, 1.76)
Baby female	0.79 (0.64, 0.98)
Baby malaria (current, asymptomatic)	0.49 (0.28, 0.80)
Baby HIV status	
Unexposed	1
Exposed, uninfected	0.96 (0.62, 1.36)
Infected	0.06 (0.02, 0.23)
Interleukin-5	
Maternal BCG scar	0.76 (0.61, 0.94)
Baby female	0.72 (0.58, 0.89)
Baby malaria (current, asymptomatic)	0.41 (0.30, 0.60)
Number of malaria episodes during infancy	0.84 (0.76, 0.95)
Baby HIV status	
Unexposed	1
Exposed, uninfected	1.01 (0.71, 1.49)
Infected	0.37 (0.25, 1.00)
Interleukin-13	
Maternal BCG scar	0.80 (0.64, 1.00)
Baby malaria (current, asymptomatic)	0.46 (0.29, 0.75)
Baby HIV status	
Unexposed	1
Exposed, uninfected	1.09 (0.74, 1.59)
Infected	0.20 (0.09, 0.53)
Interleukin-10	
Maternal <i>S. mansoni</i>	0.76 (0.58, 0.97)
Maternal <i>M. perstans</i>	
Albendazole treatment	0.93 (0.66, 1.27)
Albendazole placebo	1.57 (1.19, 2.00)
Baby height-for-age z-score at 1 year	1.08 (1.00, 1.17)
Baby HIV status	
Unexposed	1
Exposed, uninfected	0.83 (0.58, 1.14)
Infected	2.19 (1.56, 3.15)

^aFigures are geometric mean ratios of cytokine production, with bootstrap 95% confidence intervals.

Table 4

Factors remaining associated with the cytokine response to tetanus toxoid in infants who received tetanus immunisation at 6, 10 and 14 weeks of age, after adjustment for potential confounders and other crudely associated factors.

Cytokine/factor	Adjusted GMR (95% CI)^a
Interferon-γ	
Maternal education	1.25 (1.03, 1.54)
Number of doses of tetanus toxoid during pregnancy	1.44 (1.16, 1.79)
Baby female	0.69 (0.52, 0.91)
Baby malaria (current, asymptomatic)	0.47 (0.25, 0.85)
Baby HIV status	
Unexposed	1
Exposed, uninfected	0.57 (0.35, 0.94)
Infected	0.35 (0.11, 1.13)
Interleukin-5	
Maternal education	1.25 (1.04, 1.50)
Baby malaria (current, asymptomatic)	0.32 (0.21, 0.53)
Baby HIV status	
Unexposed	1
Exposed, uninfected	0.51 (0.33, 0.82)
Infected	0.16 (0.10, 0.52)
Interleukin-13	
Location of residence	
Entebbe Town	1
Manyago and Kabale	0.99 (0.64, 1.47)
Kigungu	0.75 (0.56, 0.99)
Katabi, roadside	1.30 (0.86, 1.92)
Katabi, rural	0.60 (0.35, 1.02)
Maternal hookworm	
Albendazole treatment	0.64 (0.45, 0.88)
Albendazole placebo	1.33 (0.95, 1.89)
Number of doses of tetanus toxoid during pregnancy	1.22 (1.01, 1.46)
Birth weight	1.43 (1.09, 1.89)
Weight for age z score 1 year	1.13 (1.01, 1.28)
Weight for height z score 1 year	1.13 (1.01, 1.26)
Baby malaria (current, asymptomatic)	0.50 (0.26, 0.93)
Baby HIV status	
Unexposed	1
Exposed, uninfected	0.61 (0.39, 0.95)
Infected	0.09 (0.04, 0.27)
Interleukin-10²	
Household socioeconomic status	0.90 (0.82, 0.98)

Cytokine/factor	Adjusted GMR (95% CI)^a
Maternal <i>M. perstans</i>	
Albendazole treatment	0.87 (0.60, 1.30)
Albendazole placebo	1.99 (1.35, 2.97)
Baby female	0.78 (0.63, 0.97)
Baby malaria (current, asymptomatic)	2.38 (1.48, 3.80)
Total malaria episodes in infancy	1.18 (1.03, 1.34)

^aFigures are geometric mean ratios of cytokine production, with bootstrap 95% confidence intervals.