

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Singh, A; Gupta, V; Ghosh, A; Lock, K; Ghosh-Jerath, S (2015)
Quantitative estimates of dietary intake with special emphasis on
snacking pattern and nutritional status of free living adults in ur-
ban slums of Delhi: impact of nutrition transition. *BMC nutrition*,
1. p. 22. ISSN 2055-0928 DOI: <https://doi.org/10.1186/s40795-015-0018-6>

Downloaded from: <http://researchonline.lshtm.ac.uk/2534110/>

DOI: [10.1186/s40795-015-0018-6](https://doi.org/10.1186/s40795-015-0018-6)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

Supplementary Table 3 Coefficients of variation and variance ratio for different nutrients per consumption unit

Nutrient	Coefficient of variation (%)		Variance Ratio (S_w^2/S_b^2)
	Within- Household (CV_w)	Between- Household (CV_b)	
Energy	23.68	28.08	0.71
Protein	33.80	27.34	1.53
Fat	36.37	45.44	0.64
Vitamin A	67.05	91.47	0.54
Vitamin C	92.22	49.56	3.46
Thiamine	36.61	33.46	1.20
Riboflavin	31.78	32.82	0.94
Niacin	33.62	32.88	1.05
Calcium	30.17	42.61	0.50
Folic Acid	40.41	31.33	1.66
Iron	37.00	32.98	1.26

Values in bold indicate greater within- household than between- household coefficient of variation for nutrients