

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Jordans, M; Rathod, S; Fekadu, A; Medhin, G; Kigozi, F; Kohrt, B; Luitel, N; Petersen, I; Shidhaye, R; Ssebunnya, J; +2 more... Patel, V; Lund, C; (2017) Suicidal ideation and behaviour among community and health care seeking populations in five low- and middle-income countries: a cross-sectional study. *Epidemiology and psychiatric sciences*. pp. 1-10. ISSN 2045-7960 DOI: <https://doi.org/10.1017/S2045796017000038>

Downloaded from: <http://researchonline.lshtm.ac.uk/3515653/>

DOI: <https://doi.org/10.1017/S2045796017000038>

Usage Guidelines:

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

Table 1: Sociodemographic characteristics of community study and facility detection study participants in PRIME implementation areas in Ethiopia, India, Nepal, South Africa and Uganda, 2013-2014.

	Ethiopia		India		Nepal		South Africa	Uganda
	Community Residents*	Facility attendees	Community Residents*	Facility attendees	Community Residents*	Facility attendees	Facility attendees	Facility attendees
Sample size	1486	1017	3220	760	1983	1474	1322	1897
Age, years (SD)	39.4 (15.3)	35.5 (15.6)	40.2 (15.1)	39.7 (14.8)	39.8 (15.4)	39.5 (15.0)	46.9 (13.3)	31.1 (11.7)
Female (%)	50.6	54.2	45.4	50.8	60.1	65.8	75.0	79.6
Ever married / cohabitating (%)	84.8	73.2	89.5	91.5	86.5	91.4	69.1	**
Illiterate / did not complete primary (%)	45.6	39.3	50.9	50.1	13.2	36.0	23.3	9.1

Note: *Community Survey figures are adjusted for the complex sampling design within each site; **Marital status only asked of participants who screened positive for depression or alcohol use disorders, rather than for the full sample.