

LONDON  
SCHOOL of  
HYGIENE  
& TROPICAL  
MEDICINE



Goodman, SG; Nicolau, JC; Requena, G; Maguire, A; Blankenberg, S; Chen, JY; Granger, CB; Grieve, R; Pocock, SJ; Simon, T; Yasuda, S; Vega, AM; Brieger, D; TIGRIS Study Investigators, (2017) Longer-term oral antiplatelet use in stable post-myocardial infarction patients: Insights from the long Term rIsk, clinical manaGement and healthcare Resource utilization of stable coronary artery dISease (TIGRIS) observational study. *International journal of cardiology*, 236. pp. 54-60. ISSN 0167-5273 DOI: <https://doi.org/10.1016/j.ijcard.2017.02.062>

Downloaded from: <http://researchonline.lshtm.ac.uk/3592093/>

DOI: [10.1016/j.ijcard.2017.02.062](https://doi.org/10.1016/j.ijcard.2017.02.062)

#### Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact [researchonline@lshtm.ac.uk](mailto:researchonline@lshtm.ac.uk).

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

## Supplementary Tables

**Supplementary Table 1 – Body mass index, waist circumference, heart rate, and blood pressure at enrolment by geographical region**

	Total population	Asia-Pacific/ Australia	Europe	North America	South America
Number of patients	9225 (100)	2850 (30.89)	4240 (45.96)	1024 (11.1)	1111 (12.04)
BMI, kg/m <sup>2</sup> measured	8964	2713	4138	1014	1099
Median*	26.81 (24.21-29.82)	24.77 (22.56-27.24)	27.68 (25.16-30.47)	28.59 (25.25-32.5)	27.57 (25.07-30.45)
Underweight (<20)	275 (3.07)	191 (7.04)	42 (1.01)	23 (2.27)	19 (1.73)
Normal (20-<25)	2629 (29.33)	1250 (46.07)	924 (22.33)	205 (20.22)	250 (22.75)
Overweight (25-30)	3890 (43.4)	1001 (36.9)	1976 (47.75)	391 (38.56)	522 (47.5)
Obese (>30)	2170 (24.21)	271 (9.99)	1196 (28.9)	395 (38.95)	308 (28.03)
Waist circumference (cm) measured	7711	2208	3548	882	1073
Median*	97 (89-106)	90 (83-96.85)	101 (93-109)	99.03 (91.44-109.22)	99 (92-107)
Heart rate (beats/ min)	8891	2716	4054	1014	1107
Median*	68 (60-75)	70 (64-77)	66 (60-74)	67 (60-74)	68 (62-73)
Systolic blood pressure (mm Hg) measured	9012	2774	4109	1020	1109
Median*	130 (120-140)	128 (118-138)	134 (121-146)	128 (117.5-140)	128 (120-140)
Diastolic blood pressure (mm Hg) measured	9004	2767	4109	1019	1109
Median*	77 (70-82)	75 (69-80)	80 (70-85)	74 (68-81)	80 (70-80)

All numbers are given as numbers (%), except \*median (25<sup>th</sup>, 75<sup>th</sup> percentiles)

BMI=Body Mass Index; mm Hg=millimeters of Mercury

**Supplementary Table 2 – Reasons for cessation of antiplatelet therapy prescribed following the index MI and discontinued prior to enrolment**

	ASA	Clopidogrel	Prasugrel	Ticagrelor	Ticlopidine
Number of patients	1028	3717	934	829	16
Intolerance or hypersensitivity	4 (0.4)	7 (0.2)	2 (0.2)	11 (1.3)	0 (0)
GI bleeding	16 (1.6)	15 (0.4)	5 (0.5)	5 (0.6)	0 (0)
Other bleeding complication	20 (2.0)	25 (0.7)	7 (0.8)	6 (0.7)	0 (0)
Other GI side effect	19 (1.9)	9 (0.2)	0 (0)	2 (0.2)	0 (0)
Physician advice/termination	652 (63.4)	3344 (90.0)	814 (87.1)	753 (90.8)	16 (100)
Patient wish	25 (2.4)	54 (1.5)	8 (0.9)	3 (0.4)	0 (0)
Other	290 (28.2)	262 (7.1)	98 (10.5)	49 (5.9)	0 (0)

All numbers are given as numbers (%)

GI=Gastrointestinal