

Table 3: Clinical information and disease stage at presentation for patients identified as having glaucoma or suspected glaucoma via the pilot

Diagnosis (n)	MD <sup>a</sup> Value (95% CI)	I.O.P <sup>b</sup> Value (95% CI)	C.D.R <sup>c</sup> Value (95% CI)
Suspected (8)	-2.9 (-5.8 to -0.1)	17.5 (14.9 to 20.1)	0.53 (0.37 to 0.67)
POAG or PAC(G) (4)	-13.3 (-24.9 to -1.6)	22.0 (15.5 to 28.5)	0.38 (0.04 to 0.70)
Stage of presentation (n)*	MD <sup>a</sup> Value (95% CI)	I.O.P <sup>b</sup> Value (95% CI)	C.D.R <sup>c</sup> Value (95% CI)
Early	-	-	-
Moderate (3)	-9.7 (-15.6 to -3.8)	23.3 (13.9 to 32.7)	0.30 (-0.13 to 0.73)
Advanced	-	-	-
Severe (1)	-23.8	18.0	0.60

<sup>a</sup>Mean deviation in the worst eye (defined as the eye with the poorest MD score);

<sup>b</sup>Intraocular pressure in the worst eye (mmHg); <sup>c</sup>Cup-disc ratio in the worst eye

\*Based on Mills et al [17]