

**Supplementary table 1. Stage-specific sets of weights used for age standardisation of ovarian cancer estimates**

<b>FIGO stage</b>	<b>Age category</b>	<b>Weights</b>	<b>SEER Summary Stage 2000</b>	<b>Age category</b>	<b>Weights</b>
All patients	15-44	0.10	All patients	15-44	0.10
	45-54	0.15		45-54	0.15
	55-64	0.25		55-64	0.24
	65-74	0.24		65-74	0.24
	75-84	0.20		75-84	0.20
	85-99	0.07		85-99	0.07
TNM stage I	15-44	0.21	Localised	15-44	0.24
	45-54	0.21		45-54	0.22
	55-64	0.25		55-64	0.24
	65-74	0.18		65-74	0.17
	75-84	0.11		75-84	0.11
	85-99	0.03		85-99	0.03
TNM stage II	15-44	0.10	Regional	15-44	0.13
	45-54	0.20		45-54	0.22
	55-64	0.26		55-64	0.26
	65-74	0.24		65-74	0.21
	75-84	0.16		75-84	0.15
	85-99	0.04		85-99	0.04
TNM stage III	15-44	0.06	Distant	15-44	0.05
	45-54	0.15		45-54	0.13
	55-64	0.29		55-64	0.26
	65-74	0.28		65-74	0.28
	75-84	0.19		75-84	0.21
	85-99	0.04		85-99	0.06
TNM stage IV	15-44	0.04	Missing stage	15-44	0.09
	45-54	0.11		45-54	0.11
	55-64	0.24		55-64	0.19
	65-74	0.30		65-74	0.21
	75-84	0.24		75-84	0.26
	85-99	0.07		85-99	0.14
Missing stage	15-44	0.08			
	45-54	0.11			
	55-64	0.19			
	65-74	0.21			
	75-84	0.27			
	85-99	0.14			