

**Table V: Multivariable analysis of factors associated with incident MDD and persistent MDD**

Factor	Category	Incident MDD		Persistent MDD	
		Multivariable OR (95% CI)	p-value	Multivariable OR (95% CI)	p-value
<b>Socio-economic factors (Group 1)</b>					
Study Site	Entebbe Masaka	1 6.47 (3.15-13.28)	P<0.001	1 3.87 (1.20-12.5)	0.02
Marital Status	Currently Married Widowed Separated / Divorced Single	1 0.85 (0.40-1.82) 0.67 (0.33-1.38) 0.18 (0.02-1.35)	P=0.15		
Religion	Catholic Protestant Muslim Other	-	-	1 0.33 (0.09-1.25) 1.33 (0.40-4.46) 0.46 (0.05-4.22)	0.24
<b>HIV associated clinical factors (Group 2A)</b>					
Weight (kg)	Per one s.d. increase	-	-	1.47 (0.93-2.33)	P=0.10
<b>Vulnerability/protective factors (Group 2B)</b>					
Depression score at baseline (CES-D)	0-15 16-23 24-60	1 2.54 (1.25-5.16) 2.18 (0.86-5.55)	P=0.03	1 1.79 (0.45-7.09) 4.02 (1.15-14.02)	P=0.05
<b>Stress factors (Group 2C)</b>					
Stress Score	Per one s.d. increase	1.52 (1.15-2.00)	P=0.003	-	-