

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Aggarwal, A; Lewis, D; Charman, SC; Mason, M; Clarke, N; Sullivan, R; van der Meulen, J (2017) Determinants of Patient Mobility for Prostate Cancer Surgery: A Population-based Study of Choice and Competition. *European urology*. ISSN 0302-2838 DOI: <https://doi.org/10.1016/j.eururo.2017.07.013>

Downloaded from: <http://researchonline.lshtm.ac.uk/4209919/>

DOI: [10.1016/j.eururo.2017.07.013](https://doi.org/10.1016/j.eururo.2017.07.013)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

Table 1 – Characteristics of 19,256 men undergoing radical prostatectomy between 2010 and 2014 in the English National Health Service.

	Number	%
Age (years)		
<50	702	3.7
50-59	5114	26.6
60-64	5305	27.6
65-69	5973	31.0
≥70	2162	11.0
Cancer severity		
Advanced	50	0.3
Locally advanced	6373	43.9
Intermediate localised	7613	52.4
low risk localised	488	3.4
Insufficient staging information (n=4732)		
Number of comorbidities		
0	17,821	92.6
≥1	1435	7.4
Index of Multiple Deprivation (national quintiles)		
1 (least deprived)	5312	28.0
2	4744	24.6
3	3975	20.4
4	2980	15.0
5 (most deprived)	2245	11.7
Urban- rural classification¹		
Urban	14,685	76.3
Rural	4571	23.7

Notes:

1. See text for definition