

Table 2: Characteristics of infants at 12 months¹

Variable	<i>Infant data</i>		P Value	Difference (95% CI)
	Intervention group (n=36)	Control group (n=36)		
Mean gestational age, week	39.1 (± 1.0)	39.3 (± 1.0)	0.47	-0.18 (-0.67; 0.31)
Male, n	19 (52.8%)	18 (50.0%)	0.81	
Female, n	17 (47.2%)	18 (50.0%)		
Birth Weight, g	3.1 (± 0.28)	3.1 (± 0.36)	0.34	-0.73 (-0.22; 0.079)
CF 4-5 m, n	17 (47.2%)	16 (44.5%)	0.81	
CF \geq 6 m, n	19 (52.8%)	20 (55.5%)		
Mean weight at 12 m, g	8.8 (± 0.94)	9.2 (± 1.1)	0.12	-0.38 (-0.86; 0.10)
Mean length at 12 m, cm	72.3 (± 2.6)	72.4 (± 2.7)	0.87	-0.10 (-1.35; 1.14)
WAZ at 6 m	-0.58 (± 0.9)	-0.20 (± 1.0)	0.11	-0.37 (-0.84; 0.09)
LAZ at 6 m	-1.09 (± 1.1)	-1.06 (± 1.0)	0.92	-0.27 (-0.53; 0.48)
WLZ at 6 m	-0.52 (± 0.85)	0.42 (± 0.93)	0.03	-0.47 (-0.88; -0.05)
HCZ at 6 m	0.22 (± 0.92)	0.30 (± 0.90)	0.73	-0.75 (0.21; -0.50)

¹ The definitions used in the table are: CF: complementary feeding practices, WAZ: weight-for-age Z-scores, LAZ: length-for-age Z-scores, WLZ: weight-for-length Z-scores, HCZ: head circumference-for-age Z score