

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Ferreira, JP; Girerd, N; Gregson, J; Latar, I; Sharma, A; Pfeffer, MA; McMurray, JJV; Abdul-Rahim, AH; Pitt, B; Dickstein, K; +3 more... Rossignol, P; Zannad, F; High-Risk Myocardial Infarction Database Initiative; (2018) Stroke Risk in Patients With Reduced Ejection Fraction After Myocardial Infarction Without Atrial Fibrillation. *Journal of the American College of Cardiology*, 71 (7). pp. 727-735. ISSN 0735-1097 DOI: <https://doi.org/10.1016/j.jacc.2017.12.011>

Downloaded from: <http://researchonline.lshtm.ac.uk/4646682/>

DOI: <https://doi.org/10.1016/j.jacc.2017.12.011>

Usage Guidelines:

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

Figure 1. Model calibration plot: % of observed vs predicted events at 3-years by categories of stroke risk score




Figure 2. Kaplan-Meier failure estimates curve by stroke risk score categories for the outcome of stroke with death as competing risk

