



2005 cholera epidemic in Senegal




- ▶ Peak end of March related to **pilgrimage in Touba**
- ▶ Flare and spread due to **overcrowding and travelling pilgrims**
- ▶ Peak in Dakar in autumn related to **rainfall and floods**
- ▶ **Scenarios of reduced transmission** during pilgrimage



Overview


- ▶ We analyze a dataset of **mobile phone call records** in Senegal and **extract human mobility fluxes** over a period of one year.
- ▶ The fluxes are directly used in a **spatially explicit, mechanistic epidemiological model** of the **2005 cholera epidemic in Senegal**.
- ▶ The spread of the epidemic was boosted by a **mass gathering of 3 million pilgrims**.
- ▶ This **crucial effect** could only be accounted for thanks to the first-order information about **origin, destination and number of travellers per day** not present in other data sources.

Mobile phone data analysis



- ▶ 150,000 mobile phone users over the year 2013.
- ▶ Determination of **home district** of each user using **calls made at night**
- ▶ **Time spent** at node j by users with home node i proportional to **number of calls** they made at j .
- ▶ $Q_{ij}(t)$ contains the **average fraction of time** spent by users of note i at node j during day t .

Cholera model



$$\begin{aligned} \frac{dS_i}{dt} &= \mu(H_i - S_i) - \mathcal{O}_i(t)F_i(t)S_i + \rho R_i \\ \frac{dI_i}{dt} &= \sigma \mathcal{O}_i(t)F_i(t)S_i - (\gamma + \mu + \alpha) I_i \\ \frac{dR_i}{dt} &= \gamma I_i + (1 - \sigma) \beta_i(t) \mathcal{O}_i(t)F_i(t)S_i - (\rho + \mu) R_i \\ \frac{dB_i}{dt} &= -\mu_B B_i + \frac{\theta}{H_i} [1 + \lambda J_i(t)] \mathcal{O}_i(t)G_i(t) \end{aligned}$$

- ▶ Spatially explicit SIRB type model
- ▶ Fluxes from mobile phone data employed directly within the model to account for pathogen spread
- ▶ Overcrowding effect $\mathcal{O}_i(t)$
- ▶ Rainfall
- ▶ 6 calibration parameters

$$\begin{aligned} \mathcal{O}_i(t) &= \exp\left(\frac{\omega}{H_i} \sum_{j=1}^N Q_{ji}(t)H_j\right) \\ F_i(t) &= \beta \sum_{j=1}^N Q_{ij}(t) \frac{B_j}{K + B_j} \\ G_i(t) &= \sum_{j=1}^N Q_{ji}(t)I_j. \end{aligned}$$

Targeted interventions

