

# Genome Sequences of Three *Salmonella enterica* subsp. *enterica* Serovar *Infantis* Strains from Healthy Broiler Chicks in Hungary and in the United Kingdom

Ferenc Olasz,<sup>a</sup> Tibor Nagy,<sup>a</sup> Móni Szabó,<sup>a</sup> János Kiss,<sup>a</sup> Ama Szmolka,<sup>b</sup> Endre Barta,<sup>a</sup> Andries van Tonder,<sup>c</sup> Nicholas Thomson,<sup>d</sup> Paul Barrow,<sup>e</sup> Béla Nagy<sup>b</sup>

Agricultural Biotechnology Institute of the National Agricultural Research and Innovation Centre, Gödöllő, Hungary<sup>a</sup>; Institute for Veterinary Medical Research, Centre for Agricultural Research, Hungarian Academy of Sciences, Budapest, Hungary<sup>b</sup>; Nuffield Department of Medicine, University of Oxford, Peter Medawar Building for Pathogen Research, Oxford, United Kingdom<sup>c</sup>; Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, United Kingdom<sup>d</sup>; School of Veterinary Medicine and Science, University of Nottingham, Sutton Bonington, Leicestershire, United Kingdom<sup>e</sup>

F.O. and T.N. contributed equally to this work.

**The genome sequences of three strains of *Salmonella enterica* subsp. *enterica* serovar *Infantis* isolated from broiler chickens in 1994 and 2004 in Hungary and in the 1980s in the United Kingdom are reported here. A sequence comparison should improve our understanding of the evolution of the genome and spread of *S. Infantis* in poultry.**

Received 11 December 2014 Accepted 5 January 2015 Published 12 February 2015

**Citation** Olasz F, Nagy T, Szabó M, Kiss J, Szmolka A, Barta E, van Tonder A, Thomson N, Barrow P, Nagy B. 2015. Genome sequences of three *Salmonella enterica* subsp. *enterica* serovar *Infantis* strains from healthy broiler chicks in Hungary and in the United Kingdom. *Genome Announc* 3(1):e01468-14. doi:10.1128/genomeA.01468-14.

**Copyright** © 2015 Olasz et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ferenc Olasz, [olasz@abc.hu](mailto:olasz@abc.hu), or Paul Barrow, [paul.barrow@nottingham.ac.uk](mailto:paul.barrow@nottingham.ac.uk).

*Salmonella enterica* subsp. *enterica* serovar *Infantis* spreads among broilers in Hungary (1) and elsewhere in Europe but not in the United Kingdom (2). Here, we present the genome sequences of three *Salmonella* *Infantis* strains isolated from chickens from the 1980s to the 2000s.

Fragment libraries of 500 bp were prepared from the total DNA of strains SI 69/94 and SI 54/04 (Hungary), and the 2 × 100-bp Illumina paired-end genome sequencing was performed by Aros Applied Biotechnology (Aarhus, Denmark) as a custom service, using the Illumina HiSeq 2000 platform. The numbers of reads are 34.4 million for SI 69/94 and 95 million for SI 54/04. The estimated coverages of the whole genome are 600× and 1,700×, respectively. The whole genome of *S. Infantis* strain 1326/28 (United Kingdom) (3, 4), sequenced to a coverage depth of 9× from M13mp18 (insert size, 1.4 to 2 kb) and pUC18 (insert size, 2.2 to 4.2 kb) small-insert libraries using BigDye Terminator chemistry on ABI3700 automated sequencers. The end sequences from a larger-insert plasmid (pBACe3.6, 12- to 30-kb insert size) libraries were used as a scaffold. The sequence was assembled, finished, and annotated, as described previously (5), using the program Artemis (6) to collate data and facilitate annotation.

The estimated coverage of the subsets of reads was adjusted to 30× for both Hungarian strains, and these were *de novo* assembled using Mira version 3.9.15 (7). The G+C content appeared to be same for all three strains (52.29%). The draft sequence of strain SI 54/04 contains additional scaffolds that cannot be aligned to those of SI 69/94 and the chromosome of 1326/28, suggesting the presence of a ca. 277-kbp plasmid in strain SI 54/04.

Randomly selected subsets of reads were also applied for the reference genome assisted assembly, using the assembled United Kingdom strain 1326/28 as a reference backbone. The chromosome

lengths of strains 1326/28, SI 69/94, and SI 54/04 have been determined to be 4,710,675, 4,710,832, and 4,710,839 bp, respectively.

The assembled genome sequences were submitted to the RAST annotation server (8). We set the taxon to *S. enterica*, the genetic code to 11 (*Archaea*, *Bacteria*), the sequence method to “other,” the coverage to >8×, and “automatically fix errors” to “no.” For SI 69/94, SI 54/04, and 1326/28, we obtained the following data: 4,664, 4,623, and 4,671 annotated genes, respectively; 170, 172, and 168 tRNAs, respectively; and 44 rRNAs in all three strains.

A comparison of the chromosomal nucleotide sequences revealed that the three strains exhibit marked similarity (minimum, 99.94%; maximum, 99.97%). Comparing 1326/28 to the published Israeli strains of human origin (9), 119944 (GenBank assembly ID GCA\_000506925.1) and 335-3 (GenBank assembly ID, GCA\_000506945.1), 92.63% and 100% similarities were observed, respectively. The genome sequences of the three *S. Infantis* strains revealed a strikingly limited similarity (about 83%) to the SARB27 sequences (GenBank accession no. CM001274), identified earlier as *S. Infantis* (10), but indicating an unusual evolutionary distance.

**Nucleotide sequence accession numbers.** The draft genome sequences of SI 69/94 and SI 54/04 and the assembled and annotated genome sequence of 1326/28 have been deposited in the NCBI GenBank database under the accession numbers [JRXB000000000](https://www.ncbi.nlm.nih.gov/nuclink/JRXB000000000), [JRXC000000000](https://www.ncbi.nlm.nih.gov/nuclink/JRXC000000000), and [LN649235](https://www.ncbi.nlm.nih.gov/nuclink/LN649235), respectively.

## ACKNOWLEDGMENTS

This work is supported by the Hungarian Scientific Research Fund (OTKA) K101546 and by the János Bolyai Stipend of the Hungarian Academy of Sciences to A. Szmolka, J. Kiss, and M. Szabó. N. Thomson and A. van Tonder were supported by Wellcome Trust grant 098051.

We thank Ilona Könczöl for technical assistance.

## REFERENCES

1. Nógrády N, Tóth A, Kostyák A, Pászti J, Nagy B. 2007. Emergence of multidrug-resistant clones of *Salmonella* Infantis in broiler chickens and humans in Hungary. *J Antimicrob Chemother* 60:645–648. <http://dx.doi.org/10.1093/jac/dkm249>.
2. Nógrády N, Király M, Davies R, Nagy B. 2012. Multidrug resistant clones of *Salmonella* Infantis of broiler origin in Europe. *Int J Food Microbiol* 157:108–112. <http://dx.doi.org/10.1016/j.ijfoodmicro.2012.04.007>.
3. Barrow PA, Huggins MB, Lovell MA. 1994. Host specificity of *Salmonella* infection in chickens and mice is expressed *in vivo* primarily at the level of the reticuloendothelial system. *Infect Immun* 62:4602–4610.
4. Barrow PA, Lovell MA. 1989. Invasion of Vero cells by *Salmonella* species. *J Med Microbiol* 28:59–67. <http://dx.doi.org/10.1099/00222615-28-1-59>.
5. Parkhill J, Wren BW, Mungall K, Ketley JM, Churcher C, Basham D, Chillingworth T, Davies RM, Feltwell T, Holroyd S, Jagels K, Karlyshev AV, Moule S, Pallen MJ, Penn CW, Quail MA, Rajandream MA, Rutherford KM, van Vliet AH, Whitehead S, Barrell BG. 2000. The genome sequence of the foodborne pathogen *Campylobacter jejuni* reveals hypervariable sequences. *Nature* 403:665–668. <http://dx.doi.org/10.1038/35001088>.
6. Berriman M, Rutherford K. 2003. Viewing and annotating sequence data with Artemis. *Brief Bioinform* 4:124–132. <http://dx.doi.org/10.1093/bib/4.2.124>. PubMed.
7. Chevreur B, Wetter T, Suhai S. 1999. Genome sequence assembly using trace signals and additional sequence information, p 45–56. *In* Computer science and biology. Proceedings of the German Conference on Bioinformatics, GCB '99. GCB, Hannover, Germany.
8. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
9. Aviv G, Tsyba K, Steck N, Salmon-Divon M, Cornelius A, Rahav G, Grassl GA, Gal-Mor O, Aviv G, Tsyba K, Steck N, Salmon-Divon M, Cornelius A, Rahav G, Grassl GA, Gal-Mor O. 2014. A unique megaplasmid contributes to stress tolerance and pathogenicity of an emergent *Salmonella enterica* serovar Infantis strain. *Environ Microbiol* 16:977–994. <http://dx.doi.org/10.1111/1462-2920.12351>.
10. Boyd EF, Wang FS, Beltran P, Plock SA, Nelson K, Selander RK. 1993. *Salmonella* reference collection B (SARB): strains of 37 serovars of subspecies I. *J Gen Microbiol* 139(Pt 6):1125–1132. <http://dx.doi.org/10.1099/00221287-139-6-1125>.