

Summarizing and Communicating on Survival Data According to the Audience: A Tutorial on Different Measures Illustrated with Population-Based Cancer Registry Data [Corrigendum]

Belot A, Ndiaye A, Luque-Fernandez MA, et al. *Clin Epidemiol*. 2019;11:53–65.

Page 57, Number of life years lost (NLYL) section, left column, the sentence "where the quantity $1-F_P(t)$ can be replaced by $S_P(t)$, ie, the classical survival function using the population mortality rates λ_P " placed immediately after equation 14 indicates that the quantity $1-F_P(t)$ could be replaced by $S_P(t)$. This is wrong as the one-to-one relation-

ship between hazard and risk applies only in all-cause mortality setting but not in competing risks settings (such as the cause-specific or relative survival setting). Indeed, F_P (t) depends on both hazards (cancer and other causes) through the overall survival S(t).

The authors therefore request the reader to ignore this sentence and apologize for this error.

Clinical Epidemiology

Publish your work in this journal

Clinical Epidemiology is an international, peer-reviewed, open access, online journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification,

 $\textbf{Submit your manuscript here:} \ \texttt{https://www.dovepress.com/clinical-epidemiology-journal}$

systematic reviews, risk & safety of medical interventions, epidemiology & biostatistical methods, and evaluation of guidelines, translational medicine, health policies & economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.



