

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Rehman, AM; Gladstone, BP; Verghese, VP; Muliyl, J; Jaffar, S; Kang, G (2009) Chronic growth faltering amongst a birth cohort of Indian children begins prior to weaning and is highly prevalent at three years of age. *Nutrition journal*, 8. p. 44. ISSN 1475-2891 DOI: <https://doi.org/10.1186/1475-2891-8-44>

Downloaded from: <http://researchonline.lshtm.ac.uk/4674/>

DOI: [10.1186/1475-2891-8-44](https://doi.org/10.1186/1475-2891-8-44)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

Additional file 1: Imputation of missing values

Our primary outcome was the odds of stunting at our study endpoint, 36 months of age. Of the 452 children that began follow-up, 74 dropped out of the study and 5 died. At our study endpoint, 373 children were still under follow-up. We examined the associations between not completing follow-up and each of the potential risk factors (defined *a priori*) for our multivariate model and found associations with mother's education level and the age at introduction of complementary foods. We included these predictors of missingness in our multivariate model to give us an analysis that assumes our dropouts were so-called "missing at random" (MAR) [1].

In our "completers" cohort of 373 children we had missing values in our outcome variable and in three covariates (Table 1). Our complete case analysis excluded 96 (26%) children.

Table 1 – Variables in the multivariate logistic regression model for the odds of stunting at 36 months and numbers of missing and non-missing values

Variables	Number of missing values	Number of non-missing values
Outcome		
Stunting at 36 months (stunted36)	42	331
Risk factors		
Low birth weight (lbw)	8	365
Beedi making in the household (beedi)	0	373
Mother's height (mheight)	32	341
Growth faltering at six months (growfalt6)	28	345
Not first born (firstborn)	0	373
SES Class (ses)	0	373
Sex (sex)	0	373
Age introduced complementary foods (compfood)	0	373
Mother's education attainment (medu)	0	373
Mother's age (mage)	0	373
Duration of major illness in the first three years of life (mjill)	0	373

To use more of our data, we used standard multiple imputation methods to create 10 imputations of all missing values in our "completers" cohort [2]. We used all 452 children to create the imputed values. In addition to the list of variables in our multivariate model (shown in Table 1), we included (by default) the variables associated with missingness but also included other variables related to our missing variables as suggested by van Buuren *et al.* [3] (see Table 2 for list of variables used in each imputation model). We included the height-for-age z-score from 35 months of age in the imputation model for stunting at 36 months, which was not missing in 37/42 children with missing values of stunting at 36 months of age. We included the weight of the child from 1 month of age in the imputation model for low birth weight, which was not missing in 7/8 children with missing values for low birth weight. We included no additional variables in the imputation model for mother's height. Finally, we amended the core list of variables we included in the imputation model for growth faltering at six months to include illness in the first six months of life and its interaction with gender (as these were predictors in the multivariate model with

growth faltering at six months as the outcome) and also the height-for-age, weight-for-age and weight-for-height z-scores from 5 months of age which were not missing in any children with missing values for growth faltering at 6 months.

Table 2 – Details of multiple imputation models for the imputing of missing values

Variables with missing values	Model used in imputation	Variables used in imputation model
Stunting at 36 months	Logistic regression	lbw, beedi, mheight, growfalt6, firstborn, ses, sex, compfood, medu (two dummy variables), mage, mjill (two dummy variables), height-for-age z-score at 35 months
Low birth weight	Logistic regression	stunted36, beedi, mheight, growfalt6, firstborn, ses, sex, compfood, medu (two dummy variables), mage, mjill (two dummy variables), weight at 1 month of age
Mother’s height	Logistic regression	stunted36, lbw, beedi, growfalt6, firstborn, ses, sex, compfood, medu (two dummy variables), mage, mjill (two dummy variables),
Growth faltering at six months	Logistic regression	stunted36, lbw, beedi, mheight, firstborn, ses, sex, compfood, medu (two dummy variables), mage, all illness in first six months of life (two dummy variables), interaction between sex and all illness (two dummy variables), height-for-age z-score at 5 months, weight-for-age z-score at 5 months, weight-for-height z-score at 5 months

The imputed values in our outcome, stunting at 36 months were more often “yes” the child was stunted, ranging from 25/42 to 33/42 missing values among the 10 imputation models (Table 3). Almost all missing values of low birth weight were imputed to be “no” the child did not have low birth weight, ranging from 7/8 to 8/8 (Table A.3). More often mother’s height was imputed to be ≥ 150 cm than < 150 cm, ranging from 17/32 to 26/32 (Table 3). Growth faltering at six months was more evenly split with imputed values of “no” growth faltering ranging between 13/28 to 20/28 missing values (Table 3).

For our secondary outcome of growth faltering at six months, our “completers” cohort of 403 children still under follow-up had missing values in the outcome variable (32) and in two covariates, low birth weight (10) and mother’s height (50). Our complete case analysis excluded 85 (21%) children. We investigated factors related to not completing follow-up to six months of age and found associations with duration of illness during the first six months (chi-square p-value <0.001) and the age at introduction of complementary foods (log-rank test p-value=0.0013). These predictors of missingness were included in our multivariate and imputation models by default. Distributions of imputed values were a similar pattern as for stunting at 36 months (Table 4).

Table 3 – Imputed values in each of 10 imputation models for the odds of stunting at 36 months

		IMPUTATION									
Variable	Total missing values	1	2	3	4	5	6	7	8	9	10
Stunting at 36 months	42										
No		9	11	12	11	15	11	14	10	12	17
Yes		33	31	30	31	27	31	28	32	30	25
Low birth weight	8										
No		8	8	7	8	7	7	8	8	7	7
Yes		0	0	1	0	1	1	0	0	1	1
Mother's height	32										
>=150 cm		23	18	23	21	17	18	23	26	25	18
<150 cm		9	14	9	11	15	14	9	6	7	14
Growth faltering at six months	28										
No		20	15	14	15	15	17	18	17	13	19
Yes		8	13	14	13	13	11	10	11	15	9

Table 4 – Imputed values in each of 10 imputation models for the odds of growth faltering at 6 months

		IMPUTATION									
Variable	Total missing values	1	2	3	4	5	6	7	8	9	10
Growth faltering at six months	32										
No		21	17	14	16	17	19	21	19	14	20
Yes		11	15	18	16	15	13	11	13	18	12
Low birth weight	10										
No		9	10	9	9	8	8	10	10	9	8
Yes		1	0	1	1	2	2	0	0	1	2
Mother's height	50										
>=150 cm		32	29	32	35	30	27	37	38	37	31
<150 cm		18	21	18	15	20	23	13	12	13	19

REFERENCES

1. Rubin DB: **Multiple imputation after 18+ years**. *Journal of the American Statistical Association* 1996, **91**(434):473-489.
2. Royston P: **Multiple imputation of missing values: update of ice**. *Stata Journal* 2005, **5**:527-536.
3. van Buuren S, Boshuizen HC, Knook DL: **Multiple imputation of missing blood pressure covariates in survival analysis**. *Stat Med* 1999, **18**(6):681-694.