

# ***Management of colorectal cancer explains differences in one-year relative survival between France and England for patients diagnosed 1997-2004.***

## Authors

Dejardin O <sup>1</sup>, Rachet B <sup>2</sup>, Morris E <sup>4&5</sup>, Bouvier V <sup>1</sup>, Jooste V <sup>3</sup>, Haynes R <sup>6</sup>, Coombes EG <sup>6</sup>, Forman D <sup>5&7</sup>, Jones AP <sup>6</sup>, Bouvier AM <sup>3</sup>, Launoy G <sup>1</sup>

## Author's affiliation

- 1- University Hospital of Caen, EA 3936 UCBN University of Caen, U1086 INSERM « Cancers & preventions », France
- 2- London School of Hygiene and Tropical medicine LSHTM, Cancer Research UK Cancer Survival Group, London, WC1E 7HT, UK
- 3- Digestive Cancer Registry of Burgundy; CHU Dijon; INSERM, U866, Dijon, F-21079, France; University Bourgogne, Dijon, F-21079, France.
- 4- Cancer Epidemiology Group, Leeds Institute of Molecular Medicine, University of Leeds, Level 6, Bexley Wing, St James's University Hospital, Leeds, LS9 7TF, UK
- 5- Northern and Yorkshire Cancer Registry & Information Service (NYCRIS), Level 6, Bexley Wing, St James's University Hospital, Leeds, LS9 7TF UK
- 6- School of Environmental Sciences, University of East Anglia, Norwich, UK
- 7- International Agency for Research against Cancer (IARC), Section of Cancer Information, Lyon, France

Corresponding author Launoy Guy, « Cancers & preventions » U1086 INSERM; Faculty of Medicine, Avenue de la Côte de Nacre 14032 Caen Cedex, France

Tel +33 (0)2 31 45 86 01

Mail: [guy.launoy@unicaen.fr](mailto:guy.launoy@unicaen.fr)

Word count:

Keywords: Colorectal neoplasm; Survival; France; England; Population-based study

Competing interests: None declared

## ***Summary***

### Objectives

Few international population-based studies have provided information on potential determinants of international disparities in cancer survival. This population-based study was aimed to identify the principal differences in disease characteristics and management that accounted for previously observed poorer survival in English compared to French patients with colorectal cancer

### Methods

The study population comprised all cases of colorectal cancer diagnosed between 1997 and 2004 in the areas covered by three population-based cancer registries in France and one in England (N= 40 613). To investigate the influence of clinical and treatment variables on survival, we applied multivariable excess hazard modelling based on generalised linear models with Poisson error.

### Results

Poorer survival for English patients was primarily due to a larger proportion dying within the first year after diagnosis. After controlling for inter-country differences in the use of chemotherapy and surgical resection with curative intent, country of residence was no-longer associated with 1-year survival for advanced colon cancer patients (EHR = 0.99 (0.92-1.01),  $p = 0.095$ ). Longer-term (2-5 years) excess hazards of death for colon and rectal cancer patients did not differ between France and England.

### Conclusion

This study suggests that difference in management close to diagnosis of colon and rectum cancer is related to difference in survival observed between France and England. Every efforts (Collection and standardization of additional variables such as co-morbidity) to investigate the reasons for this disparities in management between these two countries, and more generally across Europe, should be encouraged.

## ***Introduction***

Survival for patients with colorectal cancer varies notably between European countries. As reported by the CONCORD study (Coleman *et al*, 2008) in 2008, 5-year relative survival for colorectal cancer in France was 55.6% for men and 61.5% for women, which constituted the highest survival in Western Europe and second best in the world for women. By contrast, in England five-year relative survival was worse than in comparable European countries at 42.3% for men and 44.7% for women. The improvement in 5-year relative survival between 1988-1990 and 2000-2002 was comparable in both countries (+10.2% in France Vs. +10.4% in England)(Brenner *et al*, 2011). Under the scenario of equal survival in England compared to the mean European 5-year survival, between 6600 and 7500 deaths would have been avoided each year for patients diagnosed between 1985-1999 (Abdel-Rahman *et al*, 2009).

Some authors have suggested that these international disparities in cancer survival, in particular the poorer survival in England, may not be due to actual differences in survival, but rather due to differences in cancer registration modalities and quality (Beral & Peto, 2010). However, a recent study highlighted that, even under the hypothesis of extreme incorrect registration (either date of recurrence instead of date of diagnosis or under-registration of long survivors), differences in cancer registration would explain very little of the observed disparities in survival (Woods *et al*, 2011). Another study suggested that the magnitude of error in survival time due to incorrect case completeness was less than 1% for patients diagnosed with colorectal cancer in England (Møller *et al*, 2011). Moreover, registries included in the EURO CARE studies are broadly comparable in terms of registration methods, especially in terms of microscopic verification and the prevalence of death certificate only records (Berrino *et al*, 2007).

The reasons behind the international differences in survival are not well understood and many factors have been argued as potentially influential. For example, there is some evidence to show that the lower survival in England compared to Scandinavian countries is due to a high number of deaths, particularly amongst elderly patients, in the first three months following diagnosis (Engholm *et al*, 2007; Folkesson *et al*, 2009; Morris *et al*, 2011). This would suggest that a greater proportion of the population in England present with rapidly fatal disease than elsewhere in Europe. This may be due to more English cases presenting with advanced disease at diagnosis, concomitant morbidity or, perhaps, experiencing a different quality of care to those in other European countries(Gatta *et al*, 2010).

International comparisons of survival for patients with cancer are important for the planning and provision of national health services. However, with the notable exception of “EUROCARE high-resolution” studies (Ciccolallo *et al*, 2005; Gatta *et al*, 2000), very few international population-based studies have provided information on potential determinants of international disparities in cancer survival. The aim of this population-based study was to identify the principal differences in disease characteristics and cancer management that accounted for the difference in survival observed between France and England for patients with colorectal cancer.

## ***Materials & Methods***

### ***Population***

The study population comprised all cases of colorectal cancer (ICD10: C18.0 to C20.9) diagnosed between 1997 and 2004 in the areas covered by three population-based cancer registries in France (Calvados, Côte d'Or and Saône et Loire: 3% of the whole national population of France) and one in England (the Northern and Yorkshire Cancer Information Service (NYCRIS), which covers 13.3% of the national population)(Table 1). The completeness and data quality of the included registries are regularly assessed by the International Agency for Research on Cancer (IARC) or by European Network of Cancer Registries (ENCR). The final population study included 40 613 patients (7891 from France and 32722 from England). Over the period study, 919 patients in England had a 0 days survival (2.8%) versus 18/7891 (0.22%) in France.

### ***Data***

Date of birth, sex and year of diagnosis were known for all patients. Topography was classified as right colon (C18.0, C18.1, C18.2, C18.3 and C18.4); left colon (C18.5, C18.6 and C18.7); unknown location of colon cancer (C18.8 and C18.9); rectosigmoid junction (C19.9) and rectum (C20.9) (Fritz *et al*, 2000). Stage was coded using Duke's classification: Duke's A: Limited to mucosa; Duke's B: Penetrating through muscularispropria; Duke's C: lymph nodes involved; "Duke's D": a least one metastasis (Dukes, 1932). The unstaged category included: non-resected patients with no clinical evidence of metastases at diagnosis, resected patients but for whom the registry did not capture the stage, and patients who received potentially downstaging radiotherapy or chemoradiotherapy prior to surgery. Localized cancers were defined by Duke's A & B and advanced cancers were defined by Duke's C & D. stage at diagnosis was unknown for almost 18% and 7% of English and French patients respectively (proportions are comparable to previous studies)(Jones *et al*, 2009).

Information was collected on the type of treatment: surgical resection with curative intent within 6 months since diagnosis (Coded as: yes or no), chemotherapy (yes or no) and radiotherapy (yes or no). Palliative chemotherapy and palliative radiotherapy were not recorded in NYCRIS. French registries captured information on all chemotherapy and radiotherapy administered irrespective of intent.

Survival time was defined as the time duration between the date of diagnosis and the earlier of date of death or date of last information about vital status or the end of the study period on 31/12/2008.

### ***Statistical analysis***

Estimation of up-to-5-year relative survival was based on the Ederer-II approach using the user-written Stata command *strs* (Estimating and modelling relative survival, available at <http://www.pauldickman.com/>). To investigate the influence of clinical and treatment variables on survival, we applied multivariable excess hazard modelling based on generalised linear model with Poisson error (Dickman *et al*, 2004). Excess hazard ratio (HER) were calculated with 95% confident intervals (Dickman *et al*, 2004). Time since diagnosis was split in intervals as following: 0-3 months, 3-6 months, 6 months-1 year, 1-2 years, 2-3 years and 4-5 years. To take into account the influence of treatment, survival analyses were then stratified according to cancer tumour site and stage. All analyses were computed using STATA 12.1 software using a publicly available procedure (Estimating and modelling relative survival, available at <http://www.pauldickman.com/>). In the absence of reliable information on the cause of death among the cancer patients, cancer-related survival is commonly estimated by relative survival approach which removes from the observed, all-cause mortality the expected (“background”) mortality. Background mortality was provided by life tables stratified according to age, sex, year of diagnosis, and administrative area (Government Office Region for England and “Department” for France).

Multiple imputations by chained equations were performed to take into account missing values (Little & Rubin, 2002) (missing values are presented in Table 2). The imputation model incorporated the variables used in the analytical models (survival time, vital status, age, sex, topography, stage, year of diagnosis and treatments) as recently recommended (Nur *et al*, 2010). The imputation model was stratified according to country. Iterations were conducted to create 20 completed datasets, and the estimates were combined according to the Rubin rules (Rubin, 1987). Multiple imputation models were conducted using conjointly STATA 12.1 module for imputation (StataCorp. 2011. Stata: Release 12. Statistical Software. College Station, TX: StataCorp LP) and the user-written Stata command *ice* (Royston, 2009). All results presented in the next section are based on multiple imputations. In order to test the robustness of our results, all analyses were repeated on complete case analysis since the relevance of this method depends on the missingness mechanism.

## ***Results***

### ***Management***

The distribution of year of diagnosis and sex did not differ between the cancer registries (Table 1), although the proportion of right colon cancers was higher in England than in France. Distribution of stage before and after multiple imputations is presented in Table 2. For all cancer localizations, the proportion of patients diagnosed at “Duke’s D” increased in both countries after multiple imputations from 27.3% to 30.0% in France and from 27.9% to 32.7% in England for patients diagnosed with colon cancer, and from 24.0% to 31.6% in France and from 23.2% to 28.4% in England for patients diagnosed with rectal cancer. Treatment modalities (chemotherapy, radiotherapy and surgical resection with curative intent within 6 months since diagnosis) differed between the countries. Such surgical resections were performed more frequently in France, being used in 83.6% of patients in France vs. 73.1% in England. Whatever stage at diagnosis or tumour localization, the rate of patients receiving a surgical resection with curative intent within 6 months was higher in France than in England. This difference was limited for localized colon cancer (98.9% in France Vs. 96.1% in England for stage A; 99.0% in France vs. 96.3% in England for stage B) and more pronounced for advanced colon cancer (97.6% in France vs. 87.8 in England for stage C; 52.4% in France vs. 32.3% in England for stage ‘D’). Concerning rectum cancer, the different in the rate of patients with surgical resection within 6 months increased with stage (97.6% in France vs. 93.2% in England for stage A; 94.6% in France vs. 87.6% in England for stage B; 86.7% in France vs. 80.8 in England for stage C; 32.6% in France vs. 22.8% in England for stage ‘D’).

### ***Survival***

The association between country of residence and relative survival was not constant over time since diagnosis. In survival analysis, the interaction between time since diagnosis and country of residence was statistically significant ( $p < 0.001$ ), with the gain in survival for French patients decreasing with time since

diagnosis. Therefore, multivariable models were stratified according to time since diagnosis (first year since diagnosis vs. first to 5th years since diagnosis).

***One-Year survival analyses for patients diagnosed with colon cancer***

Patients diagnosed with colon cancer living in France had a better prognosis than those living in England for the first year after diagnosis (EHR = 0.70 (0.66-0.75)) (Table 3, model 1). After adjustment for sex, age and period of diagnosis, the higher survival for French patients remained unchanged (EHR = 0.70 (0.65-0.74)) (Table 3, model 2). After successive adjustment for tumour site (Table 3, model 3) and stage at diagnosis (Table 3, model 4), patients living in France still had a higher 1-year survival than those living in England (EHR = 0.68 (0.64-0.74)).

To investigate the influence of treatment, survival analyses were conducted separately on localized cancer (Duke's A and Duke's B) and advanced cancer (Duke's C and "Duke's D"). In unadjusted analyses, French patients diagnosed with a localized colon cancer had a better survival than those diagnosed in England (EHR= 0.77 (0.65-0.93)) (Table 3, model 5a). Differences in survival decreased after adjusting for surgical resection with curative intent within 6 months, but country of residence was still associated with 1-year survival (EHR = 0.82 (0.68-0.99),  $p = 0.038$ ) (Table 3, model 6a). French patients diagnosed with an advanced colon cancer also had better survival than those diagnosed in England (EHR= 0.67 (0.63-0.72)) (Table 3, model 5b) in unadjusted analyses. After adjustment for surgery with curative intent, country of residence remained associated with 1-year survival for patients diagnosed with an advanced colon cancer (EHR = 0.79 (0.74-0.85)) (Table 3, model 6b). After taking into account the use of chemotherapy, country of residence was no more associated with 1-year survival (EHR = 0.99 (0.92-1.01),  $p = 0.095$ ) (Table 3, model 7b). After considering the interaction, country of residence was associated with survival only in the presence of chemotherapy (EHR= 0.99 (0.92- 1.07),  $p=0.918$  in absence of chemotherapy and EHR=0.81 (0.69-0.94),  $p=0.008$  in presence of chemotherapy) (Table 3, model 8b).

### ***One-Year survival analyses for patients diagnosed with rectum cancer***

Similarly, after adjustment for age, sex, period of diagnosis and stage, French rectal cancer patients had a better 1-year-survival than English patients (EHR = 0.76 (0.68-0.86)) (Table 3, model 11). Country of residence was significantly associated with 1-year survival for patients diagnosed with a localized cancer (EHR = 0.70 (0.49-0.99),  $p = 0.048$ ) (Table 3, model 11a). But, again, after further adjustment for surgical resection with curative intent, country of residence was no longer statistically significant (EHR = 0.74 (0.51- 1.08)) (Table 3 model 12a).

When the analysis was restricted to include patients with advanced rectal cancer only, 1-year survival remained higher in France (EHR = 0.69 (0.61-0.77)) (Table 3, model 11b). After taking into account surgical resection with curative intent and radiochemotherapy, country of residence was no longer associated with 1-year survival (EHR = 0.96 (0.85-1.09),  $p = 0.2372$ ) (Table 3, model 13b).

### ***Patients diagnosed with colon cancer who had survived at least one year***

Amongst colon cancer patients longer-term (2-5 years) excess hazards of death were comparable in both countries even after adjusting for age, sex, tumor localization and year of diagnosis (EHR = 1.02 (0.95 – 1.11)) (Table 4, model 3). However, adjustment for stage at diagnosis shows that these patients had a better survival in France than in England (EHR = 0.92 (0.85-0.99),  $p=0.016$ ) (Table 4, model 4). Longer-term (2-5 years) excess hazards of death were comparable in both countries either for localized colon cancer (EHR =1.05 (0.85-1.31)) (Table 4, model 5a) or for advanced colon cancer (EHR =1.03 (0.95 – 1.12)) (Table 4, model 5 b).

### ***Patients diagnosed with rectum cancer who had survived at least one year***

Similarly, longer-term (2-5 years) excess hazards of death for rectal cancer patients were comparable in both countries after adjusting for age, sex and stage at diagnosis (EHR =0.91 (0.81 – 1.03)) (Table 4, Model 11). However, English patients diagnosed with an advanced rectum cancer had a better long-term survival than those living in France (EHR= 1.17 (1.03 – 1.31)) (Table 4, model 11b). After adjustment for

surgical resection with curative intent, longer-term (2-5 years) excess hazards of death were comparable in both countries (EHR= 1.06 (0.93- 1.21)) (Table 4, model 12b).

### ***Complete case analysis results***

As explained in methods section, all survival analyses were repeated in the framework of complete case analysis (Table 3, 4). The results of these analyses were comparable to those performed using multiple imputation and are therefore not repeated here. The only exception concerned the influence of treatment variables (surgical resection with curative intent, chemotherapy and radiotherapy) on the longer-term (2-5 years) excess hazards of death for patients diagnosed with an advanced rectum cancer. In the framework of complete case analyses, patients living in England still had a better survival than French patients (Table 4, model 5b, 6b and 7b).

## ***Discussion***

This study confirms that over the study period five-year colon and rectal cancer survival was higher in France than in England. This difference was primarily due to a larger proportion of English patients dying within the first year after their diagnosis and if an individual survived a year after their diagnosis then the difference in survival between the countries was no longer statistically significant. This study also demonstrates differences in the management of patients between the countries with surgery being used significantly less frequently in England compared to France, these different surgery proportions being related to the survival difference observed.

Potential differences between the datasets collected in each country need to be considered. Indeed, the lack of detail on types of surgery undertaken in the English registry data forced us to limit it to surgical resection with curative intent undertaken within the first six months after the diagnosis. Similarly, the French registries captured information on all chemotherapy and radiotherapy administered irrespective of intent whereas NYCRIS did not. Efforts have been made to address these differences in these analyses by attempting to limit the French treatment information included to that comparable to the English data. But, for future analyses, it would be highly desirable to have more detailed treatment datasets that included information on these factors to enable more robust comparisons.

Likewise, differences may exist in the English and French datasets in relation to other important variables such as stage. NYCRIS aims to collect stage at diagnosis and as neo-adjuvant treatments can downstage rectal tumours those receiving them were classified as “not staged”. In addition, accurate staging requires accurate pathological assessment of a resected tumour specimen. NYCRIS had a lower proportion of major resection, so a lower proportion of resected specimens and, therefore, a lower proportion of patients in which a robust stage could be captured. The recording of stage improved considerably in the NYCRIS data from 1998 onwards (registry merger in 1997) . To test if our findings

were influenced by this change, an analysis restricted to 1998-2004 data was conducted which provided similar findings (results not presented).

Missing data multiple imputation was undertaken in our study for dealing with missing values (Nur *et al*, 2010). Unfortunately, given the reasons for the lower staging proportion in England compared to France, the missing values for stage at diagnosis were probably not missing at random (i.e. patients with missing stage tend to have a more advanced disease than others), which could reduce the robustness of our results, since the effect of the absence of treatment in survival could be under-estimated. But, to our knowledge, no unbiased method exists for taking into account missing values not at random.

Another limitation of this study is the small proportion of the French population covered by the French digestive cancer registries (3% of the whole national population), whilst NYCRIS data represents 13.3% of the national population of England. Such a limitation is usual for population-based studies in France since information on stage at diagnosis and treatment are not routinely available in all French cancer registries but only in specialised cancer registries. The 3 specialised digestive cancer registries in France are located in Calvados, Côte d'Or and Saone et Loire, these areas being mainly rural, two of them having a reference cancer centre (University hospital and cancer care centre) in their regional capital. It is noteworthy that French colorectal cancer one- and five year-relative survival calculated in this study were very closed to those estimated on all French cancer registries, which cover about 17% of the French population (Bossard *et al*, 2007).

In a previous study, the difference in observed survival between Europe and America was mainly explained by the stage at diagnosis (Gatta *et al*, 2000). Our study does not give such importance to stage at diagnosis in explaining the difference in relative survival between England and France. In a more recent study of relative survival which included 10 European cancer registries (from Italy, France, Netherlands, Spain and UK) and 9 US cancer registries, the role of surgical practices (curative resection)

appeared to be as important as stage at diagnosis (Ciccolallo *et al*, 2005). In 1990, a randomised trial reported that chemotherapy improved survival for stage III colon cancer (Moertel *et al*, 1990). European guidelines for the management of colon cancer also recommended the prescription of adjuvant chemotherapy for stage II colon cancer for which the number of lymph sampling is <12, the tumours are poorly differentiated, vascular or lymphatic or perineural invasion has occurred, or presentation is emergency and at pT4 stage (Labianca *et al*, 2010). Chemotherapy is also recommended for metastatic colon and rectum tumours. Studies comparing adjuvant therapy uptake between countries are rare because such detailed data are not usually available at a population level. Nevertheless, a recent study highlights that proportions of adjuvant therapies differed notably between European areas either for stage II or stage III tumours (Gatta *et al*, 2010). For example, 60.6 % of patients diagnosed with stage II colon cancer received adjuvant chemotherapy in Slovakia and only 5.3% of such patients received adjuvant therapy in Estonia. Our study confirms that this variation in chemotherapy uptake across countries could be a potential explanation of difference in 1-year survival of advanced colon cancer.

If differences in data quality can be dismissed then the lower treatment proportions and their relationship to survival must be real. What is not apparent from the results is, however, why the treatment proportions in England are lower. There are two potential explanations that could account for this. Firstly, it may be that there are fundamental differences in the characteristics of the two general populations that make fewer English patients eligible for active treatments. For example, English people, including cancer patients, may have a higher prevalence and/or severity of co-morbid disease that limits how these treatments could be employed.

A second alternative explanation may be, however, that there were real differences in the overall management of the cancer patients between the countries during this time period. Such a difference in treatment could partly be explained by differences in medical practices and/or in patient behaviour.

Since indication for surgery was clearly defined by international guidelines, there is no reason to suppose that the practice of an English surgeon would differ to a French one for a given patient. Instead, health care system organization offers a likely explanation for this difference in surgery indication. Indeed, therapeutic delay (time since diagnosis to first treatment) is notably longer in England (Robertson *et al*, 2004) compared to France (Dejardin *et al*, 2004) even if this topic remains poorly investigated in France. Nevertheless, the influence of therapeutic delay on survival is highly controversial (Ramos *et al*, 2007) and, thus, is likely to explain only a part of this difference. Beside medical practices, patient behaviour towards diseases could differ between France and England. Indeed, time from first symptom to diagnosis, as well as time from diagnosis to staging is influenced by behaviour towards symptoms (Mitchell *et al*, 2008), and therefore could influence the distribution of stage between the countries. Whilst this study suggests that stage at diagnosis has no impact on the inter-country gap in survival, we cannot exclude the possibility that English patients had more co-morbidities than French ones for a given stage at diagnosis. Further studies are needed to investigate this additional potential explanation for the gap in survival.

During the period study, major reforms were initiated with the publication of national cancer plans in England in 2000 (Health, 2000) and in France in 2003 (available online at <http://www.plan-cancer.gouv.fr/>). A recent study highlights some beneficial impact of NHS cancer plan in England. For example, the 3-year relative survival for colon cancer increased from 53.5% during the period 1996-2000 to 57.6% for the period 2004-2006 (Rachet *et al*, 2009). Since survival data are not yet published for France, we cannot compare the effectiveness of the French and English cancer plan.

Every effort to investigate the reasons for such disparities between these two countries, and more generally across Europe, should be encouraged. Collection and standardization of additional variables

such as co-morbidity, anaesthetic risk and treatment pathways may be crucial for a better understanding of underlying mechanisms of inter-country differences in survival of patients with cancer.

### ***Acknowledgments***

The authors thank the French National cancer institute and the “Fondation de France”, which provided financial support for this study. BR and EM are supported by Cancer Research UK (C1336/A5735 and C23434/A9805 respectively)

## References

Abdel-Rahman M, Stockton D, Rachet B, Hakulinen T, Coleman MP (2009) What if cancer survival in Britain were the same as in Europe: how many deaths are avoidable? *Br J Cancer***101 Suppl 2**: S115-24

Beral V, Peto R (2010) UK cancer survival statistics. *BMJ***341**: c4112

Berrino F, De Angelis R, Sant M, Rosso S, Bielska-Lasota M, Lasota MB, Coebergh JW, Santaquilani M (2007) Survival for eight major cancers and all cancers combined for European adults diagnosed in 1995-99: results of the EUROCARE-4 study. *Lancet Oncol***8**(9): 773-83

Bossard N, Velten M, Remontet L, Belot A, Maarouf N, Bouvier AM, Guizard AV, Tretarre B, Launoy G, Colonna M, Danzon A, Molinie F, Troussard X, Bourdon-Raverdy N, Carli PM, Jaffre A, Bessaguet C, Sauleau E, Schvartz C, Arveux P, Maynadie M, Grosclaude P, Esteve J, Faivre J (2007) Survival of cancer patients in France: A population-based study from the Association of the French Cancer Registries (FRANCIM). *European Journal of Cancer***43**(1): 149-160

Brenner H, Bouvier AM, Foschi R, Hackl M, Larsen IK, Lemmens V, Mangone L, Francisci S, Group tEW (2011) Progress in colorectal cancer survival in Europe, from the late 1980s to the early 21st century: The EUROCARE study. *Int J Cancer*

Ciccolallo L, Capocaccia R, Coleman MP, Berrino F, Coebergh JWW, Damhuis RAM, Faivre J, Martinez-Garcia C, Moller H, de Leon MP, Launoy G, Raverdy N, Williams EMI, Gatta G (2005) Survival differences between European and US patients with colorectal cancer: role of stage at diagnosis and surgery. *Gut***54**(2): 268-273

Coleman MP, Quaresma M, Berrino F, Lutz JM, De Angelis R, Capocaccia R, Baili P, Rachet B, Gatta G, Hakulinen T, Micheli A, Sant M, Weir HK, Elwood JM, Tsukuma H, Koifman S, E Silva GA, Francisci S, Santaquilani M, Verdecchia A, Storm HH, Young JL (2008) Cancer survival in five continents: a worldwide population-based study (CONCORD). *Lancet Oncol***9**(8): 730-56

Dejardin O, Herbert C, Velten M, Buemi A, Menegoz F, Maarouf N, Launoy G (2004) Social and geographical factors influencing the delay in treatment for colorectal cancer. *British Journal of Cancer***91**(9): 1751-1752

Dickman PW, Sloggett A, Hills M, Hakulinen T (2004) Regression models for relative survival. *Stat Med***23**(1): 51-64

Dukes C (1932) The classification of cancer of the rectum. *Journal of Pathology & Bacteriology***35**: 9

Engholm G, Kejs AM, Brewster DH, Gaard M, Holmberg L, Hartley R, Iddenden R, Møller H, Sankila R, Thomson CS, Storm HH (2007) Colorectal cancer survival in the Nordic countries and the United Kingdom: excess mortality risk analysis of 5 year relative period survival in the period 1999 to 2000. *Int J Cancer***121**(5): 1115-22

Folkesson J, Engholm G, Ehrnrooth E, Kejs AM, Pålman L, Harling H, Wibe A, Gaard M, Thornorvaldur J, Tryggvadottir L, Brewster DH, Hakulinen T, Storm HH (2009) Rectal cancer survival in the Nordic countries and Scotland. *Int J Cancer***125**(10): 2406-12

Fritz A, Percy C, Jack A, Shanmugaratnam K, Sobin L, Parkin Dm, Whelan S (2000) *International Classification of diseases for Oncology - Third Edition*. Geneva: World Health Organization

Gatta G, Capocaccia R, Sant M, Bell CM, Coebergh JW, Damhuis RA, Faivre J, Martinez-Garcia C, Pawlega J, Ponz de Leon M, Pottier D, Raverdy N, Williams EM, Berrino F (2000) Understanding variations in survival for colorectal cancer in Europe: a EURO CARE high resolution study. *Gut***47**(4): 533-8

Gatta G, Zigon G, Aareleid T, Ardanaz E, Bielska-Lasota M, Galceran J, Gózdź S, Hakulinen T, Martinez-Garcia C, Plesko I, Zakelj MP, Rachtan J, Tagliabue G, Vercelli M, Faivre J (2010) Patterns of care for European colorectal cancer patients diagnosed 1996-1998: a EURO CARE high resolution study. *Acta Oncol***49**(6): 776-83

Health Do (2000) The NHS Cancer plan, Health Do (ed). London: Department of Health

Jones AM, Morris E, Thomas J, Forman D, Melia J, Moss SM (2009) Evaluation of bowel cancer registration data in England, 1996-2004. *Br J Cancer***101**(8): 1269-73

Labianca R, Nordlinger B, Beretta GD, Brouquet A, Cervantes A, Group EGW (2010) Primary colon cancer: ESMO Clinical Practice Guidelines for diagnosis, adjuvant treatment and follow-up. *Ann Oncol***21 Suppl 5**: v70-7

Little R, Rubin D (2002) *Statistical analysis with missing data*. Hoboken, New Jersey: John Wiley & Sons, Inc.

Mitchell E, Macdonald S, Campbell NC, Weller D, Macleod U (2008) Influences on pre-hospital delay in the diagnosis of colorectal cancer: a systematic review. *Br J Cancer***98**(1): 60-70

Moertel CG, Fleming TR, Macdonald JS, Haller DG, Laurie JA, Goodman PJ, Ungerleider JS, Emerson WA, Tormey DC, Glick JH (1990) Levamisole and fluorouracil for adjuvant therapy of resected colon carcinoma. *N Engl J Med***322**(6): 352-8

Morris EJ, Sandin F, Lambert PC, Bray F, Klint A, Linklater K, Robinson D, Pahlman L, Holmberg L, Møller H (2011) A population-based comparison of the survival of patients with colorectal cancer in England, Norway and Sweden between 1996 and 2004. *Gut***60**(8): 1087-93

Møller H, Richards S, Hanchett N, Riaz SP, Lichtenborg M, Holmberg L, Robinson D (2011) Completeness of case ascertainment and survival time error in English cancer registries: impact on 1-year survival estimates. *Br J Cancer***105**(1): 170-6

Nur U, Shack LG, Rachet B, Carpenter JR, Coleman MP (2010) Modelling relative survival in the presence of incomplete data: a tutorial. *Int J Epidemiol***39**(1): 118-28

Rachet B, Maringe C, Nur U, Quaresma M, Shah A, Woods LM, Ellis L, Walters S, Forman D, Steward J, Coleman MP (2009) Population-based cancer survival trends in England and Wales up to 2007: an assessment of the NHS cancer plan for England. *Lancet Onco***10**(4): 351-69

Ramos M, Esteva M, Cabeza E, Campillo C, Llobera J, Aguiló A (2007) Relationship of diagnostic and therapeutic delay with survival in colorectal cancer: a review. *Eur J Cancer***43**(17): 2467-78

Robertson R, Campbell NC, Smith S, Donnan PT, Sullivan F, Duffy R, Ritchie LD, Millar D, Cassidy J, Munro A (2004) Factors influencing time from presentation to treatment of colorectal and breast cancer in urban and rural areas. *British Journal of Cancer***90**(8): 1479-1485

Royston P (2009) Multiple imputation of missing values: Further update of *ice*, with an emphasis on categorical variables. *STATA Journal***9**(2): 11

Rubin D (1987) *Multiple Imputation for Nonresponse in Surveys*. New-York: John Wiley & Sons

Woods LM, Coleman MP, Lawrence G, Rashbass J, Berrino F, Rachet B (2011) Evidence against the proposition that "UK cancer survival statistics are misleading": simulation study with National Cancer Registry data. *BMJ***342**: d3399

**Table 1 : Patients diagnosed with colorectal cancer in two European countries between 1997-2004 (N=40 613)**

Variables	Calvados		Côte d'Or		Saone et Loire		England (Nycris)		P-values
	N= 2481		N= 2329		N= 3081		N= 32722		
	N	%	N	%	N	%	N	%	
<b>Sex</b>									0.402
Men	1359	54.8%	1300	55.8%	1683	54.6%	18176	55.5%	
Women	1122	45.2%	1029	44.2%	1398	45.4%	14546	44.5%	
<b>Age</b>									0.001
<61	523	21.1%	482	20.7%	508	16.5%	5865	17.9%	
61-69	475	19.1%	452	19.4%	672	21.8%	7200	22.0%	
70-79	712	28.7%	641	27.5%	893	29.0%	9418	28.8%	
>=79	771	31.1%	754	32.4%	1008	32.7%	10239	31.3%	
<b>Topography</b>									<0.001
Right colon	765	30.8%	740	31.8%	1024	33.2%	8881	27.1%	
Left Colon	715	28.8%	717	30.8%	923	30.0%	7917	24.2%	
Unknown Colon *	65	2.6%	18	0.8%	7	0.2%	2879	8.8%	
Recto-sigmoidejunction	235	9.5%	355	15.2%	326	10.6%	3269	10.0%	
Rectum	701	28.3%	499	21.4%	801	26.0%	9776	29.9%	
<b>Duke's stage</b>									<0.001
A	390	15.7%	409	17.6%	567	18.4%	3724	11.4%	
B	676	27.2%	713	30.6%	872	28.3%	8567	26.2%	
C	592	23.9%	477	20.5%	657	21.3%	7686	23.5%	
"D"	644	26.0%	543	23.3%	743	24.1%	7234	22.1%	
Unstaged	179	7.2%	187	8.0%	242	7.9%	5511	16.8%	
<b>Year of diagnosis</b>									0.007
1997	287	11.6%	267	11.5%	393	12.8%	3782	11.6%	
1998	305	12.3%	258	11.1%	389	12.6%	4079	12.5%	
1999	300	12.1%	287	12.3%	351	11.4%	4143	12.7%	
2000	308	12.4%	280	12.0%	418	13.6%	4185	12.8%	
2001	304	12.3%	267	11.5%	374	12.1%	4069	12.4%	
2002	299	12.1%	296	12.7%	356	11.6%	4191	12.8%	
2003	345	13.9%	324	13.9%	416	13.5%	4096	12.5%	
2004	333	13.4%	350	15.0%	384	12.5%	4177	12.8%	
<b>Surgical resection within 6 months since diagnosis</b>									<0,001
Yes	2065	83.2%	1944	83.5%	2592	84.1%	23908	73.1%	
No	416	16.8%	385	16.5%	489	15.9%	8814	26.9%	
<b>Chemotherapy</b>									<0.001
Yes	905	36.5%	751	32.2%	1069	34.7%	8542	26.1%	
No	1559	62.8%	1564	67.2%	1982	64.3%	24180	73.9%	
unknown	17	0.7%	14	0.6%	30	1.0%	0	0.0%	
<b>Radiotherapy</b>									<0.001
Yes	375	15.1%	374	16.1%	586	19.0%	4027	12.3%	
No	2095	84.4%	1943	83.4%	2467	80.1%	28695	87.7%	
unknown	11	0.4%	12	0.5%	28	0.9%	0	0.0%	

\* Including C18,8

**Table 2: Distribution of Duke's stage before and after multiple imputation by site and country of residence**

	<b>Complete Case analysis</b>				<b>Multiple Imputation</b>			
	France		England		France		England	
	Colon	Rectum	Colon	Rectum	Colon	Rectum	Colon	Rectum
Duke's A	14.9%	31.7%	10.4%	22.1%	14.3%	27.2%	9.7%	19.2%
Duke's B	33.3%	23.4%	34.0%	25.1%	31.9%	21.0%	31.0%	23.7%
Duke's C	24.5%	20.9%	27.7%	29.6%	23.7%	20.2%	26.6%	28.7%
"Duke's D"	27.2%	24.0%	27.9%	23.2%	30.0%	31.5%	32.7%	28.4%

**Table 3 : Excess hazard ratios of death for country of residence during the first year since diagnosis, colorectal cancer, 1997-2004**

N°	Variable	Multiple imputations for missing values				Complete case analysis			
		EHR	95%CI		P-values	EHR	95%CI		P-values
<b>Colon Cancer</b>									
<b>Model 1</b>	countries	England	1.00			<0,001	1.00		<0,001
		France	0.70	0.66	0.75		0.73	0.68	0.78
<b>Model 2</b>	=Model 1 + Age, sex, period								
		England	1.00			<0,001	1.00		<0,001
		France	0.70	0.65	0.74		0.73	0.68	0.78
<b>Model 3</b>	=Model 2 +Cancer localisation*								
		England	1.00			<0,001	1.00		<0,001
		France	0.70	0.65	0.74		0.81	0.75	0.86
<b>Model 4</b>	=Model 3 +Stage								
		England	1.00			<0,001	1.00		<0,001
		France	0.68	0.64	0.74		0.75	0.70	0.80
<b>Model 5 a</b>	=Model 2 for Duke's A & B only								
		England	1.00			0.006	1.00		0.033
		France	0.77	0.65	0.93		0.79	0.66	0.95
<b>Model 6 a</b>	=Model 5a for Duke's A & B + resection within 6 months								
		England	1.00			0.038	1.00		0.038
		France	0.82	0.68	0.99		0.82	0.68	0.98
<b>Model 5 b</b>	=model 3 for Duke's C & D cancer only								
		England	1.00			<0,001	1.00		<0,001
		France	0.67	0.63	0.72		0.74	0.69	0.80
<b>Model 6 b</b>	=model 5b for Duke's C & D cancer only + resection within 6 months								
		England	1.00			<0,001	1.00		<0,001
		France	0.79	0.74	0.85		0.83	0.77	0.90
<b>Model 7 b</b>	=model 5b for Duke's C & D cancer only + resection within 6 months+ chemotherapy								
		England	1.00			0.095	1.00		0.862
		France	0.94	0.88	1.01		1.00	0.92	1.06
<b>Model 8 b</b>	=model 5b for Duke's C & D cancer only + resection within 6 months+ chemotherapy+ interaction (countries-chemotherapy)								
		England	1.00			0.918	1.00		0.082
Withoutchemotherapy		France	0.99	0.92	1.07		1.07	0.99	1.17
		England	1.00			0.008	1.00		0.001
Withchemotherapy		France	0.81	0.69	0.94		0.76	0.65	0.89
<b>Rectum cancer</b>									
<b>Model 9</b>		England	1.00			<0,001	1.00		<0,001
		France	0.76	0.68	0.86		0.74	0.64	0.85
<b>Model 10</b>	=Model 9 + Age, sex, period								
		England	1.00			<0,001	1.00		<0,001
		France	0.76	0.67	0.85		0.73	0.63	0.84
<b>Model 11</b>	=Model 10 + Stage								
		England	1.00			<0,001	1.00		<0,001
		France	0.69	0.61	0.77		0.67	0.58	0.77
<b>Model 11a</b>	=Model 10 for duke's A & B cancer only								
		England	1.00			0.048	1.00		0.065
		France	0.70	0.49	0.99		0.71	0.50	1.02
<b>Model 12a</b>	=Model 10 for Duke's A & B cancer only + resection within 6 months								
		England	1.00			0.13	1.00		0.353
		France	0.74	0.51	1.08		0.83	0.57	1.22
<b>Model 11b</b>	=Model 10 for Duke's C & D cancer only								
		England	1.00			<0,001	1.00		<0,001
		France	0.69	0.61	0.77		0.66	0.57	0.77
<b>Model 12b</b>	=Model 11a for Duke's C & D cancer only + resection within 6 months								
		England	1.00			<0,001	1.00		<0,001
		France	0.73	0.64	0.82		0.77	0.66	0.89
<b>Model 13b</b>	=Model 11a for Duke's C & D cancer only + resection within 6 months + Radiotherapy + Chemotherapy								
		England	1.00			0.575	1.00		0.637
		France	0.96	0.85	1.09		1.01	0.86	1.10

\* Unknow Colon Cancer (C18.8 or C18.9) were kept in complete case analysis

**Table 4 : Excess hazard ratios of death for country of residence after the first year since diagnosis (2-5 years), colorectal cancer, 1997-2004**

N°	Variable	<i>Multiple imputations for missing values</i>			<i>Complete case analysis</i>				
		EHR	95%CI	P-values	EHR	95%CI	P-values		
<b>Colon Cancer</b>									
<b>Model 1</b>	countries	England	1.00			0.571	1.00		0.407
		France	1.02	0.95	1.11		1.03	0.95	1.11
<b>Model 2</b>	=Model 1 + Age, sex, period								
		England	1.00			0.518	1.00		0.458
		France	1.03	0.95	1.11		1.03	0.95	1.12
<b>Model 3</b>	=Model 2 +Cancer localisation								
		England	1.00			0.472	1.00		0.414
		France	1.03	0.95	1.11		1.03	0.95	1.12
<b>Model 4</b>	=Model 3 +Stage								
		England	1.00			0.032	1.00		0.036
		France	0.92	0.85	0.99		0.92	0.85	0.99
<b>Model 5 a</b>	=Model 3 for duke's A & B cancer only								
		England	1.00			0.623	1.00		0.829
		France	1.05	0.85	1.31		1.02	0.82	1.28
<b>Model 6 a</b>	=Model 5a for Duke's A & B cancer only + resection within 6 months								
		England	1.00			0.318	1.00		0.737
		France	1.07	0.86	1.33		1.04	0.84	1.30
<b>Model 5 b</b>	=model 3 for Duke's C & D cancer only								
		England	1.00			0.449	1.00		0.199
		France	1.03	0.95	1.12		1.06	0.97	1.15
<b>Model 6 b</b>	=model 5b for Duke's C & D cancer only + resection surgery within 6 months								
		England	1.00			0.25	1.00		0.449
		France	1.04	0.97	1.14		1.03	0.95	1.12
<b>Rectum cancer</b>									
<b>Model 9</b>		England	1.00			0.733	1.00		0.891
		France	0.98	0.87	1.10		1.01	0.90	1.14
<b>Model 10</b>	=Model 9 + Age, sex, period								
		England	1.00			0.807	1.00		0.829
		France	0.99	0.88	1.11		1.01	0.89	1.15
<b>Model 11</b>	=Model 10 + Stage								
		England	1.00			0.127	1.00		0.542
		France	0.91	0.81	1.03		1.04	0.92	1.18
<b>Model 11a</b>	=Model 9 for Duke's A & B cancer only								
		England	1.00			0.161	1.00		0.734
		France	0.81	0.60	1.09		0.89	0.45	1.76
<b>Model 11b</b>	=Model 11a for Duke's C & D cancer only								
		England	1.00			0.011	1.00		<0.001
		France	1.17	1.03	1.31		1.36	1.19	1.56
<b>Model 12b</b>	=Model 11a for Duke's C & D cancer only + resection within 6 months								
		England	1.00			0.331	1.00		<0.001
		France	1.06	0.93	1.21		1.39	1.22	1.60
<b>Model 13b</b>	=Model 12a for Duke's C & D cancer only + radiotherapy + chemotherapy								
		England	1.00			0.218	1.00		<0.001
		France	1.09	0.95	1.24		1.45	1.26	1.66

\* Unknow Colon Cancer (C18.8 or C18.9) were kept in complete case analysis