

# Associations between health and sexual lifestyles in Britain: findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3)



Nigel Field, Catherine H Mercer, Pam Sonnenberg, Clare Tanton, Soazig Clifton, Kirstin R Mitchell, Bob Erens, Wendy Macdowall, Frederick Wu, Jessica Datta, Kyle G Jones, Amy Stevens, Philip Prah, Andrew J Copas, Andrew Phelps, Kaye Wellings\*, Anne M Johnson\*



## Summary

**Background** Physical and mental health could greatly affect sexual activity and fulfilment, but the nature of associations at a population level is poorly understood. We used data from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3) to explore associations between health and sexual lifestyles in Britain (England, Scotland, and Wales).

**Methods** Men and women aged 16–74 years who were resident in households in Britain were interviewed between Sept 6, 2010, and Aug 31, 2012. Participants completed the survey in their own homes through computer-assisted face-to-face interviews and self-interview. We analysed data for self-reported health status, chronic conditions, and sexual lifestyles, weighted to account for unequal selection probabilities and non-response to correct for differences in sex, age group, and region according to 2011 Census figures.

**Findings** Interviews were done with 15162 participants (6293 men, 8869 women). The proportion reporting recent sexual activity (one or more occasion of vaginal, oral, or anal sex with a partner of the opposite sex, or oral or anal sex or genital contact with a partner of the same sex in the past 4 weeks) decreased with age after the age of 45 years in men and after the age of 35 years in women, while the proportion in poorer health categories increased with age. Recent sexual activity was less common in participants reporting bad or very bad health than in those reporting very good health (men: 35.7% [95% CI 28.6–43.5] vs 74.8% [72.7–76.7]; women: 34.0% [28.6–39.9] vs 67.4% [65.4–69.3]), and this association remained after adjusting for age and relationship status (men: adjusted odds ratio [AOR] 0.29 [95% CI 0.19–0.44]; women: 0.43 [0.31–0.61]). Sexual satisfaction generally decreased with age, and was significantly lower in those reporting bad or very bad health than in those reporting very good health (men: 45.4% [38.4–52.7] vs 69.5% [67.3–71.6], AOR 0.51 [0.36–0.72]; women: 48.6% [42.9–54.3] vs 65.6% [63.6–67.4], AOR 0.69 [0.53–0.91]). In both sexes, reduced sexual activity and reduced satisfaction were associated with limiting disability and depressive symptoms, and reduced sexual activity was associated with chronic airways disease and difficulty walking up the stairs because of a health problem. 16.6% (95% CI 15.4–17.7) of men and 17.2% (16.3–18.2) of women reported that their health had affected their sex life in the past year, increasing to about 60% in those reporting bad or very bad health. 23.5% (20.3–26.9) of men and 18.4% (16.0–20.9) of women who reported that their health affected their sex life reported that they had sought clinical help (>80% from general practitioners; <10% from specialist services).

**Interpretation** Poor health is independently associated with decreased sexual activity and satisfaction at all ages in Britain. Many people in poor health report an effect on their sex life, but few seek clinical help. Sexual lifestyle advice should be a component of holistic health care for patients with chronic ill health.

**Funding** Grants from the UK Medical Research Council and the Wellcome Trust, with support from the Economic and Social Research Council and Department of Health.

## Introduction

Physical and mental health disorders, the drugs used to treat them, and long-term disability could greatly affect sexual lifestyles<sup>1–4</sup>—ie, sexual activity, sexual behaviours, sexual problems, the formation and maintenance of relationships, and sexual satisfaction.<sup>5–7</sup> For example, poor self-assessed general health is associated with reduced sexual activity and frequency of sexual activity in older people.<sup>4</sup> However, most population-based studies investigating these associations have not been specifically designed to measure sexual behaviour, and have included only one or a few measures of health,<sup>8–12</sup> surveyed only men,<sup>13–15</sup> or been focused on older people.<sup>4,8,13,14,16</sup> In the

past 5 years, some studies have included both self-reported and biological or physical measures of a few specific disorders to explore associations with sexual function.<sup>17,18</sup> However, some chronic conditions—eg, arthritis, stroke, or heart disease—cannot easily be measured with biological or physical measures in large surveys of this kind. Therefore, the associations between ill health or disabilities and sexual lifestyles are not well described across the sexually active adult life at a population level. Moreover, the effect of health on people's sex lives is seldom considered in clinical practice,<sup>19,20</sup> and the evidence base that can be used to guide clinical management is small.

Published Online  
November 26, 2013  
[http://dx.doi.org/10.1016/S0140-6736\(13\)62222-9](http://dx.doi.org/10.1016/S0140-6736(13)62222-9)

See Online/Comment  
[http://dx.doi.org/10.1016/S0140-6736\(13\)62378-8](http://dx.doi.org/10.1016/S0140-6736(13)62378-8)

Copyright © Field et al. Open Access article distributed under the terms of CC BY

\*Joint senior authors

See Online for video infographic

Research Department of Infection and Population Health, University College London, London, UK

(N Field MBPhD, CH Mercer PhD, P Sonnenberg PhD,

C Tanton PhD, S Clifton BSc,

B Erens MA, K G Jones MSc,

A Stevens BSc, P Prah MSc,

A J Copas PhD,

Prof A M Johnson MD); NatCen

Social Research, London, UK

(S Clifton, A Phelps BA);

Department of Social and

Environmental Health Research

(K R Mitchell PhD,

W Macdowall MSc, J Datta MSc,

Prof K Wellings FRCOG) and

Health Services Research and

Policy (B Erens), London School

of Hygiene and Tropical

Medicine, London, UK; and

Centre for Endocrinology and

Diabetes, Institute of Human

Development, University of

Manchester, Manchester, UK

(Prof F Wu FRCP)

Correspondence to:

Dr Nigel Field, Research

Department of Infection and

Population Health, University

College London, Mortimer

Market Centre, London

WC1E 6JB, UK

[nigel.field@ucl.ac.uk](mailto:nigel.field@ucl.ac.uk)

A decrease in sexual activity and frequency of sexual activity is associated with increasing age, although about half of people aged 65–74 years report sexual activity in the past year.<sup>4,5,8,9,13,21</sup> As the strongest driver of chronic ill health and long-term disability, age is therefore an important confounder in studies of the associations between health and sexual lifestyles. Likewise, the availability of partners—which could be limited by widowhood, no stable partner, or other reasons—affects these associations.<sup>1</sup>

In the past two decades, detailed information about sexual lifestyles across the population in Britain (England, Scotland, and Wales) has been obtained in the National Surveys of Sexual Attitudes and Lifestyles (Natsal). However, these large probability sample surveys have not previously included detailed questions about health.<sup>5–7</sup> In the third Natsal survey (Natsal-3), participants were asked to provide information about their health, including self-assessed general health, disability, functional impairment, and specific chronic conditions. Furthermore, people aged 16–74 years were included, which provides a rare opportunity to investigate the associations between sexual lifestyles and health across the life course in men and women.

We tested the hypothesis that poor health, disability, and specific chronic conditions would remain associated with three measures of sexual lifestyles after adjustment for age

and relationship status: recent sexual activity, sexual satisfaction, and sexual response problems specific to men or women (erectile difficulties and vaginal dryness). To provide actionable information for clinicians and policy makers, we investigated whether people felt that any health condition or disability had affected their sex lives and the extent to which clinical advice had been sought.

**Methods**

**Participants and procedures**

Full details of the methods of Natsal-3 have been reported elsewhere.<sup>5,22,23</sup> Briefly, we used a multistage, clustered, and stratified probability sample design. Men and women aged 16–74 years who were resident in households in Britain were interviewed between Sept 6, 2010, and Aug 31, 2012. An anonymised dataset will be deposited with the UK Data Archive, and the complete questionnaire and technical report will be available on the Natsal website on the day of publication.

Participants were interviewed in their own homes by professional interviewers without clinical qualifications using computer-assisted personal interview. Participants were asked about self-assessed general health with a widely cited and validated question,<sup>24</sup> and about any longstanding and restricting illness or disability. To measure self-reported functional ability, participants were

For the questionnaire and more information about Natsal-3 see <http://www.natsal.ac.uk>

	16–24 years	25–34 years	35–44 years	45–54 years	55–64 years	65–74 years	All age groups
<b>Men</b>							
Self-reported general health status							
Very good	49.4% (46.6–52.2)	47.8% (45.0–50.6)	41.6% (37.9–45.4)	35.8% (40.6–48.8)	26.6% (38.2–46.3)	24.6% (37.7–46.2)	38.8% (37.4–40.3)
Good	42.4% (39.6–45.1)	41.0% (38.2–43.9)	45.1% (41.3–48.9)	44.6% (40.6–48.8)	42.2% (38.2–46.3)	41.9% (37.7–46.2)	43.0% (41.4–44.6)
Fair	7.5% (6.2–9.0)	9.9% (8.2–11.9)	10.9% (9.0–13.3)	15.2% (12.6–18.2)	21.7% (18.5–25.4)	27.1% (23.5–31.0)	14.4% (13.3–15.5)
Bad or very bad	0.7% (0.4–1.2)	1.2% (0.8–1.9)	2.4% (1.6–3.6)	4.4% (3.2–6.0)	9.5% (7.4–12.1)	6.4% (4.6–8.9)	3.8% (3.3–4.4)
Unweighted denominator	1689	1473	783	747	697	603	5992
Weighted denominator	1210	1332	1386	1334	1093	779	7133
Longstanding illnesses or disability							
None	86.2% (84.1–88.1)	82.6% (80.4–84.6)	76.2% (72.8–79.3)	63.8% (59.9–67.6)	46.6% (42.7–50.5)	43.2% (39.1–47.3)	68.6% (67.2–70.0)
Non-limiting	8.3% (7.0–10.0)	8.4% (7.0–10.1)	10.6% (8.4–13.3)	18.2% (15.2–21.7)	24.4% (21.1–28.1)	27.3% (23.8–31.2)	15.2% (14.1–16.3)
Limiting	5.4% (4.3–6.8)	9.0% (7.5–10.8)	13.2% (11.0–15.8)	17.9% (15.2–21.0)	29.0% (25.5–32.7)	29.5% (25.9–33.4)	16.2% (15.1–17.3)
Unweighted denominator	1689	1472	783	748	697	603	5992
Weighted denominator	1210	1332	1386	1335	1094	779	7135
Number of self-reported chronic conditions*							
0	90.6% (89.0–92.1)	82.4% (80.0–84.6)	71.8% (68.1–75.2)	58.0% (54.1–61.8)	38.5% (34.7–42.4)	30.0% (26.1–34.1)	64.7% (63.2–66.2)
1	8.5% (7.1–10.1)	14.5% (12.6–16.7)	21.4% (18.4–24.8)	28.3% (24.9–32.0)	33.0% (29.5–36.7)	33.8% (29.8–38.0)	22.3% (21.0–23.7)
≥2	0.9% (0.6–1.4)	3.1% (2.1–4.6)	6.8% (5.3–8.7)	13.7% (11.2–16.6)	28.5% (25.2–32.1)	36.3% (32.3–40.5)	13.0% (11.9–14.0)
Unweighted denominator	1687	1471	783	748	697	602	5988
Weighted denominator	1208	1331	1386	1335	1094	778	7132
Difficulty walking up stairs because of a health problem							
No difficulty	98.2% (97.4–98.8)	96.5% (95.0–97.5)	94.4% (92.5–95.8)	88.2% (85.5–90.5)	79.6% (76.3–82.7)	72.4% (68.3–76.2)	89.6% (88.6–90.5)
Some difficulty	1.7% (1.1–2.5)	3.1% (2.1–4.6)	3.8% (2.6–5.5)	9.7% (7.7–12.2)	13.1% (10.6–16.0)	19.5% (16.3–23.1)	7.6% (6.8–8.4)
Much difficulty or unable	0.1% (0.0–0.4)	0.4% (0.2–0.8)	1.8% (1.1–3.0)	2.1% (1.3–3.3)	7.3% (5.4–9.7)	8.1% (5.9–10.9)	2.8% (2.4–3.4)
Unweighted denominator	1689	1473	782	748	698	603	5993
Weighted denominator	1210	1332	1384	1335	1095	779	7135

(Continues on next page)

	16–24 years	25–34 years	35–44 years	45–54 years	55–64 years	65–74 years	All age groups
(Continued from previous page)							
<b>Women</b>							
Self-reported general health status							
Very good	45.7% (43.4–48.1)	46.1% (43.8–48.3)	47.7% (44.6–50.9)	36.5% (39.1–45.7)	32.6% (36.5–43.0)	26.0% (37.6–44.8)	40.0% (38.8–41.2)
Good	43.5% (41.1–45.8)	42.8% (40.6–45.1)	39.4% (36.4–42.6)	42.4% (39.1–45.7)	39.7% (36.5–43.0)	41.2% (37.6–44.8)	41.5% (40.3–42.7)
Fair	8.9% (7.7–10.2)	9.5% (8.2–10.9)	10.0% (8.3–12.0)	15.5% (13.2–18.1)	19.6% (17.2–22.3)	23.6% (20.7–26.7)	13.9% (13.0–14.8)
Bad or very bad	1.9% (1.4–2.7)	1.7% (1.2–2.3)	2.8% (2.0–4.0)	5.7% (4.4–7.2)	8.1% (6.4–10.2)	9.3% (7.3–11.8)	4.6% (4.1–5.2)
Unweighted denominator	2081	2379	1158	1057	974	803	8452
Weighted denominator	1172	1325	1393	1366	1166	856	7278
Longstanding illnesses or disability							
None	82.6% (80.7–84.4)	78.7% (77.0–80.4)	73.6% (70.9–76.1)	63.7% (60.4–66.8)	51.4% (47.8–55.0)	41.4% (37.6–45.2)	66.8% (65.5–68.0)
Non-limiting	8.5% (7.1–10.0)	11.0% (9.8–12.3)	12.8% (10.9–15.0)	12.9% (10.8–15.4)	20.2% (17.5–23.2)	26.2% (22.9–29.7)	14.5% (13.6–15.5)
Limiting	8.9% (7.7–10.3)	10.3% (9.0–11.7)	13.7% (11.7–15.9)	23.4% (20.8–26.3)	28.4% (25.3–31.7)	32.5% (29.0–36.2)	18.7% (17.7–19.7)
Unweighted denominator	2081	2378	1158	1057	974	803	8451
Weighted denominator	1172	1324	1393	1366	1166	856	7278
Number of self-reported chronic conditions*							
0	81.9% (80.0–83.6)	72.9% (70.9–74.9)	66.6% (63.7–69.4)	51.2% (47.8–54.7)	36.6% (33.3–40.1)	23.5% (20.5–26.8)	57.5% (56.2–58.7)
1	15.8% (14.2–17.5)	21.3% (19.5–23.2)	23.0% (20.5–25.7)	28.6% (25.6–31.8)	31.9% (28.6–35.3)	30.9% (27.5–34.5)	24.9% (23.8–26.0)
≥2	2.4% (1.8–3.2)	5.7% (4.8–6.8)	10.4% (8.7–12.3)	20.2% (17.7–22.9)	31.5% (28.4–34.8)	45.6% (41.9–49.4)	17.6% (16.6–18.6)
Unweighted denominator	2080	2378	1156	1055	974	803	8446
Weighted denominator	1171	1324	1391	1364	1166	856	7273
Difficulty walking up stairs because of health problem							
No difficulty	96.4% (95.5–97.2)	95.7% (94.7–96.6)	92.0% (90.2–93.6)	82.5% (79.8–84.8)	74.5% (71.4–77.5)	61.4% (57.7–65.1)	85.2% (84.2–86.2)
Some difficulty	3.2% (2.5–4.1)	3.4% (2.6–4.3)	6.1% (4.7–7.7)	12.2% (10.2–14.6)	17.5% (15.1–20.2)	25.9% (22.8–29.2)	10.4% (9.6–11.3)
Much difficulty or unable	0.4% (0.2–0.7)	0.9% (0.6–1.4)	1.9% (1.2–2.9)	5.3% (4.1–6.8)	7.9% (6.2–10.0)	12.7% (10.4–15.5)	4.3% (3.8–4.9)
Unweighted denominator	2081	2379	1158	1057	973	803	8451
Weighted denominator	1172	1325	1393	1366	1165	856	7276

Data in parentheses are 95% CIs. Denominators are all individuals reporting one or more sexual partner over the lifetime and vary across variables because of item non-response. For all variables and both sexes, the  $\chi^2$  p value for association with age was <0.0001. \*Measure of comorbidity includes arthritis, heart attack, coronary heart disease, angina, other forms of heart disease, hypertension, stroke, diabetes, broken hip or pelvis bone or hip replacement ever, backache lasting longer than 3 months, any other muscle or bone disease lasting longer than 3 months, depression, cancer, and any thyroid condition treated in the past year.

**Table 1: General health characteristics of participants reporting at least one sexual partner over the lifetime, by age group and sex**

asked whether they had any difficulty walking up a flight of stairs because of a health problem.<sup>25</sup> Body-mass index (BMI) was calculated from self-reported height and weight. To obtain information about self-reported clinical diagnoses of a range of chronic conditions, interviewers used show cards listing different disorders to ask participants whether they had any of the conditions listed. The first card asked whether a doctor had ever told them that they had cardiac or vascular diseases, hypertension, diabetes, chronic airways disease, and arthritis. Separate cards asked whether participants had ever had prostate diseases, broken hip or pelvis, and hip replacement, and whether they had received treatment from a health-care professional for any backache or bone or muscle disease lasting for more than 3 months in the past year.

Participants subsequently completed a computer-assisted self-interview, which included a validated two-question patient health questionnaire (PHQ-2), with which depressive symptoms in the past 2 weeks were assessed with two screening questions (scored 0–3).<sup>26</sup> Participants were deemed to have depressive symptoms if they had a total score of 3 or more, a cutoff which has been previously validated.<sup>27</sup> Participants were asked

whether they had had any health condition or disability in the past year that they felt had affected their sexual activity or enjoyment in any way. Additionally, participants were asked whether they had sought help or advice about their sex life from a range of sources in the past year; more than one answer was allowed. The questionnaire underwent thorough cognitive testing and piloting, as previously reported.<sup>23,28</sup>

The computer-assisted self-interview included many questions about sexual practices, partners, and activity.<sup>22</sup> Here, we focus on four measures: recent sexual activity, satisfaction with sex life, sexual response problems, and relationship status. Recent sexual activity was defined as reporting of one or more occasion of vaginal, oral, or anal sex with a partner of the opposite sex, or oral or anal sex or genital contact with a partner of the same sex in the past 4 weeks. Participants were asked to think about their sex life in the past year in response to the statement “I feel satisfied with my sex life”. We deemed that those answering that they agreed or agreed strongly were satisfied with their sex life. Participants reporting at least one sexual partner in the past year were asked to report which, if any, of a range of sexual

	Men					Women				
	Percentage reporting sexual activity	AOR	p value	Unweighted denominator	Weighted denominator	Percentage reporting sexual activity	AOR	p value	Unweighted denominator	Weighted denominator
All	67.4% (66.0-68.8)	..	..	5994	7137	61.6% (60.4-62.9)	..	..	8452	7278
<b>Demographic characteristics</b>										
Age group			<0.0001					<0.0001		
16-24 years	58.2% (55.5-60.8)	1.00	..	1689	1210	63.4% (60.9-65.7)	1.00	..	2081	1172
25-34 years	79.4% (77.1-81.5)	1.05 (0.82-1.35)	..	1473	1332	80.0% (78.1-81.7)	0.82 (0.65-1.03)	..	2379	1325
35-44 years	79.4% (76.3-82.2)	0.62 (0.46-0.82)	..	783	1386	75.8% (73.1-78.4)	0.45 (0.34-0.58)	..	1158	1393
45-54 years	74.9% (71.4-78.2)	0.53 (0.39-0.72)	..	748	1335	68.0% (64.9-70.9)	0.30 (0.23-0.39)	..	1057	1366
55-64 years	58.8% (54.7-62.7)	0.24 (0.18-0.32)	..	698	1095	43.2% (39.7-46.6)	0.10 (0.08-0.13)	..	974	1166
65-74 years	39.3% (35.2-43.6)	0.09 (0.07-0.13)	..	603	779	22.7% (19.6-26.1)	0.04 (0.03-0.06)	..	803	856
Relationship status			<0.0001					<0.0001		
Living with a partner	78.2% (76.5-79.9)	1.00	..	2938	4649	73.5% (71.9-75.1)	1.00	..	4329	4646
In a steady relationship, not cohabiting	90.7% (88.2-92.7)	1.35 (0.98-1.84)	..	955	766	88.4% (86.3-90.3)	1.18 (0.93-1.49)	..	1355	784
No steady relationship, previously cohabited	27.7% (24.2-31.4)	0.09 (0.07-0.12)	..	752	669	17.9% (15.9-20.1)	0.07 (0.05-0.08)	..	1560	1118
No steady relationship, never cohabited	27.8% (25.1-30.7)	0.02 (0.02-0.03)	..	1297	1003	23.6% (20.8-26.6)	0.02 (0.01-0.02)	..	1162	697
<b>General health</b>										
Self-reported general health status			<0.0001					<0.0001		
Very good	74.8% (72.7-76.7)	1.00	..	2411	2771	67.4% (65.4-69.3)	1.00	..	3429	2912
Good	68.0% (65.9-69.9)	0.73 (0.62-0.87)	..	2532	3067	62.7% (60.8-64.6)	0.92 (0.79-1.07)	..	3545	3020
Fair	54.5% (50.4-58.5)	0.51 (0.40-0.66)	..	826	1025	50.9% (47.6-54.3)	0.81 (0.65-1.01)	..	1125	1010
Bad or very bad	35.7% (28.6-43.5)	0.29 (0.19-0.44)	..	223	270	34.0% (28.6-39.9)	0.43 (0.31-0.61)	..	353	335
Longstanding illnesses or disability			<0.0001					<0.0001		
None	71.7% (70.1-73.2)	1.00	..	4269	4897	67.1% (65.7-68.6)	1.00	..	5856	4859
Non-limiting	64.1% (60.3-67.7)	0.92 (0.74-1.14)	..	803	1084	55.3% (51.8-58.7)	0.86 (0.71-1.04)	..	1131	1058
Limiting	52.6% (48.9-56.2)	0.62 (0.51-0.76)	..	920	1155	46.8% (43.8-49.9)	0.43 (0.38-0.50)	..	1464	1360
Number of self-reported chronic conditions*			<0.0001					<0.0001		
0	71.7% (70.1-73.2)	1.00	..	4140	4615	68.5% (66.9-70.0)	1.00	..	5196	4179
1	66.0% (62.8-69.0)	0.91 (0.74-1.11)	..	1178	1593	60.3% (57.7-62.9)	0.96 (0.81-1.13)	..	1993	1813
≥2	48.9% (44.6-53.2)	0.58 (0.46-0.74)	..	670	924	41.1% (38.0-44.3)	0.67 (0.55-0.81)	..	1257	1282
Body-mass index			0.0134					0.0011		
Normal: 18.5-25 kg/m <sup>2</sup>	67.3% (65.1-69.4)	1.00	..	2633	2820	67.0% (65.2-68.7)	1.00	..	3326	4020
Underweight: <18.5 kg/m <sup>2</sup>	46.0% (35.9-56.5)	0.51 (0.28-0.92)	..	128	105	59.3% (53.2-65.2)	0.80 (0.53-1.22)	..	315	211
Overweight: 25-30 kg/m <sup>2</sup>	70.8% (68.5-73.0)	1.02 (0.85-1.23)	..	2009	2660	59.9% (57.5-62.3)	0.87 (0.74-1.03)	..	2062	1918
Obese: 30-35 kg/m <sup>2</sup>	67.5% (63.6-71.2)	0.87 (0.69-1.10)	..	711	965	55.6% (51.9-59.3)	0.78 (0.63-0.96)	..	937	879
Obese: >35 kg/m <sup>2</sup>	56.3% (48.8-63.6)	0.60 (0.40-0.90)	..	263	353	48.4% (43.5-53.3)	0.57 (0.43-0.76)	..	535	484
Difficulty walking up stairs because of health problem			<0.0001					0.0008		
No difficulty	69.8% (68.4-71.2)	1.00	..	5432	6393	65.3% (64.0-66.7)	1.00	..	7385	6201
Some difficulty	52.7% (47.2-58.1)	0.68 (0.53-0.89)	..	406	540	42.8% (38.8-47.0)	0.72 (0.57-0.90)	..	749	759
Much difficulty or unable	32.5% (24.6-41.7)	0.34 (0.21-0.54)	..	155	202	34.1% (28.3-40.5)	0.56 (0.40-0.78)	..	317	316
<b>Specific health conditions</b>										
Any cardiac or vascular disease†			0.4173					0.2769		
No	68.7% (67.2-70.1)	1.00	..	5697	6738	62.7% (61.4-64.0)	1.00	..	8193	7013
Yes	46.6% (40.4-52.9)	0.88 (0.63-1.21)	..	296	398	32.4% (26.3-39.2)	0.82 (0.57-1.17)	..	256	263
Hypertension			0.4103					0.6093		
No	69.2% (67.7-70.6)	1.00	..	5401	6257	64.2% (62.9-65.5)	1.00	..	7612	6365
Yes	55.0% (50.6-59.3)	0.90 (0.70-1.16)	..	592	879	43.5% (39.9-47.2)	1.06 (0.85-1.31)	..	837	911
Diabetes			0.0384					0.1933		
No	68.5% (67.1-69.9)	1.00	..	5719	6746	62.6% (61.3-63.9)	1.00	..	8158	6973
Yes	48.3% (41.7-55.0)	0.69 (0.49-0.98)	..	274	391	38.8% (32.9-45.1)	0.80 (0.56-1.12)	..	291	303

(Continues on next page)

	Men					Women				
	Percentage reporting sexual activity	AOR	p value	Unweighted denominator	Weighted denominator	Percentage reporting sexual activity	AOR	p value	Unweighted denominator	Weighted denominator
(Continued from previous page)										
Chronic airways disease			0.0041					0.0017		
No	67.9% (66.5–69.3)	1.00	..	5931	7052	62.0% (60.8–63.3)	1.00	..	8365	7191
Yes	27.3% (16.0–42.4)	0.35 (0.17–0.71)	..	62	84	25.3% (16.8–36.3)	0.41 (0.23–0.71)	..	84	84
Arthritis			0.4949					0.0230		
No	69.0% (67.6–70.4)	1.00	..	5535	6490	65.5% (64.2–66.8)	1.00	..	7495	6248
Yes	51.3% (46.4–56.3)	0.91 (0.68–1.20)	..	458	646	37.8% (34.2–41.5)	0.76 (0.60–0.96)	..	954	1028
Broken hip or pelvis or hip replacement			0.8368					0.0886		
No	67.7% (66.3–69.1)	1.00	..	5909	7020	62.2% (60.9–63.4)	1.00	..	8326	7146
Yes	53.8% (41.8–65.4)	0.94 (0.51–1.74)	..	81	114	31.2% (22.5–41.3)	0.62 (0.35–1.08)	..	123	129
Backache, or bone or muscle disease for >3 months in past year			0.1032					0.7451		
No	67.7% (66.2–69.2)	1.00	..	5367	6286	63.4% (62.0–64.7)	1.00	..	7292	6195
Yes	65.4% (61.1–69.4)	1.23 (0.96–1.58)	..	625	849	51.5% (48.2–54.9)	1.04 (0.84–1.28)	..	1158	1081
Depressive symptoms‡			<0.0001					0.0178		
No	70.2% (68.8–71.6)	1.00	..	5214	6331	63.6% (62.3–65.0)	1.00	..	7292	6374
Yes	52.3% (47.5–57.0)	0.55 (0.43–0.72)	..	607	660	53.5% (50.0–57.0)	0.76 (0.61–0.95)	..	983	780
Prostate disease or surgery			0.8643							
No	68.0% (66.6–69.4)	1.00	..	5805	6873	..	..	..	..	..
Yes	52.4% (44.3–60.3)	0.97 (0.66–1.42)	..	185	261	..	..	..	..	..
Menopause§								0.0060		
No	..	..	..	..	..	23.7% (21.6–25.9)	1.00	..	1007	1092
Yes	..	..	..	..	..	32.8% (30.6–35.1)	0.61 (0.42–0.87)	..	1006	1418

Data in parentheses are 95% CIs. Sexual activity was defined as one or more occasion of vaginal, oral, or anal sex with a partner of the opposite sex, or oral or anal sex or genital contact with a partner of the same sex in the past 4 weeks. All models were adjusted for age and relationship status. Models investigating age and relationship status also adjusted for self-assessed general health status. Models investigating specific conditions were also adjusted for comorbidity, for which comorbidity was coded as 0=0–1 specific conditions and 1=≥2 specific conditions. AOR=adjusted odds ratio. \*Measure of comorbidity and includes arthritis, heart attack, coronary heart disease, angina, other forms of heart disease, hypertension, stroke, diabetes, broken hip or pelvis bone or hip replacement ever, backache lasting longer than 3 months, any other muscle or bone disease lasting longer than 3 months, depression, cancer, and any thyroid condition treated in the past year. †Heart attack, coronary heart disease, angina, other forms of heart disease, and stroke. ‡Respondents were asked whether they had often been bothered by feeling down, depressed, or hopeless in the past 2 weeks, and whether they had often been bothered by little interest or pleasure in doing things in the past 2 weeks, with a validated two-question patient health questionnaire (PHQ-2). §Women deemed to be postmenopausal when they had not menstruated in the past year, with analysis restricted to those aged 45–64 years.

**Table 2: Reporting of sexual activity in the past 4 weeks in relation to demographic and health characteristics, by sex**


difficulties they had experienced for at least 6 months in the past year (the duration of symptoms corresponded to criteria in the 2013 *Diagnostic and Statistical Manual of Mental Disorders*, fifth edition<sup>29</sup>). Here, we focus on trouble achieving or maintaining an erection in men and an uncomfortably dry vagina in women, because these difficulties capture both physiological and psychological elements of sexual response and are associated with ageing.<sup>30</sup> Finally, participants were divided into four categories on the basis of their relationship status at the time of interview: cohabiting with a partner (including marriage and civil partnerships); in a steady relationship (ie, expected to engage in sexual activity again with their partner) but not cohabiting with their partner; no steady relationship but reported previous cohabitation; and no steady relationship and never cohabited.

The Natsal-3 study was approved by the Oxfordshire Research Ethics Committee A (reference: 09/H0604/27). Participants provided oral informed consent for interviews.

### Statistical analysis

We did all analyses with the complex survey functions of Stata (version 12.1), accounting for stratification, clustering, and weighting of data. We included only participants who reported having had one or more sexual partner over the lifetime in our analysis, because those reporting no previous sexual experience did not complete the computer-assisted self-interview.

For sexual activity, satisfaction, response, and the perception of health effects on sex life, we report prevalences and 95% CIs in men and women by age group, relationship status, and each health variable. We weighted Natsal-3 data to adjust for the unequal probabilities of selection in terms of age and the number of adults in the eligible age range at an address. After application of these selection weights, the Natsal-3 sample was broadly representative of the British population compared with 2011 Census figures,<sup>22,31</sup> although men and London residents were slightly under-represented. Therefore, we also applied a non-response



**Figure 1: Reporting of sexual activity in past 4 weeks by sex and self-assessed general health status**

Bars show 95% CIs. Sexual activity was defined as one or more occasion of vaginal, oral, or anal sex with a partner of the opposite sex, or oral or anal sex or genital contact with a partner of the same sex in the past 4 weeks.

post-stratification weight to correct for differences in sex, age and Government Office Region between the achieved sample and the 2011 Census.

We used logistic regression models to calculate adjusted odds ratios (AORs) for associations with demographic and health variables, adjusting for age and relationship status. We additionally adjusted models investigating age and relationship status for self-assessed general health to understand the associations with age and relationship status independently of health status. We adjusted models investigating specific chronic health conditions for age, relationship status, and comorbidity (using categories of zero or one condition and more than one condition) to take account of the clustering of illnesses.

#### Role of the funding source

The sponsors of the study had no role in study design, data collection, data analysis, data interpretation, or

writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

## Results

Interviews were done with 15 162 participants (6293 men, 8869 women). 5994 men and 8452 women reported having had one or more sexual partner over the lifetime. The proportion of people in the worst self-reported health categories increased with age (table 1). The proportion of individuals reporting sexual activity in the past 4 weeks was highest in men aged 25–44 years and in women aged 25–34 years and decreased with age thereafter (table 2). The proportion of individuals reporting sexual activity in the past 4 weeks also varied by relationship status, with single people much less likely to report sexual activity than those in a steady relationship (table 2).

Overall, the proportion of men and women reporting sexual activity in the past 4 weeks was lower in those reporting bad or very bad health than in those reporting very good health, and this association remained after adjustment for age and relationship status (table 2). We recorded significant differences in the proportion reporting sexual activity in the past 4 weeks by health status in men aged 35–44 years and older, and in women aged 45–64 years (figure).

Similarly, we recorded strong associations between other broad measures of self-reported health and sexual activity in the past 4 weeks. After adjustment for age and relationship status, men and women reporting a limiting longstanding illness, two or more chronic conditions, a BMI of more than 35 kg/m<sup>2</sup>, or difficulty walking up stairs because of a health problem were significantly less likely to report sexual activity in the past 4 weeks than were those who did not report these issues (table 2). Although fewer individuals reporting specific health conditions reported sexual activity in the past 4 weeks than did those without these disorders, after adjustment for age, relationship status, and comorbidities, the associations were significant only for men and women with chronic airways disease or depressive symptoms, men with diabetes, and women with arthritis or who were postmenopausal (table 2). In sensitivity analyses, the patterns of association were broadly in the same direction and of a similar magnitude for reporting of sexual activity in the past 6 months and when the analysis was restricted to individuals aged 45–74 (data not shown).

More than 60% of men and women reported being satisfied with their sex life (table 3). Overall, after adjustment for relationship status and health, satisfaction decreased significantly after the age of 35 years in men and after the age of 25 years in women (table 3), but the association with age was weaker than for sexual activity. The percentage of men reporting sexual satisfaction was significantly lower in those reporting good, fair, or bad or very bad health than in those reporting very good health

	Men					Women				
	Percentage reporting sexual satisfaction	AOR	p value	Unweighted denominator	Weighted denominator	Percentage reporting sexual satisfaction	AOR	p value	Unweighted denominator	Weighted denominator
All	61.7% (60.3-63.1)	..	..	5933	7110	61.4% (60.2-62.7)	..	..	8428	7278
Demographic characteristics										
Age group			<0.0001					<0.0001		
16-24 years	64.8% (62.2-67.3)	1.00	..	1601	1150	68.3% (66.0-70.5)	1.00	..	1985	1109
25-34 years	68.7% (66.1-71.2)	0.88 (0.72-1.08)	..	1477	1332	68.5% (66.4-70.6)	0.70 (0.59-0.83)	..	2409	1334
35-44 years	63.1% (59.4-66.7)	0.62 (0.49-0.79)	..	788	1390	64.0% (61.0-66.9)	0.53 (0.43-0.64)	..	1179	1416
45-54 years	61.2% (57.2-65.0)	0.61 (0.47-0.78)	..	759	1355	59.8% (56.5-63.0)	0.46 (0.38-0.56)	..	1085	1395
55-64 years	54.3% (50.2-58.3)	0.50 (0.39-0.64)	..	707	1106	54.8% (51.5-58.1)	0.41 (0.33-0.50)	..	977	1174
65-74 years	53.9% (49.5-58.2)	0.51 (0.39-0.66)	..	601	778	48.9% (45.1-52.8)	0.38 (0.30-0.47)	..	793	849
Relationship status			<0.0001					<0.0001		
Living with a partner	64.9% (63.0-66.8)	1.00	..	2957	4685	67.0% (65.4-68.6)	1.00	..	4377	4709
In a steady relationship, not cohabiting	83.0% (80.0-85.6)	2.04 (1.62-2.56)	..	959	768	79.6% (76.8-82.1)	1.47 (1.22-1.78)	..	1375	797
No steady relationship, previously cohabited	38.5% (34.8-42.3)	0.37 (0.30-0.44)	..	791	704	34.9% (32.3-37.6)	0.29 (0.25-0.34)	..	1599	1131
No steady relationship, never cohabited	45.7% (42.7-48.8)	0.31 (0.26-0.37)	..	1196	929	43.7% (40.5-47.0)	0.25 (0.21-0.30)	..	1042	616
General health										
Self-reported general health status			<0.0001					0.0018		
Very good	69.5% (67.3-71.6)	1.00	..	2382	2752	65.6% (63.6-67.4)	1.00	..	3408	2902
Good	60.1% (57.8-62.3)	0.69 (0.59-0.80)	..	2502	3063	61.7% (59.7-63.6)	0.90 (0.79-1.01)	..	3533	3022
Fair	50.0% (45.8-54.1)	0.52 (0.42-0.65)	..	816	1013	53.1% (49.8-56.4)	0.75 (0.63-0.89)	..	1132	1016
Bad or very bad	45.4% (38.4-52.7)	0.51 (0.36-0.72)	..	231	280	48.6% (42.9-54.3)	0.69 (0.53-0.91)	..	355	338
Longstanding illnesses or disability			0.0052					<0.0001		
None	64.6% (63.0-66.3)	1.00	..	4209	4865	64.8% (63.3-66.3)	1.00	..	5828	4840
Non-limiting	58.2% (53.9-62.3)	0.88 (0.72-1.08)	..	798	1085	57.7% (54.4-60.9)	0.86 (0.74-1.01)	..	1130	1057
Limiting	52.5% (48.8-56.2)	0.77 (0.64-0.92)	..	924	1159	52.3% (49.4-55.1)	0.59 (0.52-0.68)	..	1469	1381
Number of self-reported chronic conditions*			0.0075					<0.0001		
0	65.1% (63.4-66.7)	1.00	..	4068	4577	67.0% (65.5-68.6)	1.00	..	5141	4142
1	58.2% (55.0-61.3)	0.84 (0.71-1.00)	..	1179	1599	58.1% (55.6-60.6)	0.77 (0.67-0.88)	..	2020	1841
≥2	51.0% (46.8-55.3)	0.74 (0.60-0.92)	..	679	929	48.1% (45.1-51.2)	0.62 (0.52-0.73)	..	1262	1291
Body-mass index			0.0045					0.1190		
Normal: 18.5-25 kg/m <sup>2</sup>	63.6% (61.4-65.8)	1.00	..	2599	2800	63.7% (61.9-65.5)	1.00	..	3329	4013
Underweight: <18.5 kg/m <sup>2</sup>	58.2% (48.1-67.6)	0.94 (0.61-1.44)	..	120	100	65.7% (59.1-71.7)	1.20 (0.88-1.64)	..	302	200
Overweight: 25-30 kg/m <sup>2</sup>	61.3% (58.9-63.6)	0.90 (0.77-1.04)	..	2007	2669	60.9% (58.5-63.3)	0.98 (0.86-1.12)	..	2084	1941
Obese: 30-35 kg/m <sup>2</sup>	61.7% (57.6-65.7)	0.95 (0.77-1.17)	..	714	967	56.2% (52.7-59.8)	0.86 (0.72-1.02)	..	935	879
Obese: >35 kg/m <sup>2</sup>	46.4% (39.5-53.6)	0.53 (0.38-0.74)	..	259	347	54.3% (49.4-59.2)	0.81 (0.64-1.02)	..	534	483
Difficulty walking up stairs because of health problem			0.0119					0.0873		
No difficulty	62.8% (61.3-64.3)	1.00	..	5366	6361	63.1% (61.7-64.4)	1.00	..	7361	6197
Some difficulty	56.1% (50.5-61.6)	1.01 (0.78-1.30)	..	413	550	53.8% (49.8-57.7)	0.94 (0.78-1.14)	..	749	764
Much difficulty or unable	42.2% (33.8-51.0)	0.61 (0.42-0.90)	..	153	198	47.6% (41.3-54.0)	0.77 (0.58-1.04)	..	317	316
Specific health conditions										
Any cardiac or vascular disease†			0.7804					0.3598		
No	62.3% (60.9-63.7)	1.00	..	5630	6706	61.9% (60.6-63.2)	1.00	..	8174	7015
Yes	51.4% (45.2-57.5)	1.04 (0.77-1.41)	..	302	404	49.3% (42.4-56.2)	1.17 (0.84-1.64)	..	251	261
Hypertension			0.3162					0.2841		
No	62.9% (61.5-64.4)	1.00	..	5334	6226	62.7% (61.3-64.0)	1.00	..	7581	6358
Yes	52.8% (48.4-57.2)	0.89 (0.71-1.12)	..	598	884	52.7% (49.0-56.3)	1.11 (0.92-1.35)	..	844	918
Diabetes			0.4124					0.3036		
No	62.3% (60.8-63.7)	1.00	..	5661	6719	62.1% (60.8-63.4)	1.00	..	8134	6968
Yes	51.7% (44.7-58.7)	0.87 (0.63-1.21)	..	271	390	46.1% (39.8-52.5)	0.85 (0.62-1.16)	..	291	307

(Continues on next page)

	Men					Women				
	Percentage reporting sexual satisfaction	AOR	p value	Unweighted denominator	Weighted denominator	Percentage reporting sexual satisfaction	AOR	p value	Unweighted denominator	Weighted denominator
(Continued from previous page)										
Chronic airways disease			0.5281					0.1941		
No	61.8% (60.3–63.2)	1.00	..	5868	7026	61.6% (60.4–62.9)	1.00	..	8341	7191
Yes	55.2% (41.8–67.9)	1.22 (0.66–2.23)	..	64	84	42.1% (31.2–53.8)	0.72 (0.44–1.18)	..	84	85
Arthritis			0.2272					0.4645		
No	62.3% (60.8–63.7)	1.00	..	5472	6464	63.5% (62.2–64.8)	1.00	..	7476	6247
Yes	55.6% (50.5–60.6)	1.18 (0.90–1.54)	..	460	646	48.9% (45.5–52.4)	0.93 (0.75–1.14)	..	949	1029
Broken hip or pelvis or hip replacement			0.4519					0.1319		
No	61.7% (60.3–63.1)	1.00	..	5847	6994	61.8% (60.5–63.0)	1.00	..	8301	7143
Yes	59.3% (47.0–70.5)	1.24 (0.71–2.14)	..	81	112	41.5% (32.4–51.3)	0.72 (0.46–1.11)	..	125	134
Backache, or bone or muscle disease for >3 months in past year			0.7165					0.0782		
No	62.1% (60.6–63.6)	1.00	..	5301	6256	63.1% (61.7–64.4)	1.00	..	7252	6179
Yes	58.7% (54.3–63.0)	1.04 (0.84–1.30)	..	630	853	52.1% (48.9–55.4)	0.85 (0.70–1.02)	..	1174	1079
Depressive symptoms‡			<0.0001					<0.0001		
No	63.8% (62.4–65.3)	1.00	..	5303	6430	64.0% (62.7–65.3)	1.00	..	7400	6466
Yes	40.6% (36.1–45.3)	0.42 (0.33–0.52)	..	621	667	40.6% (37.1–44.2)	0.41 (0.34–0.49)	..	1020	804
Prostate disease or surgery			0.3774							
No	62.1% (60.7–63.6)	1.00	..	5737	6837	..	..	..	..	..
Yes	49.9% (42.4–57.5)	0.85 (0.60–1.21)	..	191	269	..	..	..	..	..
Menopause§								0.7000		
No	..	..	..	..	..	19.6% (17.7–21.7)	1.00	..	644	831
Yes	..	..	..	..	..	37.8% (35.4–40.2)	1.05 (0.79–1.40)	..	1339	1714
Sexual activity								..		
Sexually active in the past 4 weeks			<0.0001					<0.0001		
No	37.1% (34.8–39.6)	1.00	..	1995	2174	38.0% (35.9–40.0)	1.00	..	3035	2645
Yes	73.7% (71.9–75.4)	3.69 (3.13–4.35)	..	3801	4789	75.9% (74.5–77.3)	3.73 (3.22–4.32)	..	5184	4465

Data in parentheses are 95% CIs. All models were adjusted for age and relationship status. Models investigating age and relationship status also adjusted for self-assessed general health status. Models investigating specific conditions were also adjusted for comorbidity, for which comorbidity was coded as 0=0–1 specific conditions and 1=≥2 specific conditions. AOR=adjusted odds ratio. \*Measure of comorbidity and includes arthritis, heart attack, coronary heart disease, angina, other forms of heart disease, hypertension, stroke, diabetes, broken hip or pelvis bone or hip replacement ever, backache lasting longer than 3 months, any other muscle or bone disease lasting longer than 3 months, depression, cancer, and any thyroid condition treated in the past year. †Heart attack, coronary heart disease, angina, other forms of heart disease, and stroke. ‡Respondents were asked whether they had often been bothered by feeling down, depressed, or hopeless in the past 2 weeks, and whether they had often been bothered by little interest or pleasure in doing things in the past 2 weeks, with a validated two-question patient health questionnaire (PHQ-2). §Women deemed to be postmenopausal when they had not menstruated in the past year, with analysis restricted to those aged 45–64 years.

Table 3: Reporting of satisfaction with sex life in the past year in relation to demographic and health characteristics, by sex

(table 3). The percentage of women reporting sexual satisfaction was also significantly lower in those reporting fair, bad, or very bad health than in those reporting very good health (table 3). Similar associations were recorded for other broad measures of health (table 3). However, after adjustment for age, relationship status, and comorbidities, depressive symptoms was the only specific health condition to be significantly associated with low sexual satisfaction (table 3).

The percentage of individuals reporting sexual response problems increased with age: erectile difficulties were most common in men aged 65–74 years (affecting 26.2%, 95% CI 21.5–31.6) and vaginal dryness was most common in women aged 55–64 years (affecting 22.1%, 18.7–25.9; appendix). After adjustment for age and relationship status, participants with poorer general health, a limiting disability, or two or more chronic conditions were more likely than those who did

not report these issues to have erectile difficulties or vaginal dryness (appendix). For specific health conditions in men, reporting of erectile difficulties was associated only with depressive symptoms after adjustment for age, relationship status, and comorbidities. In women, reporting of vaginal dryness was associated with a previous broken hip or pelvis or a hip replacement, depressive symptoms, being postmenopausal, and backache or bone or muscle disease for more than 3 months in the past year (appendix).

The percentage of male participants reporting that a health condition had affected their sex life in the past year increased in each successive age group (table 4). In women, the pattern was different: the percentage was highest in those aged 45–54 years, and lowest in those aged 16–24 years and 65–74 years (table 5). After adjustment for age and relationship status, we recorded strong associations in men and women between

reporting that a health condition affected one's sex life and poorer self-assessed general health, limiting disabilities, one or more comorbidity, and difficulty with walking up stairs (tables 4, 5). By contrast with the small number of specific chronic conditions associated with sexual activity, satisfaction, and sexual response problems, reporting of an effect of health conditions on one's sex life was associated with a range of chronic health conditions: cardiac or vascular disease, chronic airways disease, backache, and depressive symptoms in

both men and women; diabetes and prostate disease in men; and arthritis and hip or pelvis fracture or hip replacement in women (tables 4, 5).

Overall, 7.0% (95% CI 6.2–7.8) of men and 6.7% (6.1–7.4) of women reported having sought clinical help or advice about their sex life. Seeking clinical help was more common in individuals reporting that a health condition affected their sex lives than in those who did not (men: 23.5% [95% CI 20.3–26.9] vs 3.7% [3.2–4.4]; women: 18.4% [17.5–22.8] vs 4.3% [3.8–4.9]). Most

	All men			Men whose health affects sex life				
	Percentage reporting that their health affects sex life	Adjusted odds ratio	p value	Unweighted denominator	Weighted denominator	Percentage reporting that they sought clinical advice*	Unweighted denominator	Weighted denominator
All	16.6% (15.4–17.7)	..	..	5621	6870	23.5% (20.3–26.9)	845	1134
Demographic characteristics								
Age group			<0.0001					
16–24 years	6.1% (4.9–7.7)	1.00	..	1362	995	11.3% (5.7–21.2)	87	61
25–34 years	10.6% (9.1–12.3)	1.64 (1.20–2.24)	..	1445	1296	15.9% (10.7–23.1)	161	137
35–44 years	14.2% (11.7–17.0)	2.10 (1.47–3.01)	..	782	1378	23.6% (16.1–33.1)	121	195
45–54 years	18.1% (15.2–21.4)	2.48 (1.76–3.49)	..	749	1343	25.2% (17.8–34.4)	139	240
55–64 years	25.4% (22.0–29.0)	3.03 (2.17–4.21)	..	698	1095	25.5% (19.2–33.1)	178	278
65–74 years	29.2% (25.2–33.5)	3.83 (2.73–5.38)	..	585	762	26.9% (20.3–34.6)	159	222
Relationship status			0.4449					
Living with a partner	17.5% (16.0–19.1)	1.00	..	2944	4665	23.5% (19.6–27.8)	496	813
In a steady relationship, not cohabiting	12.6% (10.2–15.4)	1.05 (0.80–1.39)	..	955	766	34.6% (24.2–46.8)	109	96
No steady relationship, previously cohabited	21.5% (18.4–25.0)	0.86 (0.67–1.11)	..	796	709	22.0% (15.0–31.1)	159	152
No steady relationship, never cohabited	9.9% (7.8–12.5)	0.84 (0.62–1.14)	..	920	725	11.4% (5.7–21.5)	81	72
General health								
Self-reported health status			<0.0001					
Very good	7.6% (6.4–9.1)	1.00	..	2241	2638	21.7% (15.1–30.3)	154	201
Good	14.6% (13.0–16.3)	1.88 (1.49–2.38)	..	2367	2963	22.7% (17.8–28.6)	311	432
Fair	33.8% (30.2–37.6)	5.05 (3.90–6.54)	..	787	993	24.9% (19.2–31.6)	250	332
Bad or very bad	60.7% (53.0–67.9)	14.57 (9.86–21.52)	..	224	273	24.8% (17.4–34.0)	129	166
Longstanding illness or disability			<0.0001					
None	9.0% (8.0–10.2)	1.00	..	3957	4669	20.5% (15.7–26.4)	319	421
Non-limiting	21.2% (18.1–24.7)	2.19 (1.70–2.81)	..	765	1061	26.0% (18.9–34.6)	154	225
Limiting	43.1% (39.4–46.9)	6.22 (4.99–7.76)	..	897	1139	24.8% (20.3–29.9)	372	488
Number of self-reported chronic conditions†			<0.0001					
0	8.3% (7.3–9.4)	1.00	..	3812	4381	17.7% (12.8–24.1)	297	364
1	22.7% (20.2–25.5)	2.82 (2.26–3.52)	..	1123	1540	27.3% (21.5–34.1)	253	347
≥2	44.6% (40.6–48.8)	7.16 (5.59–9.17)	..	686	948	25.2% (20.2–30.9)	295	423
Body-mass index			<0.0001					
Normal: 18.5–25 kg/m <sup>2</sup>	13.4% (11.8–15.1)	1.00	..	2428	2669	22.3% (17.2–28.3)	296	356
Underweight: <18.5 kg/m <sup>2</sup>	12.3% (6.4–22.4)	1.28 (0.62–2.65)	..	95	84	33.0% (7.8–74.3)	11	10
Overweight: 25–30 kg/m <sup>2</sup>	16.4% (14.6–18.4)	1.06 (0.86–1.30)	..	1948	2615	25.1% (20.0–31.1)	299	428
Obese: 30–35 kg/m <sup>2</sup>	20.2% (17.1–23.8)	1.30 (1.01–1.68)	..	696	954	25.8% (18.4–34.9)	133	190
Obese: >35 kg/m <sup>2</sup>	31.4% (25.4–38.2)	2.27 (1.62–3.17)	..	254	342	22.2% (13.5–34.2)	77	107
Difficulty walking up stairs because of health problem			<0.0001					
No difficulty	12.6% (11.5–13.8)	1.00	..	5069	6133	23.9% (20.0–28.3)	584	773
Some difficulty	43.4% (38.0–49.1)	4.06 (3.13–5.28)	..	401	542	24.7% (18.3–32.5)	169	232
Much difficulty or unable	65.6% (56.9–73.3)	9.80 (6.51–14.76)	..	150	193	17.7% (10.4–28.4)	91	127

(Continues on next page)

	All men			Men whose health affects sex life				
	Percentage reporting that their health affects sex life	Adjusted odds ratio	p value	Unweighted denominator	Weighted denominator	Percentage reporting that they sought clinical advice*	Unweighted denominator	Weighted denominator
(Continued from previous page)								
Specific health conditions								
Any cardiac or vascular disease‡			<0.0001					
No	14.9% (13.8–16.0)	1.00	..	5376	6528	22.0% (18.7–25.7)	735	967
Yes	48.8% (42.0–55.6)	2.14 (1.57–2.91)	..	244	341	31.9% (23.5–41.7)	110	166
Hypertension			0.7823					
No	14.1% (13.0–15.3)	1.00	..	5031	5994	23.6% (19.9–27.7)	652	843
Yes	33.2% (29.1–37.6)	1.04 (0.81–1.32)	..	589	876	23.0% (17.4–29.8)	193	291
Diabetes			0.0045					
No	15.1% (14.0–16.3)	1.00	..	5357	6486	22.6% (19.2–26.3)	732	976
Yes	41.1% (34.7–47.9)	1.55 (1.15–2.09)	..	263	383	28.8% (20.3–39.2)	113	158
Chronic airways disease			<0.0001					
No	16.1% (15.0–17.2)	1.00	..	5558	6787	23.8% (20.6–27.4)	812	1087
Yes	56.5% (42.0–69.9)	4.12 (2.09–8.12)	..	62	83	15.1% (6.1–32.8)	33	47
Arthritis			0.5600					
No	14.8% (13.6–16.0)	1.00	..	5172	6235	24.1% (20.6–28.0)	694	917
Yes	34.1% (29.4–39.2)	1.08 (0.83–1.41)	..	448	635	20.6% (14.4–28.5)	151	217
Broken hip or pelvis or hip replacement			0.1012					
No	16.1% (15.0–17.3)	1.00	..	5536	6754	23.2% (20.0–26.8)	813	1086
Yes	41.4% (29.9–53.9)	1.55 (0.92–2.61)	..	81	112	29.9% (15.9–49.0)	31	46
Backache, or bone or muscle disease for >3 months in past year			0.0277					
No	14.2% (13.1–15.4)	1.00	..	5013	6035	24.9% (21.1–29.0)	649	855
Yes	33.3% (29.3–37.5)	1.31 (1.03–1.67)	..	606	834	19.1% (13.8–25.9)	195	278
Depressive symptoms§			<0.0001					
No	14.8% (13.7–16.0)	1.00	..	5027	6215	22.6% (19.1–26.5)	664	916
Yes	33.7% (29.4–38.4)	2.62 (2.05–3.34)	..	582	638	26.6% (19.8–34.8)	179	215
Prostate disease or surgery			0.0026					
No	15.7% (14.6–16.9)	1.00	..	5427	6598	22.6% (19.4–26.3)	768	1032
Yes	37.6% (30.3–45.5)	1.82 (1.23–2.68)	..	190	269	32.0% (22.0–43.9)	76	101
Sexual activity								
Sexually active in the past 4 weeks			0.0275					
No	21.2% (19.0–23.5)	1.00	..	1704	1952	20.3% (15.7–25.9)	314	413
Yes	14.3% (13.0–15.7)	0.80 (0.65–0.97)	..	3780	4769	25.1% (21.0–29.7)	492	682
Satisfied with sex life			<0.0001					
No	23.6% (21.5–25.8)	1.00	..	2099	2586	27.4% (23.1–32.3)	458	610
Yes	12.3% (11.0–13.6)	0.50 (0.42–0.59)	..	3510	4269	18.8% (14.7–23.8)	387	524

Data in parentheses are 95% CIs. All models were adjusted for age and relationship status. Models investigating age and relationship status also adjusted for self-assessed general health status. Models investigating specific conditions were also adjusted for comorbidity, for which comorbidity was coded as 0=0–1 specific conditions and 1=≥2 specific conditions. \*General practitioner, sexual health clinic, other clinics or doctors, or psychiatrist or psychologist. †Measure of comorbidity and includes arthritis, heart attack, coronary heart disease, angina, other forms of heart disease, hypertension, stroke, diabetes, broken hip or pelvis bone or hip replacement ever, backache lasting longer than 3 months, any other muscle or bone disease lasting longer than 3 months, depression, cancer, and any thyroid condition treated in the past year. ‡Heart attack, coronary heart disease, angina, other forms of heart disease, and stroke. §Respondents were asked whether they had often been bothered by feeling down, depressed, or hopeless in the past 2 weeks, and whether they had often been bothered by little interest or pleasure in doing things in the past 2 weeks, with a validated two-question patient health questionnaire (PHQ-2).

**Table 4: Reporting of health conditions affecting sexual activity or enjoyment in the past year and whether clinical advice has been sought, in relation to demographic and health characteristics of men**

participants who reported seeking help asked a family doctor for advice (men: 85.4%, 95% CI 78.7–90.3; women: 80.7%, 74.9–85.4), with smaller proportions reporting visiting a sexual health clinic (men: 7.6%, 4.5–12.5; women: 11.2%, 8.0–15.5), another type of

clinic (men: 6.8%, 4.0–11.3; women: 8.8%, 5.5–13.7), or a psychiatrist or psychologist (men: 6.5%, 3.2–12.5; women: 11.3%, 7.6–16.4; data not shown). A higher proportion of men than women who had reported that their health affected their sex life reported that they had

	All women			Women whose health affects sex life				
	Percentage reporting that their health affects sex life	Adjusted odds ratio	p value	Unweighted denominator	Weighted denominator	Percentage reporting that they sought clinical advice*	Unweighted denominator	Weighted denominator
All	17.2% (16.3–18.2)	..	..	8097	7071	18.4% (16.0–20.9)	1322	1215
Demographic characteristics								
Age group			<0.0001					
16–24 years	13.2% (11.5–15.2)	1.00	..	1716	957	24.4% (18.7–31.2)	223	126
25–34 years	18.5% (16.8–20.4)	1.27 (1.03–1.56)	..	2376	1312	19.9% (15.9–24.6)	425	243
35–44 years	16.6% (14.4–19.0)	1.02 (0.80–1.29)	..	1174	1404	16.4% (11.7–22.4)	190	233
45–54 years	19.6% (17.1–22.3)	1.03 (0.81–1.31)	..	1064	1374	16.8% (11.9–23.2)	218	269
55–64 years	18.8% (16.2–21.7)	0.86 (0.66–1.12)	..	975	1175	22.5% (16.3–30.1)	165	219
65–74 years	14.8% (12.1–17.9)	0.58 (0.43–0.77)	..	792	848	9.2% (4.8–16.9)	101	125
Relationship status			<0.0001					
Living with a partner	19.1% (17.8–20.4)	1.00	..	4357	4678	18.5% (15.6–21.7)	809	889
In a steady relationship, not cohabiting	17.2% (14.9–19.8)	0.81 (0.66–1.00)	..	1373	796	19.2% (14.2–25.5)	220	137
No steady relationship, previously cohabited	12.2% (10.5–14.0)	0.42 (0.35–0.52)	..	1620	1153	17.1% (12.5–22.9)	214	140
No steady relationship, never cohabited	10.9% (8.5–14.0)	0.42 (0.32–0.57)	..	738	439	17.7% (10.2–29.0)	78	48
General health								
Self-reported health status			<0.0001					
Very good	9.6% (8.4–10.8)	1.00	..	3245	2799	21.4% (16.3–27.5)	299	267
Good	15.3% (14.0–16.7)	1.80 (1.51–2.15)	..	3392	2928	18.0% (14.5–22.1)	516	448
Fair	30.6% (27.7–33.8)	4.99 (4.05–6.15)	..	1110	1009	20.5% (15.9–26.0)	319	307
Bad or very bad	57.5% (51.5–63.4)	17.29 (12.96–23.05)	..	350	335	11.7% (7.8–17.1)	188	193
Longstanding illness or disability			<0.0001					
None	11.3% (10.3–12.3)	1.00	..	5550	4666	18.7% (15.3–22.7)	621	525
Non-limiting	14.3% (12.2–16.8)	1.48 (1.19–1.85)	..	1107	1042	25.2% (18.4–33.5)	155	149
Limiting	39.8% (36.8–42.8)	6.29 (5.28–7.50)	..	1439	1363	16.2% (13.0–19.8)	546	540
Number of self-reported chronic conditions†			<0.0001					
0	10.6% (9.6–11.7)	1.00	..	4865	3976	18.2% (14.5–22.5)	510	421
1	20.0% (18.0–22.2)	2.47 (2.06–2.96)	..	1953	1788	20.0% (16.0–24.7)	408	358
≥2	33.5% (30.6–36.4)	6.26 (5.11–7.68)	..	1278	1306	17.2% (13.5–21.7)	404	435
Body-mass index			0.0039					
Normal: 18.5–25 kg/m <sup>2</sup>	16.5% (15.1–18.0)	1.00	..	3844	3220	20.0% (16.5–24.1)	601	532
Underweight: <18.5 kg/m <sup>2</sup>	18.8% (13.8–25.1)	1.32 (0.89–1.95)	..	261	178	17.3% (7.4–35.4)	46	33
Overweight: 25–30 kg/m <sup>2</sup>	15.9% (14.1–17.8)	0.93 (0.78–1.10)	..	2039	1906	14.2% (10.5–18.9)	311	302
Obese: 30–35 kg/m <sup>2</sup>	20.4% (17.4–23.6)	1.28 (1.03–1.60)	..	920	874	22.3% (15.9–30.4)	178	176
Obese: >35 kg/m <sup>2</sup>	22.0% (17.9–26.6)	1.43 (1.09–1.88)	..	529	482	19.3% (12.5–28.6)	110	105
Difficulty walking up stairs because of health problem			<0.0001					
No difficulty	13.8% (12.9–14.8)	1.00	..	7041	5995	20.2% (17.3–23.5)	949	827
Some difficulty	30.1% (26.5–34.0)	3.34 (2.72–4.11)	..	742	760	19.6% (14.6–25.9)	225	227
Much difficulty or unable	51.3% (44.8–57.7)	8.81 (6.65–11.68)	..	313	315	7.0% (4.0–12.1)	148	161
Specific health conditions								
Any cardiac or vascular disease‡			0.0043					
No	16.8% (15.8–17.8)	1.00	..	7900	6864	18.7% (16.3–21.3)	1269	1148
Yes	32.1% (25.0–40.2)	1.71 (1.18–2.46)	..	195	205	12.4% (4.8–28.3)	52	66
Hypertension			0.1057					
No	16.5% (15.5–17.6)	1.00	..	7262	6161	18.9% (16.4–21.7)	1162	1018
Yes	21.8% (18.7–25.2)	0.83 (0.67–1.04)	..	833	909	15.5% (10.0–23.1)	159	196
Diabetes			0.0637					
No	16.7% (15.7–17.7)	1.00	..	7808	6766	18.1% (15.7–20.7)	1243	1125
Yes	29.2% (23.5–35.7)	1.35 (0.98–1.85)	..	287	304	21.8% (13.9–32.5)	78	89

(Continues on next page)

	All women			Women whose health affects sex life				
	Percentage reporting that their health affects sex life	Adjusted odds ratio	p value	Unweighted denominator	Weighted denominator	Percentage reporting that they sought clinical advice*	Unweighted denominator	Weighted denominator
(Continued from previous page)								
Chronic airways disease			0.0002					
No	16.9% (16.0-17.9)	1.00	..	8009	6983	18.5% (16.2-21.2)	1292	1180
Yes	38.9% (27.7-51.4)	2.69 (1.62-4.49)	..	86	86	11.0% (3.7-28.5)	29	34
Arthritis			0.0177					
No	15.6% (14.7-16.7)	1.00	..	7146	6038	19.2% (16.6-22.2)	1078	942
Yes	26.4% (23.3-29.8)	1.29 (1.05-1.60)	..	949	1031	15.2% (10.9-20.8)	243	272
Broken hip or pelvis or hip replacement			0.0069					
No	16.9% (15.9-17.9)	1.00	..	7972	6939	18.9% (16.5-21.5)	1285	1168
Yes	35.0% (25.8-45.4)	1.85 (1.19-2.88)	..	122	130	6.3% (1.9-18.6)	36	46
Backache, or bone or muscle disease for >3 months in past year			<0.0001					
No	14.5% (13.5-15.5)	1.00	..	6942	5988	18.6% (15.9-21.6)	962	867
Yes	32.3% (29.3-35.5)	1.70 (1.40-2.06)	..	1153	1081	17.8% (13.7-22.8)	360	347
Depressive symptoms‡			<0.0001					
No	14.9% (13.9-15.9)	1.00	..	7102	6278	17.7% (15.1-20.7)	1002	935
Yes	35.8% (32.2-39.5)	2.75 (2.28-3.33)	..	985	781	20.6% (16.2-25.9)	320	279
Menopause¶			0.2097					
No	16.9% (14.0-20.3)	1.00	..	632	822	9.6% (4.8-18.2)	109	139
Yes	20.3% (18.1-22.8)	1.25 (0.88-1.76)	..	1389	1705	23.1% (18.0-29.1)	270	345
Sexual activity								
Sexually active in the past 4 weeks			0.0681					
No	16.8% (15.2-18.5)	1.00	..	2736	2469	14.7% (11.3-18.9)	409	414
Yes	17.2% (16.0-18.4)	0.85 (0.71-1.01)	..	5168	4449	20.6% (17.6-24.1)	866	760
Satisfied with sex life			<0.0001					
No	23.2% (21.5-25.0)	1.00	..	2999	2669	20.6% (17.3-24.3)	669	619
Yes	13.5% (12.5-14.7)	0.45 (0.39-0.52)	..	5061	4365	16.0% (13.0-19.5)	650	591

Data in parentheses are 95% CIs. All models were adjusted for age and relationship status. Models investigating age and relationship status also adjusted for self-assessed general health status. Models investigating specific conditions were also adjusted for comorbidity, for which comorbidity was coded as 0=0-1 specific conditions and 1=≥2 specific conditions. \*General practitioner, sexual health clinic, other clinics or doctors, or psychiatrist or psychologist. †Measure of comorbidity and includes arthritis, heart attack, coronary heart disease, angina, other forms of heart disease, hypertension, stroke, diabetes, broken hip or pelvis bone or hip replacement ever, backache lasting longer than 3 months, any other muscle or bone disease lasting longer than 3 months, depression, cancer, and any thyroid condition treated in the past year. ‡Heart attack, coronary heart disease, angina, other forms of heart disease, and stroke. §Respondents were asked whether they had often been bothered by feeling down, depressed, or hopeless in the past 2 weeks, and whether they had often been bothered by little interest or pleasure in doing things in the past 2 weeks, with a validated two-question patient health questionnaire (PHQ-2). ¶Women deemed to be postmenopausal when they had not menstruated in the past year, with analysis restricted to those aged 45-64 years.

**Table 5: Reporting of health conditions affecting sexual activity or enjoyment in the past year and whether clinical advice has been sought, in relation to demographic and health characteristics of women**

sought clinical advice (tables 4, 5). However, the proportions did not vary substantially or consistently by health status (tables 4, 5).

**Discussion**

In our large, population-based study, we have shown that poorer physical health, limiting disabilities, functional impairment, and depressive symptoms are associated with decreased sexual activity and sexual satisfaction and increased reporting of sexual response problems (erectile difficulties and vaginal dryness) in men and women aged 16-74 years in Britain (panel). Furthermore, about 17% of all men and women reported that their health had affected their sex life in the past year. This proportion rose to roughly 60% in participants with bad or very bad health. Less than a quarter of men and a fifth of women who

reported that their health affected their sex life had sought clinical help, which was usually from general practitioners rather than specialists. Overall, we have identified strong associations between health and sexual lifestyles, and established that many people are aware of an effect of ill health on their sex life.

Our sample encompasses most of the sexually active lifespan, adding to previous work by emphasising the differences in reported sexual activity according to health status, including at younger ages. For example, the proportion of men aged 35-44 years who reported sexual activity in the past 4 weeks was much lower for those with fair, bad, or very bad health than for those with very good or good health, and similar to the proportion in men reporting good or very good health but 20 years older. We recorded a similar pattern in women, although we noted

no difference by health status in the oldest women. This finding could be a result of the low prevalence of sexual activity in older women, and possibly related to their partnership status and their partners' age and health. An area of future research will be to investigate the role of partners' characteristics—eg, through studies of both partners in a relationship. Nevertheless, about a third of men and women in bad or very bad health reported sexual activity in the past 4 weeks, and just less than half of the same group reported satisfaction with their sex lives. Beckman and colleagues<sup>21</sup> suggested that more accepting attitudes have contributed to increased reporting of sexual activity in older people; the message that poor health need not mean the end of an active or satisfying sex life is an important one.

Although we sought to minimise reporting bias by using computer-assisted self-interview for the most sensitive questions in addition to non-response weighting,<sup>30</sup> our cross-sectional data should be interpreted with caution. We relied on self-reported diagnoses, which can lead to misclassification errors, although previous studies have shown good levels of agreement between self-reported chronic disease and medical records,<sup>32,33</sup> and self-report remains the primary method to assess health status in many large population surveys. Furthermore, the patterns of disease prevalence that we recorded by age and sex were similar to those reported in the Health Survey for England.<sup>34</sup> Participants with undiagnosed or preclinical disease could not be identified in our study, which could have meant the effect size was underestimated. Our study did not include physical or biological measures of health, which can be used to validate self-reporting.<sup>35</sup> Additionally, we were unable to account for specific drugs or partners' health status, or to explore severity within specific conditions, and we cannot attribute directional causality.

The Natsal-3 sample was broadly representative of the general population of Britain in terms of self-reported general health, marital status, and ethnic origin, on the basis of 2011 census data, after weighting for key sociodemographic characteristics.<sup>22,31</sup> However, the data are susceptible to participation biases—eg, poor health could affect willingness to participate—and the sampling frame did not include individuals in residential or nursing care. Although we have referred to the life course, our findings are not generalisable to people older than 74 years. Our findings at different ages could be partly affected by differences in sexual behaviour between generations.<sup>5</sup> In the future, longitudinal studies and investigations including biological and physical measures of health will be important in the refinement of our understanding and establishment of causality. Nevertheless, the strength and novelty of our study lie in its size, population representativeness, and broad range of detailed sexual lifestyle measures. Importantly, we could explore the associations between sexual lifestyles and health, with measures of both broad health and specific conditions, independently of the key confounders of age

### Panel: Research in context

#### Systematic review

A range of health conditions, disabilities, and drugs affects sexual health and lifestyles.<sup>7,11</sup> The effects are important throughout life, but could be particularly important in older age, when chronic ill health is most common. However, population-based studies in which these issues have been investigated have often had narrow age ranges, few measures of health, and inconsistent findings.<sup>3</sup> Moreover, these issues are seldom discussed in clinical practice,<sup>11</sup> and the evidence base to guide clinical management and self-help is small.

#### Interpretation

As far as we are aware, our analysis is the first in which the relationships between sexual lifestyles and health have been studied across a wide age range in men and women, and with a broad range of self-reported health measures. We have shown that health and sexual lifestyles are associated in most sexually active age groups, even after adjustment for age and relationship status. However, we have also shown that many people in poor health and at older ages report an active or satisfying sex life, or both. Many people reported that health conditions had had an effect on their sex lives in the past year—an effect that persisted into older age. This association was significantly higher in participants reporting ill health. However, of individuals reporting a health condition that affected their sex life, only a quarter of men and a fifth of women had sought clinical help or advice. Our findings should help clinicians, their patients, and policy makers to consider the continuation of sexual activity and enjoyment in the face of ill health. Practitioners should consider giving appropriate advice about sexual lifestyles to promote the overall wellbeing of patients with chronic conditions.

and partnership status. This Article builds a compelling and consistent picture showing the strong associations between sexual lifestyles and health across the life course.

Although our findings are consistent with previous reports,<sup>4,8,10,13,14,36</sup> not all studies have reached the same conclusions about the associations with general health, particularly in women. A large Spanish cross-sectional survey of people older than 65 years<sup>8</sup> showed that sexual activity was associated with self-assessed general health status in men but not women. A US study<sup>11</sup> showed that poor physical health did not contribute to the age-related decrease in sexual frequency in women aged 44–72 years, although health did contribute in men. Additionally, another US study<sup>37</sup> showed that ill health was unrelated to frequency of sexual activity in retired men and women older than 45 years. However, these studies were restricted in age range, and, by including a wide age range and using a large sample size, we were able to identify the associations with ill health separately from those with age. With adjustment for age and additionally for the availability of partners and comorbidities, we noted strong associations between sexual lifestyles and some specific chronic health conditions. Our data support other studies linking mental ill health and sexual problems.<sup>8,13,14</sup> Diabetes was associated with reduced sexual activity in men but not women, but we note other studies showed associations only for women,<sup>4,17</sup> which could be a result of differences in methods used. In our study, hypertension was not associated with sexual activity, satisfaction, or sexual responses, although other studies—in some of which investigators used sphygmomanometry—have shown

that sexual activity is reduced in women with hypertension,<sup>18</sup> but not in men.<sup>13,18</sup>

To our knowledge, no other large population survey of sexual lifestyles has investigated whether people felt that a health condition affected their sexual activity or enjoyment. Reporting of a health condition that affected an individual's sex life increased in each age group in men but not women. These findings are consistent with previous work suggesting that an impaired sexual response is less likely to be problematic for older women than younger women.<sup>38</sup>

Most people reporting a condition affecting their sex life in the past year had not sought clinical advice. When they had, they consulted general practitioners or family doctors and seldom specialists, which corroborates findings that general practitioners are often the first point of contact for sexual concerns.<sup>39</sup> Evidence suggests that, although many major health conditions are recognised to affect sexual function,<sup>40</sup> clinical advice about the effect of ill health on sexual activity is often not given, even when the condition is gynaecological or genital,<sup>41,42</sup> and doctors rarely ask patients about their sexual function.<sup>43</sup> Not only are patients sometimes unwilling to discuss problems, but health-care professionals might also have poor awareness and little training about advising and communicating with patients about sexual problems.<sup>44</sup> This issue needs to be addressed, because sexual problems are common and can cause distress,<sup>30</sup> can indicate underlying physical or mental health problems, and can be caused iatrogenically (eg, by drugs or surgery).<sup>45</sup> Our findings indicate that many patients with chronic ill health are already aware that their health has an effect on their sex lives. Therefore, awareness needs to increase, guidance improve, and communication skills built so that problems are acknowledged, ameliorated, and openly discussed in appropriate ways.

On the one hand, people are living longer and have expectations of continued sexual fulfilment as they age.<sup>46,47</sup> But on the other hand, ill health increases in old age; the proportion of the English population reporting one or more chronic conditions rises from 17% in individuals younger than 40 years to 60% in those older than 60 years.<sup>48</sup> The sexual health of older people has been a neglected area of research.<sup>44,47,49</sup> More recent public health policy, such as a WHO 2013 report,<sup>50</sup> reflects an increasing awareness of the interdependence between sexual health and general health. A UK Department of Health 2013 report<sup>51</sup> drew attention to the effects of long-term health conditions on erectile function and the effect of cancer on sexual health in older age. Our study suggests that strategic frameworks could go further and include associations between health—eg, disabilities or mental health—and sexual lifestyles at younger ages, as well as the broader associations between sexual lifestyles and chronic ill health.

Sexual lifestyles are strongly linked to overall health and wellbeing. The finding that many people who felt that their health affected their sex life seldom sought

clinical help suggests an unmet clinical need that is inadequately recognised in present medical practice and health policy. Our data should help to challenge stereotypes and inform sexual policy for all ages.

#### Contributors

NF, CHM, PS, and AMJ conceived this Article. NF wrote the first draft of the Article, with further contributions from CHM, PS, CT, SC, KRM, BE, WM, FW, JD, KW, and AMJ. NF did statistical analyses, with support from CHM, CT, KGJ, and PP. CHM, PS, BE, WM, FW, AJC, KW, and AMJ, initial applicants for Natsal-3, wrote the study protocol and obtained funding. NF, CHM, PS, CT, SC, KRM, BE, WM, FW, JD, AJC, AP, KW, and AMJ designed the Natsal-3 questionnaire, applied for ethics approval, and undertook piloting of the questionnaire. SC, BE, and AP were responsible for data collection and delivery. CHM, CT, SC, PP, and AP managed data. All authors contributed to data interpretation, reviewed successive drafts, and approved the final version of the report.

#### Conflicts of interest

AMJ has been a Governor of the Wellcome Trust since 2011. The other authors declare that they have no conflicts of interest.

#### Acknowledgments

Natsal-3 is a collaboration between University College London (London, UK), the London School of Hygiene and Tropical Medicine (London, UK), NatCen Social Research, Public Health England (formerly the Health Protection Agency), and the University of Manchester (Manchester, UK). The study was supported by grants from the Medical Research Council (G0701757) and the Wellcome Trust (084840), with contributions from the Economic and Social Research Council and Department of Health. NF is supported by a National Institute for Health Research Academic Clinical Lectureship, and KGJ is supported by a National Institute for Health Research Methodology Fellowship. We thank the study participants; the team of interviewers from NatCen Social Research who did the interviews; Heather Wardle, Vicki Hawkins, Cathy Coshall, and operations and computing staff from NatCen Social Research; and Graham Hart and the reviewers for valuable comments on early drafts of the report.

#### References

- DeLamater J. Sexual expression in later life: a review and synthesis. *J Sex Res* 2012; **49**: 125–41.
- Davison SL, Bell RJ, LaChina M, Holden SL, Davis SR. The relationship between self-reported sexual satisfaction and general well-being in women. *J Sex Med* 2009; **6**: 2690–97.
- Tan HM, Tong SF, Ho CCK. Men's health: sexual dysfunction, physical, and psychological health—is there a link? *J Sex Med* 2012; **9**: 663–71.
- Lindau ST, Schumm LP, Laumann EO, Levinson W, O'Muirheartaigh CA, Waite LJ. A study of sexuality and health among older adults in the United States. *N Engl J Med* 2007; **357**: 762–74.
- Mercer CH, Tanton C, Prah P, et al. Changes in sexual attitudes and lifestyles in Britain through the life course and trends over time: findings from the National Surveys of Sexual Attitudes and Lifestyles (Natsal). *Lancet* 2013; published online Nov 26. [http://dx.doi.org/10.1016/S0140-6736\(13\)62035-8](http://dx.doi.org/10.1016/S0140-6736(13)62035-8).
- Johnson AM, Wadsworth J, Field J, Wellings K. Sexual attitudes and lifestyles. Oxford: Blackwell, 1994.
- Johnson AM, Mercer CH, Erens B, et al. Sexual behaviour in Britain: partnerships, practices, and HIV risk behaviours. *Lancet* 2001; **358**: 1835–42.
- Palacios-Ceña D, Carrasco-Garrido P, Hernández-Barrera V, Alonso-Blanco C, Jiménez-García R, Fernández-de-las-Peñas C. Sexual behaviors among older adults in Spain: results from a population-based national sexual health survey. *J Sex Med* 2012; **9**: 121–29.
- Herbenick D, Reece M, Schick V, Sanders SA, Dodge B, Fortenberry JD. Sexual behavior in the United States: results from a national probability sample of men and women ages 14–94. *J Sex Med* 2010; **7** (suppl 5): 255–65.
- Richters J, Grulich AE, de Visser RO, Smith AMA, Rissel CE. Sex in Australia: sexual difficulties in a representative sample of adults. *Aust N Z J Public Health* 2003; **27**: 164–70.

- 11 Karraker A, Delamater J, Schwartz CR. Sexual frequency decline from midlife to later life. *J Gerontol B Psychol Sci Soc Sci* 2011; **66**: 502–12.
- 12 Kontula O, Haavio-Mannila E. The impact of aging on human sexual activity and sexual desire. *J Sex Res* 2009; **46**: 46–56.
- 13 Corona G, Lee DM, Forti G, et al. Age-related changes in general and sexual health in middle-aged and older men: results from the European Male Ageing Study (EMAS). *J Sex Med* 2010; **7**: 1362–80.
- 14 Hyde Z, Flicker L, Hankey GJ, et al. Prevalence of sexual activity and associated factors in men aged 75 to 95 years: a cohort study. *Ann Intern Med* 2010; **153**: 693–702.
- 15 Bacon CG, Mittleman MA, Kawachi I, Giovannucci E, Glasser DB, Rimm EB. Sexual function in men older than 50 years of age: results from the health professionals follow-up study. *Ann Intern Med* 2003; **139**: 161–68.
- 16 Laumann EO, Paik A, Glasser DB, et al. A cross-national study of subjective sexual well-being among older women and men: findings from the Global Study of Sexual Attitudes and Behaviors. *Arch Sex Behav* 2006; **35**: 145–61.
- 17 Lindau ST, Tang H, Gomero A, et al. Sexuality among middle-aged and older adults with diagnosed and undiagnosed diabetes: a national, population-based study. *Diabetes Care* 2010; **33**: 2202–10.
- 18 Spatz ES, Canavan ME, Desai MM, Krumholz HM, Lindau ST. Sexual activity and function among middle-aged and older men and women with hypertension. *J Hypertens* 2013; **31**: 1096–105.
- 19 Gott M, Hinchliff S, Galena E. General practitioner attitudes to discussing sexual health issues with older people. *Soc Sci Med* 2004; **58**: 2093–103.
- 20 Gott M, Galena E, Hinchliff S, Elford H. “Opening a can of worms”: GP and practice nurse barriers to talking about sexual health in primary care. *Fam Pract* 2004; **21**: 528–36.
- 21 Beckman N, Waern M, Gustafson D, Skoog I. Secular trends in self reported sexual activity and satisfaction in Swedish 70 year olds: cross sectional survey of four populations, 1971–2001. *BMJ* 2008; **337**: a279.
- 22 Erens B, Phelps A, Clifton S. Methodology of the third British National Survey of Sexual Attitudes and Lifestyles (Natsal-3). *Sex Transm Infect* 2013; published online Nov 26. DOI:10.1136/sextrans-2013-051359.
- 23 Erens B, Phelps A, Clifton S. The third National Survey of Sexual Attitudes and Lifestyles (Natsal-3): technical report. <http://www.natsal.ac.uk/natsal-3/methodology> (accessed Nov 26, 2013).
- 24 Jylhä M. What is self-rated health and why does it predict mortality? Towards a unified conceptual model. *Soc Sci Med* 2009; **69**: 307–16.
- 25 Young Y, Boyd CM, Guralnik JM, Fried LP. Does self-reported function correspond to objective measures of functional impairment? *J Am Med Dir Assoc* 2010; **11**: 645–53.
- 26 Arroll B. Screening for depression in primary care with two verbally asked questions: cross sectional study. *BMJ* 2003; **327**: 1144–46.
- 27 Arroll B, Goodyear-Smith F, Crengle S, et al. Validation of PHQ-2 and PHQ-9 to screen for major depression in the primary care population. *Ann Fam Med* 2010; **8**: 348–53.
- 28 Gray M, Nicholson S. National Survey of Sexual Attitudes and Lifestyles 2010: findings and recommendations from cognitive question testing. April, 2009. [http://www.surveynet.ac.uk/index/\\_search1099%5cCognitive%5cNatcennatsal2009.pdf#search=%22survey%22](http://www.surveynet.ac.uk/index/_search1099%5cCognitive%5cNatcennatsal2009.pdf#search=%22survey%22) (accessed Sept 13, 2013).
- 29 American Psychiatric Association. Diagnostic and statistical manual of mental disorders, fifth edition. Arlington, VA: American Psychiatric Publishing, 2013.
- 30 Mitchell KR, Mercer CH, Ploubidis GB, et al. Sexual function in Britain: findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3). *Lancet* 2013; published online Nov 25. [http://dx.doi.org/10.1016/S0140-6736\(13\)62366-1](http://dx.doi.org/10.1016/S0140-6736(13)62366-1).
- 31 Prah P, Copas AJ, Mercer CH, et al. Consistency in reporting sensitive sexual behaviours in Britain: change in reporting bias in the second and third National Surveys of Sexual Attitudes and Lifestyles (Natsal-2 and Natsal-3). *Sex Transm Infect* 2013; published online Nov 26. DOI:10.1136/sextrans-2013-051360.
- 32 Haapanen N, Miilunpalo S, Pasanen M, Oja P, Vuori I. Agreement between questionnaire data and medical records of chronic diseases in middle-aged and elderly Finnish men and women. *Am J Epidemiol* 1997; **145**: 762–69.
- 33 Okura Y, Urban LH, Mahoney DW, Jacobsen SJ, Rodeheffer RJ. Agreement between self-report questionnaires and medical record data was substantial for diabetes, hypertension, myocardial infarction and stroke but not for heart failure. *J Clin Epidemiol* 2004; **57**: 1096–103.
- 34 Health and Social Care Information Centre. Health Survey for England—2011, trend tables. Dec 20, 2012. <http://www.hscic.gov.uk/searchcatalogue?productid=10152&q=hypertension&topics=1%2fPublic+health%2fHealth+status&sort=Relevance&size=10&page=1#top> (accessed Oct 19, 2013).
- 35 Banks J, Marmot M, Oldfield Z, Smith JP. Disease and disadvantage in the United States and in England. *JAMA* 2006; **295**: 2037–45.
- 36 Lindau ST, Gavrilova N. Sex, health, and years of sexually active life gained due to good health: evidence from two US population based cross sectional surveys of ageing. *BMJ* 2010; **340**: c810.
- 37 DeLamater J, Moorman SM. Sexual behavior in later life. *J Aging Health* 2007; **19**: 921–45.
- 38 Bancroft J, Loftus J, Long JS. Distress about sex: a national survey of women in heterosexual relationships. *Arch Sex Behav* 2003; **32**: 193–208.
- 39 Gott M, Hinchliff S. Barriers to seeking treatment for sexual problems in primary care: a qualitative study with older people. *Fam Pract* 2003; **20**: 690–95.
- 40 Lindau ST, Abramssohn E, Gosch K, et al. Patterns and loss of sexual activity in the year following hospitalization for acute myocardial infarction (a United States national multisite observational study). *Am J Cardiol* 2012; **109**: 1439–44.
- 41 Stead ML, Fallowfield L, Brown JM, Selby P. Communication about sexual problems and sexual concerns in ovarian cancer: qualitative study. *BMJ* 2001; **323**: 836–37.
- 42 Hill EK, Sandbo S, Abramssohn E, et al. Assessing gynecologic and breast cancer survivors’ sexual health care needs (sexual care needs of cancer survivors). *Cancer* 2011; **117**: 2643–51.
- 43 Hinchliff S, Gott M. Seeking medical help for sexual concerns in mid- and later life: a review of the literature. *J Sex Res* 2011; **48**: 106–17.
- 44 Taylor A, Gosney MA. Sexuality in older age: essential considerations for healthcare professionals. *Age Ageing* 2011; **40**: 538–43.
- 45 Nusbaum MRH, Hamilton CD. The proactive sexual health history. *Am Fam Physician* 2002; **66**: 1705–12.
- 46 Gott M, Hinchliff S. How important is sex in later life? The views of older people. *Soc Sci Med* 2003; **56**: 1617–28.
- 47 Goodson P. Sexual activity in middle to later life. *BMJ* 2010; **340**: c850.
- 48 UK Department of Health. Raising the profile of long term conditions care: a compendium of information. January, 2008. [http://webarchive.nationalarchives.gov.uk/20080814090357/http://dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH\\_082069?IdcService=GET\\_FILE&dID=157746&Rendition=Web](http://webarchive.nationalarchives.gov.uk/20080814090357/http://dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_082069?IdcService=GET_FILE&dID=157746&Rendition=Web) (accessed May 17, 2013).
- 49 WHO. Defining sexual health: report of a technical consultation on sexual health, 28–31 January 2002, Geneva. 2006. [http://www.who.int/reproductivehealth/publications/sexual\\_health/defining\\_sexual\\_health.pdf](http://www.who.int/reproductivehealth/publications/sexual_health/defining_sexual_health.pdf) (accessed April 5, 2013).
- 50 Lusti-Narasimhan M, Beard JR. Sexual health in older women. *Bull World Health Organ* 2013; **91**: 707–09.
- 51 UK Department of Health. A framework for sexual health improvement in England. March, 2013. [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/142592/9287-2900714-TSO-SexualHealthPolicyNW\\_ACCESSIBLE.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/142592/9287-2900714-TSO-SexualHealthPolicyNW_ACCESSIBLE.pdf) (accessed June 27, 2013).