

LONDON  
SCHOOL of  
HYGIENE  
& TROPICAL  
MEDICINE



LSHTM Research Online

Harron, KL; Doidge, JC; Knight, HE; Gilbert, RE; Goldstein, H; Cromwell, DA; van der Meulen, JH; (2017) A guide to evaluating linkage quality for the analysis of linked data. International journal of epidemiology. ISSN 0300-5771 DOI: <https://doi.org/10.1093/ije/dyx177>

Downloaded from: <http://researchonline.lshtm.ac.uk/4539077/>

DOI: <https://doi.org/10.1093/ije/dyx177>

**Usage Guidelines:**

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact [researchonline@lshtm.ac.uk](mailto:researchonline@lshtm.ac.uk).

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

**A guide to evaluating linkage quality for the analysis of linked data**

Journal:	<i>International Journal of Epidemiology</i>
Manuscript ID	IJE-2017-03-0289.R1
Manuscript Type:	Education Corner
Date Submitted by the Author:	11-Jul-2017
Complete List of Authors:	Harron, Katie; London School of Hygiene and Tropical Medicine, Health Services Research and Policy Doidge, James; Administrative Data Research Centre for England Knight, Hannah; London School of Hygiene and Tropical Medicine, Health Services Research and Policy; Royal College of Obstetricians and Gynaecologists Gilbert, Ruth; University College London, Great Ormond Street Institute of Child Health Goldstein, H; University of Bristol, Graduate School of Education Cromwell, David; London School of Hygiene and Tropical Medicine van der Meulen, J; <none> ,
Key Words:	record linkage, bias, selection bias, linkage error, hospital records, data accuracy

17 February 2017

## A guide to evaluating linkage quality for the analysis of linked data

Katie Harron, James Doidge, Hannah Knight, Ruth Gilbert, Harvey Goldstein, David Cromwell, Jan van der Meulen

Corresponding author: Dr Katie Harron, 15-17 Tavistock Place, London, WC1 H 9SH, +44 207 927 2876; Katie.harron@lshtm.ac.uk

### Abstract

Linked datasets are an important resource for epidemiological and clinical studies, but linkage error can lead to biased results. For data security reasons, linkage of personal identifiers is often performed by a third party, making it difficult for researchers to assess the quality of the linked dataset in the context of specific research questions. This is compounded by a lack of guidance on how to determine the potential impact of linkage error. We describe how linkage quality can be evaluated and provide widely applicable guidance for both data providers and researchers.

Using an illustrative example of a linked dataset of maternal and baby hospital records, we demonstrate three approaches for evaluating linkage quality: applying the linkage algorithm to a subset of gold-standard data to quantify linkage error; comparing characteristics of linked and unlinked data to identify potential sources of bias; and evaluating the sensitivity of results to changes in the linkage procedure. These approaches can inform our understanding of the potential impact of linkage error and provide an opportunity to select the most appropriate linkage procedure for a specific analysis. Evaluating linkage quality in this way will improve the quality and transparency of epidemiological and clinical research using linked data.

**Key words:** record linkage; linkage error; bias; hospital records; data accuracy; sensitivity and specificity; selection bias; data linkage; administrative data

Word count: 4657

17 February 2017

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Key messages:

- Errors in data linkage are a potential source of bias in results of studies using linked data, yet researchers using linked data often find it difficult to assess the extent of such bias due to the separation of linkage and analysis processes.
- We describe three methods for evaluating data linkage quality and identifying potential sources of bias: applying the linkage algorithm to a subset of gold-standard data to quantify linkage error; comparing characteristics of linked and unlinked data to identify potential sources of bias; and evaluating the sensitivity of results to changes in the linkage procedure.
- These methods are relevant however and by whoever the linkage is conducted, and can provide a better understanding of the pattern of any bias, the extent to which linkage error may affect our results, and determinants of the amount of bias that is likely to be introduced.
- When linkage error is identified as a possible source of bias, methods to adjust for these biases should be used, which can help provide more robust results.

17 February 2017

## Introduction

Epidemiological and clinical research is increasingly based on datasets created by linking data from different sources such as administrative hospital datasets, clinical databases, and national death registers.[1, 2] Due to the nature of many of these data sources, which are typically collected for financial or clinical management, a unique person identifier (e.g. National Health Service number in the UK) may not be available for linkage. Therefore, linkage is often based on a series of identifiers that are not unique, are prone to errors or missing values, or are dynamic (i.e. may change over time such as postcode or name). Errors in the linkage process that arise from imperfect identifiers are increasingly being recognised as a potential source of bias in results from studies using linked data.[3, 4] However, assessing the extent to which linkage error affects results can be difficult for users of linked data who do not have access to the identifiable data used for linkage, which is often the case if the data linkage is performed by a third party. Separation of linkage and analysis processes in this way is recommended to preserve data security and personal confidentiality.[5]

Studies that evaluate linkage quality are therefore often restricted to estimates of the match rate (the proportion of records that were linked), sensitivity (the proportion of true links that were detected), or positive predictive value (the proportion of detected links that were true), which can be obtained, for example, by comparing a linked dataset with a 'gold-standard' or reference dataset, where true match status is known.[6] However, these metrics are limited in their ability to tell us the degree to which linkage error might produce bias in outcomes of interest. In some instances, we can assume that effect sizes will be underestimated but in most scenarios, it is not straightforward to predict the direction of bias that may result from linkage error. There are several factors that determine how linkage error affects an estimate but one key factor is the distribution of errors with respect to variables of interest, and this is usually unknown. In complex analyses incorporating multiple variables, different variables can be affected by linkage error in different ways.

17 February 2017

1  
2  
3 There is a lack of guidance on how to explore the extent to which error impacts upon analysis, and  
4  
5 this area has been identified as a priority for research.[7, 8] This Education Corner article describes  
6  
7 three simple approaches for evaluating quality of linkage, using an illustrative example of a linked  
8  
9 dataset of maternal and baby hospital records. We aim to provide guidance that is applicable to both  
10  
11 data providers and researchers, and to encourage application of these methods amongst  
12  
13 researchers using linked data.[9]  
14  
15

### 16 Why is linkage error important?

17 Linkage error can occur in two ways: false matches and missed matches. False matches occur when  
18  
19 records belonging to different individuals are erroneously linked together. False matches typically  
20  
21 (but not always) add noise to estimates, diluting the association between variables captured in  
22  
23 different datasets and biasing effect estimates towards zero.[3] Missed matches occur when records  
24  
25 belonging to the same individual are not linked. When unlinked records are excluded from analyses,  
26  
27 one consequence is reduced sample size and statistical power. If linkage is 'informative' (e.g. linkage  
28  
29 to a disease register indicating the presence of a particular condition), a consequence of missed  
30  
31 matches can be under-ascertainment of exposures or outcomes.[10, 11]  
32  
33  
34  
35

36 An important further issue is that linkage errors do not always occur randomly, meaning that  
37  
38 particular subgroups of individuals are often over- or under-represented amongst records affected  
39  
40 by linkage error. Systematic reviews of studies comparing the characteristics of linked and unlinked  
41  
42 records have identified that more vulnerable or hard to reach populations are often missed, with the  
43  
44 probability of a missed match being associated with a range of characteristics including gender, age,  
45  
46 ethnicity, deprivation and health status.[12, 13] Consequently, the linked data may not be  
47  
48 representative of the population of interest, which can reduce the study's external validity, or may  
49  
50 not capture subgroups that are of particular interest. As these demographic variables are often  
51  
52 associated with exposures or outcomes of interest, differential rates of linkage error may also  
53  
54 introduce bias. For example, unlinked mortality records in one particular ethnic group could lead to a  
55  
56 distorted comparison of mortality rates by ethnicity.[14-16]  
57  
58  
59  
60

17 February 2017

1  
2  
3 If unlinked records are to be excluded from analysis, selection bias (or collider-bias) can occur if  
4 selection into the linked dataset is related to both an exposure and outcome of interest.[17, 18] For  
5 example, suppose it is more difficult to link records for low birth weight babies and also more  
6 difficult to link records from mothers who smoke. In this case, records for low birth weight babies  
7 that are successfully linked are more likely to be from mothers who do not smoke (since, in this  
8 example, records from mothers who smoke are more difficult to link). Conditioning on linked records  
9 could therefore induce a protective relationship between maternal smoking and low birth weight,  
10 analogous to the birthweight paradox described in epidemiological literature.[19]  
11  
12  
13  
14  
15  
16  
17  
18  
19

## 20 Evaluating the impact of data linkage error

21  
22 The following sections describe three approaches to evaluating linkage quality (see Box 1 for a  
23 summary). The use of these methods can help researchers using linked data to understand the  
24 potential impact of linkage error on results.  
25  
26  
27  
28  
29

- 30 1. Using a gold-standard dataset to quantify false matches and missed matches;
- 31 2. Comparing characteristics of linked and unlinked data to identify potential sources of bias;
- 32 3. Sensitivity analyses to evaluate how sensitive results are to changes in linkage procedure.

33  
34 We use an illustrative example of linkage between hospital records for mothers and babies to  
35 demonstrate how these approaches can be implemented by researchers using linked data.  
36  
37

38  
39 *Box 1 here*  
40  
41  
42

### 43 Illustrative example: linking hospital records for mothers and babies

44  
45 The two most popular approaches to linkage have previously been described in an Education Corner  
46 article: deterministic (or rule-based) methods and probabilistic (or score-based) methods.[20]  
47  
48

49  
50 Alternative methods also exist.[21, 22] In a previous study, we used a combination of these  
51 techniques to create a mother-baby cohort of records from Hospital Episode Statistics (HES), an  
52 administrative data resource that holds detailed information of all admissions to National Health  
53 Service (NHS) hospitals in England.[23] The methods are described in full elsewhere, but comprised  
54  
55  
56  
57  
58  
59  
60

17 February 2017

1  
2  
3 deterministic and probabilistic linkage of de-identified information in data items contained in both  
4  
5 the mother's delivery record and the baby's birth record.[24] Using a deterministic algorithm based  
6  
7 on exact matching of hospital, maternal age, gestational age, baby's sex, birth order and GP practice  
8  
9 code, 42% of baby records were linked to a maternal record. The match rate increased to 98%  
10  
11 through the use of probabilistic linkage and additional variables (e.g. admission dates and mode of  
12  
13 delivery). However, the extent to which these missed matches (or any false matches) affect analyses,  
14  
15 has not yet been explored.

### 16 17 18 19 1. Evaluating linkage error using 'gold-standard' data

20  
21 If data are available where the true match status of each pair of records is known, these 'gold-  
22  
23 standard' data can be used to test linkage algorithms and estimate rates of linkage error. There are  
24  
25 various ways in which gold-standard datasets can be derived, for example from an additional data  
26  
27 source with complete identifiers, from a subsample of records that have been manually reviewed or  
28  
29 otherwise determined to be matches (or non-matches), or from a representative synthetic dataset  
30  
31 (e.g. generated through simulating data).[11, 25, 26] Gold-standard datasets allow us to identify  
32  
33 where errors have occurred in our linkage; where we have failed to linked records that should have  
34  
35 been linked (missed matches), or where we have linked together records belonging to different  
36  
37 entities (false matches). Since gold-standard data should be linked in the same way as the study data  
38  
39 (for comparison), involvement of data linkers is required. Unlike the second two approaches, gold-  
40  
41 standard comparisons cannot be readily implemented by researchers who do not have access to the  
42  
43 identifying data (e.g. in a 'trusted third party' model).  
44  
45  
46

#### 47 48 *Creating the gold-standard dataset*

49  
50 In order to create a gold-standard for assessing the quality of linkage within the HES mother-baby  
51  
52 cohort, we needed a dataset where the true match status of HES maternal and baby records was  
53  
54 known. This was possible due to another study, which collected electronic records from maternity  
55  
56 information systems (MIS) within 15 English obstetric units for births between April 2012 and March  
57  
58  
59  
60



17 February 2017

1  
2  
3 2013.[27] The MIS data captured NHS numbers for both mothers and babies together on the same  
4  
5 record (this is not the case for HES). The MIS records were linked by NHS Digital to corresponding  
6  
7 maternal and baby records in HES, using a deterministic approach based on NHS number, date of  
8  
9 birth, sex and postcode. After excluding a number of uncertain links (see Supplementary Figure 1),  
10  
11 the MIS-HES links provided a gold-standard dataset that could be used to validate the same subset  
12  
13 of births in the linked HES mother-baby cohort (Figure 1).  
14  
15

16  
17 *Figure 1 here*

#### 18 19 *Evaluating linkage using the gold-standard dataset*

20  
21 We compared linked records in the gold-standard dataset (linked using direct patient identifiers  
22  
23 including NHS number), with those in our HES mother-baby cohort (linked using linkage of indirect  
24  
25 identifiers captured for birth and delivery records). To enable this comparison, we applied our  
26  
27 original linkage algorithm to the same set of HES records captured in the MIS data. This allowed us  
28  
29 to quantify the number of false matches or missed matches that our linkage algorithm produced,  
30  
31 and to derive standard measures of linkage quality (sensitivity / recall, false match rate, and positive  
32  
33 predictive value / precision).[28] These metrics are thought to be more useful than specificity (due  
34  
35 to the imbalance between the number of non-matches versus matches in a linkage) or the F-  
36  
37 measure (due to issues when comparing linked datasets of different sizes).  
38  
39

40  
41 The gold-standard dataset comprised records for 72,817 babies (Supplementary Figure 1). Of these,  
42  
43 72,520 (99.6%) were linked using the original linkage algorithm and 297 (0.4%) were unlinked. Of the  
44  
45 72,520 linked records, 71,884 were true matches, giving a positive predictive value of 99.1%, a false  
46  
47 match rate of 0.9% (636 false matches/72,520 linked records), and a sensitivity of 98.7%  
48  
49 (71,884/72,817 true matches).  
50  
51

52  
53 The low error rates observed in this evaluation demonstrate that the original linkage algorithm was  
54  
55 highly accurate. We might therefore assume that the impact of linkage error will be negligible, since,  
56  
57 for example, excluding such a small proportion of the target population is unlikely to affect the  
58  
59  
60

17 February 2017

1  
2  
3 generalisability of results, or to dramatically reduce precision. However, further evaluation is needed  
4  
5 to assess whether selection bias could be present, e.g. if records from a particular subgroup were  
6  
7 more likely to be missed; or whether the 0.9% of false matches could have introduced enough noise  
8  
9 to bias results.  
10

## 11 2. Comparing characteristics of linked and unlinked data

12  
13  
14 In order to identify particular subgroups of records that are differentially missed during linkage, we  
15  
16 can compare the characteristics of linked and unlinked records. This method of quality appraisal can  
17  
18 only be implemented when all of the records in at least one of the files (or within the target sample  
19  
20 within one of the files) are expected to link; it would not be useful, for example, when linking to a  
21  
22 register of deaths to determine mortality, as there would be expected systematic differences  
23  
24 between those who link and those who do not. This approach can be implemented if researchers  
25  
26 have access to record-level or aggregate information on the characteristics of unlinked records.  
27  
28  
29 Since data linkage studies are often characterised by large sample sizes, standardised differences can  
30  
31 be more informative than  $p$ -values for comparing unlinked and linked records. Standardised  
32  
33 differences are calculated as the mean difference divided by the standard deviation and can be  
34  
35 easily calculated in statistical software packages (e.g. using the 'stddiff' command in Stata).[29, 30]  
36  
37 Standardised differences of 0.2, 0.5, and 0.8 represent small, moderate, and large standardized  
38  
39 differences, respectively.[29, 31] This helps us to identify variables that may have been more  
40  
41 affected by linkage error and are therefore potential sources of bias.[32]  
42  
43  
44  
45

### 46 *Evaluating linkage using comparisons of linked and unlinked records*

47  
48 Compared with true-matches in the gold-standard, the 297 records that failed to link (missed  
49  
50 matches) and 636 baby records that linked to the wrong mother (false matches) were more likely to  
51  
52 be multiple births, or babies with lower gestational age, lower birth weight, more neonatal medical  
53  
54 conditions, or babies born by caesarean section, or those of non-white ethnic background (Table 1).  
55  
56 Records with linkage errors were also more likely to be from babies born to nulliparous mothers, or  
57  
58  
59  
60

17 February 2017

1  
2  
3 mothers without pregnancy risk-factors. Linkage errors were strongly driven by data quality, since  
4 records with one or more missing values were less likely to link.  
5  
6

7  
8 *Table 1 here*  
9

10 The results in this example indicate that although linkage error rates were low, there was still some  
11 potential for bias, as particular subgroups of records were more often affected than others. Whether  
12 these differences were large enough to introduce bias into results depends on the relationship  
13 between these variables and the parameters of interest. It is therefore helpful to explore how  
14 results of interest might change according to different levels of error.  
15  
16  
17  
18  
19  
20  
21

### 22 3. Sensitivity analyses 23

24 In order to assess the sensitivity of results to different linkage procedures, we can perform  
25 sensitivity analyses, aiming to assess the extent to which results vary and the direction of likely bias.  
26 This can involve changing the linkage algorithm or varying the match weight threshold for  
27 probabilistic linkage, and re-running analyses to evaluate any impact on results.[3, 14] The aim of  
28 this approach is to determine whether decisions about the design of the linkage procedure could  
29 have had a substantial impact on inferences drawn from the linked data. These types of sensitivity  
30 analysis can be implemented by researchers without access to identifying data, if they are provided  
31 with match weights (in probabilistic linkage) or decision-steps (in deterministic linkage). As these are  
32 not sensitive data, data providers and linkers are usually able to share these with researchers.[9]  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

#### 44 *Evaluating linkage using sensitivity analyses* 45

46 We conducted a sensitivity analysis to evaluate the mother-baby linkage by changing the threshold  
47 used in our linkage algorithm, and comparing results across different sets of linked records. We  
48 compared linkage results from our original probabilistic algorithm using a threshold weight of 20 for  
49 classifying records as links, with results from an algorithm that minimised false links by using a  
50 considerably higher threshold of 45. These thresholds were selected based on examination of the  
51 observed distribution of weights in our analysis; this distribution can differ substantially depending  
52  
53  
54  
55  
56  
57  
58  
59  
60

17 February 2017

1  
2  
3 on the number and quality of matching variables so thresholds are generally selected in the context  
4  
5 of a specific linkage or analysis. The aim of this type of sensitivity analysis is to select thresholds that  
6  
7 are likely to reflect plausible limits for the trade-off between false matches and missed matches. We  
8  
9 also compared results with those from the initial deterministic linkage only, i.e. where records  
10  
11 agreed exactly on hospital, maternal age, gestational age, baby's sex, birth order and GP practice  
12  
13 code.  
14

15  
16 As expected, increasing the match weight threshold in probabilistic linkage, or using deterministic  
17  
18 linkage only, produced linkages that introduced fewer false matches but more missed matches  
19  
20 (Table 2). This is because stricter linkage criteria make it less likely that records belonging to  
21  
22 different entities will link by chance, but more likely that records with missing or incorrect linking  
23  
24 variables will remain unlinked.  
25  
26

27  
28 *Table 2 here*  
29

30  
31 We expected that impact of linkage errors in each of these linkage scenarios would depend on the  
32  
33 research question, and therefore assessed four different outcomes: proportion of stillbirths,  
34  
35 proportion of preterm births (<37 weeks of gestation), the association between neonatal survival to  
36  
37 discharge and delivery risk factors, and the association between delivery risk factors and ethnic  
38  
39 group. Odds ratios were estimated from logistic regression models, adjusting for a number of  
40  
41 maternal and neonatal risk-factors (listed in Table 1, based on ICD-10 diagnosis codes listed in  
42  
43 Supplementary Table 1). Analysis was performed in Stata 14.[33]  
44

45  
46  
47 i. Proportions of stillbirths and preterm births  
48

49 We expected that the generalisability (i.e. external validity) of the data would be affected by missed  
50  
51 matches. In particular, we expected that records of preterm births or stillbirths would be less likely  
52  
53 to link than those of later gestations or live births, and that the ascertainment of these outcomes  
54  
55 would therefore be lower in datasets more affected by missed matches.  
56  
57  
58  
59  
60

17 February 2017

1  
2  
3 By comparing results across different linkage algorithms (Table 3), we observed that for preterm  
4 birth (7.65% of records in gold-standard), ascertainment was similar for the original linkage  
5 algorithm (7.64%), but ascertainment was a little lower when using a higher match weight threshold  
6 of 45 (7.31%) or deterministic linkage only (7.43%). All confidence intervals for proportions of  
7 stillbirths and preterm births estimated in the linked datasets overlapped with those in the gold-  
8 standard, indicating that in this example, linkage error is unlikely to have resulted in substantial bias  
9 for these outcomes.  
10  
11  
12  
13  
14  
15  
16

17  
18  
19 *Table 3 here*

20  
21 ii. Association between neonatal survival to discharge and recording of delivery risk-factors  
22

23  
24 We expected that statistical power would be affected either through missed matches (due to a  
25 reduction in the size of the study population), or a lack of precision introduced by false matches  
26 (leading to increased noise in the association between variables). Given the large sample size, we  
27 assumed that power implications would be most important for identifying associations with rare  
28 outcomes (e.g. mortality).  
29  
30  
31  
32  
33  
34  
35

36 As expected, we found that as the number of linked records decreased due to more missed matches  
37 at higher thresholds (Figure 2), there was a reduction in the precision of estimates of association  
38 between delivery risk-factors and survival (Table 3). Although an association between delivery risk-  
39 factors and survival was observed in the gold-standard data (adjusted odds ratio = 0.40; 95% CI 0.17-  
40 0.95), this association was no longer observed in the deterministic linkage where less than half of  
41 records were linked (adjusted odds ratio = 0.52; 95% CI 0.22-1.25).  
42  
43  
44  
45  
46  
47  
48

49 *Figure 2 here*

50  
51  
52 These results demonstrate that there can be power implications when large numbers of unlinked  
53 records are excluded from analysis; a problem analogous to that of complete case analysis in the  
54 presence of missing data.  
55  
56  
57  
58  
59  
60

17 February 2017

iii. Association between delivery risk-factors and ethnicity

We expected that selection bias could be introduced if selection into the linked dataset were associated with both the outcome and exposure. For example, earlier comparisons between linked and unlinked data had indicated that that records for mothers with delivery risk-factors were less likely to link, and also that mothers in the Black ethnic group were less likely to link. If this was the case, mothers with delivery risk-factors who were successfully linked would be more likely to be from other Ethnic groups (since, in this example, those from the Black ethnic group were harder to link). Conditioning on linked records could therefore induce a spurious protective relationship between Black ethnicity and delivery risk-factors.

In the gold-standard data, we observed that 6.5% of mothers with delivery risk-factors were from the Black ethnic group, whilst in the deterministic linkage, only 4.7% of mothers with delivery risk-factors were from the Black ethnic group. There was no true association between ethnicity and delivery risk-factors. However, within the deterministically-linked data, there appeared to be a protective effect (odds ratio = 0.80; 95% CI 0.66-0.96, Table 3). These results indicate that in this example, some inferences from linked data could be incorrect due to selection bias in the presence of missed matches.

## Discussion

We describe three approaches for evaluating linkage quality, and demonstrate how these methods can be used determine the extent to which linkage error may introduce bias for a specific research question. Our illustrative example showed that even with high linkage rates, particular subgroups of individuals are disproportionately affected by linkage error, as has been observed in previous literature.[13] However, we demonstrate that sensitivity analyses can help us to understand the direction of any bias, and to assess whether linkage errors may influence inferences from the linked dataset.

17 February 2017

### Access to information required to evaluate linkage quality

In many jurisdictions, linkage is carried out by an independent body, and information about the linkage processes is not readily available to researchers. Comprehensive guidelines on information that should be shared between data providers, linkers and researchers are available elsewhere.[9] All of the approaches described in this article require some level of collaboration between data linkers and the researchers aiming to evaluate linkage quality but only one (gold-standard data) requires direct involvement of the linkers; approaches 2 and 3 can be implemented by researchers, provided that certain non-sensitive information is shared.

Our example of HES mother-baby linkage was supported by the availability of a subset of gold-standard data, which is one of the most useful tools for quantifying linkage error. Consideration should always be given to the representativeness of the gold-standard dataset. In our example, the proportion of unlinked records in the gold-standard data was lower than in the HES mother-baby cohort overall, indicating that data quality in the hospitals contributing MIS data may have been slightly better than average. The same applies to gold-standard datasets created using subsets of records that have complete data either on a single unique identifier, or on a sufficient set of partially identifying variables; records with high quality data may differ systematically from those of poorer quality data.

In practice, gold-standard datasets are rarely available. Even those that are generated (e.g. through manually reviewing a sample of linked and unlinked records, or by creating a synthetic dataset with the same characteristics as the original data), are often only available to the data linkers and not to researchers.[34] In practice, this means that data linkers would need to evaluate linkage quality using a gold-standard (as researchers generally would not have access to gold-standard data).

However, when gold-standard datasets are not available, researchers can consider alternative approaches: comparisons of characteristics of linked and unlinked data, and sensitivity analyses. These methods can be easily implemented, but require data linkers to provide information on the characteristics of unlinked records, and/or the quality of each potential link.[9] Providing record-

17 February 2017

1  
2  
3 level or aggregate characteristics of unlinked records allows researchers to compare linked and  
4  
5 unlinked records, to identify any potential sources of bias, where particular subgroups of records  
6  
7 were missed from the linkage.  
8  
9

10 Sensitivity analyses can be performed if measures of linkage certainty (e.g. match weights in  
11  
12 probabilistic matching or matching ranks/criteria in hierarchical deterministic matching) are  
13  
14 provided by data linkers alongside a linked dataset.[9] This makes any uncertainty or subjectivity in  
15  
16 the linkage process more transparent and allows researchers to run analysis on different sets of  
17  
18 linked records in turn. It should be noted that caution is needed when interpreting results of this  
19  
20 type of sensitivity analysis, as in most cases, none of the linkage algorithms compared will be 100%  
21  
22 accurate. The trade-off between false matches and missed matches will vary depending on the  
23  
24 linkage algorithm, and these errors may impact on results in opposing or compounding ways. In our  
25  
26 example, inferences about the associations between delivery risk-factors and survival to discharge,  
27  
28 and between ethnicity and delivery risk-factors, differed between the methods. Exploring  
29  
30 differences in results over a range of linkage algorithms in this way can help researchers consider the  
31  
32 pattern of any bias, and to identify scenarios which are particularly likely to produce substantial bias.  
33  
34  
35  
36

### 37 Further methods for evaluating linkage quality

38 In our example, we expected all babies to link with a mother, which made comparisons between  
39  
40 linked and unlinked records easily interpretable, and allowed us to directly estimate the proportion  
41  
42 of missed matches. However, careful consideration needs to be given to appropriate reference  
43  
44 populations when all records are not expected to link. For example, we would not expect all hospital  
45  
46 records to link with a mortality record and vice versa; rather a successful link indicates that an  
47  
48 individual has died ('informative linkage'). If that is the case, comparing the characteristics of the  
49  
50 individuals whose records were and were not linked would also be affected by differences in the  
51  
52 groups for whom no linked records were available (i.e. the difference between those who died and  
53  
54 those who survived). In such situations, external reference data (e.g. age-specific mortality rates) can  
55  
56 allow us to assess how linkage rates might differ for different subgroups.[10]  
57  
58  
59  
60



17 February 2017

1  
2  
3 Further methods not covered in this article can also be used to evaluate linkage quality in the  
4  
5 absence of a gold-standard. For example, estimates of false match rates can be derived by applying  
6  
7 linkage algorithms to records known to have no match (e.g. attempting to link with mortality records  
8  
9 for individuals known to be alive, or attempting to link male patient records with maternity  
10  
11 records).[3] Alternatively, inconsistency checks, such as checks for admissions following a patient's  
12  
13 death, linkage between a male patient and a caesarean section, or linkage of one mortality record to  
14  
15 two different individuals, can be performed post-linkage in de-identified data to identify false  
16  
17 matches.[35, 36] Linkage error rates estimated in this way should be interpreted with caution, as not  
18  
19 all errors may have been identified, distinguishing between linkage errors and data coding errors can  
20  
21 be difficult.[35] Nevertheless, these methods can reveal useful information about the *relative*  
22  
23 distribution of errors across subgroups or with respect to variables of interest (i.e. while the absolute  
24  
25 error rate may remain unknown, higher rates of inconsistencies may be observed with respect to  
26  
27 some variable of interest, implicating likely bias).[35]

### 31 Handling bias in the analysis of linked data

32 Evaluation of linkage quality can guide decisions about appropriate study design. For example, if  
33  
34 linkage is used to identify individuals with a particular condition or disease (informative linkage),  
35  
36 high levels of missed matches will lead to under ascertainment, meaning that cohort study designs  
37  
38 may be unsuitable (particularly for deriving estimates of prevalence or incidence). Where linkage  
39  
40 rates are too low, researchers may conclude that linked data are not fit for these purposes. On the  
41  
42 other hand, a case-control study may still be valid, whereby a high threshold is used to identify cases  
43  
44 and a low threshold is used to identify controls (assuming no other biases are present).[37] In this  
45  
46 scenario, records for which there is uncertainty about linkage would not be included in analysis.

47  
48 An alternative, which still makes use of all available records, is to use multiple imputation to handle  
49  
50 missing values due to unlinked or equivocal records.[38] Furthermore, information from match  
51  
52 weights can be incorporated into imputation procedures, making use of variable distributions in  
53  
54 candidate links (known as 'prior-informed imputation').[39, 40] This method incorporates  
55  
56  
57  
58  
59  
60

17 February 2017

1  
2  
3 information from 'auxiliary' variables, such as individual characteristics associated with linkage  
4  
5 quality (e.g. birthweight or ethnicity), to help correct for selection biases.  
6  
7

8 In situations in which we have information about how linkage error affects the distribution of  
9  
10 outcomes and exposures in our data, it may be possible to use well-established techniques for  
11  
12 quantitative bias analysis, to adjust for these errors.[41, 42] This is particularly relevant for simple  
13  
14 analyses, but becomes more complex with complicated designs involving more than two data  
15  
16 sources and/or a number of covariates. Developing appropriate methods to handle bias arising from  
17  
18 linkage error is a priority for methodological research.[43]  
19

20  
21 Studies of linked data are often based on administrative data that has not been collected primarily  
22  
23 for research. In addition to linkage error, researchers should also consider other issues specifically  
24  
25 relevant to these types of data (e.g. missing data, coding changes, service changes, etc.), and explore  
26  
27 methods to handle any potential bias that is identified.[43, 44]  
28  
29

### 30 31 Summary

32  
33 We describe three methods for evaluating linkage quality: applying the linkage algorithm to a subset  
34  
35 of gold-standard data to quantify linkage error; comparing characteristics of linked and unlinked  
36  
37 data to identify potential sources of bias; and evaluating the sensitivity of results to changes in the  
38  
39 linkage procedure. These methods are generalisable to many other linkage situations and can be  
40  
41 used as a guide for evaluating the quality of linkage for population-based analyses of linked data.  
42  
43

44 Researchers using linked data should collaborate with data providers to understand the data linkage  
45  
46 process, including data extraction and cleaning, linkage methods, and resulting data quality.[9]  
47

48 Ultimately, this will improve transparency and enhance the value of linked data for epidemiological  
49  
50 and clinical research.[44]  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

17 February 2017

## Figure legends

### Figure 1: Creation of a gold-standard dataset for evaluating linkage quality in the HES mother-baby cohort.

<sup>1</sup> 83,195 records of births/deliveries from 15 English hospitals, April 2012-March 2013; <sup>2</sup> 672,955 records of births/deliveries from all NHS hospitals in England, April 2012-March 2013; <sup>3</sup> 72,817 records in the HES mother-baby validation cohort (gold-standard).

### Figure 2: Number of linked records and percentage of missed matches and false matches for a range of linkage criteria. W=threshold used to classify links in probabilistic linkage.

## Acknowledgements

We would like to thank Lynn Copley for managing the required data. We would also like to thank the NHS Trusts that participated in the RCOG MIS pilot project and have agreed to be named in resulting academic outputs: Birmingham Women's NHS Foundation Trust, Bradford Teaching Hospitals NHS Foundation Trust, Cambridge University Hospitals NHS Foundation Trust, Central Manchester University Hospitals NHS Foundation Trust, Chelsea and Westminster Hospital NHS Foundation Trust, Homerton University Hospital NHS Foundation Trust, Liverpool Women's NHS Foundation Trust, Mid Cheshire Hospitals NHS Foundation Trust, Norfolk & Norwich University Hospitals NHS Foundation Trust, Nottingham University Hospitals NHS Trust\*, Southern Health and Social Care Trust, St George's Healthcare NHS Trust, The Newcastle Upon Tyne Hospitals NHS Foundation Trust, University College London Hospitals NHS Foundation Trust, and Western Sussex Hospitals NHS Foundation Trust\*.

\*These trusts contributed data for more than one hospital.

Approvals for the use of HES data were obtained as part of the standard Hospitals Episode Statistics approval process. Hospital Episode Statistics were made available by NHS Digital. All rights reserved.

17 February 2017

## Funding

This work was supported by the Wellcome Trust [grant number 103975/Z/14/Z]. Jan van der Meulen is supported by the NIHR CLAHRC North Thames. The work was also supported by the Economic and Social Research Council through the Administrative Data Research Centre for England (ES/L007617/1).

## Conflict of interest

The authors state no relevant conflicts of interest.

For Review Only

17 February 2017

## References

1. Jutte, DP, L Roos, and MD Brownell, *Administrative record linkage as a tool for public health research*. *Annu Rev Public Health*, 2011. **32**: 91-108.
2. Holman, C, et al., *A decade of data linkage in Western Australia: strategic design, applications and benefits of the WA data linkage system*. *Aust Health Rev*, 2008. **32**(4): 766-777.
3. Moore, CL, et al., *A new method for assessing how sensitivity and specificity of linkage studies affects estimation*. *PLoS One*, 2014. **9**(7): e103690.
4. Baldi, I, et al., *The impact of record linkage bias in the Cox model*. *J Eval Clin Pract*, 2010. **16**(1): 92-96.
5. Kelman, C, A Bass, and C Holman, *Research use of linked health data - a best practice protocol*. *Aust N Z J Public Health*, 2002. **26**(3): 251 - 255.
6. Pinto da Silveira, D and E Artmann, *Accuracy of probabilistic record linkage applied to health databases: systematic review*. *Rev Saúde Públ*, 2009. **43**(5): 875-82.
7. Jorm, L, *Routinely collected data as a strategic resource for research: priorities for methods and workforce*. *Public Health Res Pract*, 2015. **25**(4): e2541540.
8. Boyd, JH, et al., *Understanding the origins of record linkage errors and how they affect research outcomes*. *Aust N Z J Public Health*, 2016. **41**(2): 215.
9. Gilbert, R, et al., *GUILD: Guidance for Information about Linking Datasets*. *J Public Health*, 2017. **1-8**.
10. Schmidlin K, et al., *Impact of unlinked deaths and coding changes on mortality trends in the Swiss National Cohort*. *BMC Med Inform Decis Mak*, 2013. **13**(1): 1-11.
11. Harron, K, et al., *Linkage, evaluation and analysis of national electronic healthcare data: application to providing enhanced blood-stream infection surveillance in paediatric intensive care*. *PLoS One*, 2013. **8**(12): e85278.
12. Bohensky, M, *Chapter 4: Bias in data linkage studies*, in *Methodological Developments in Data Linkage*, K. Harron, C. Dibben, and H. Goldstein, Editors. 2015, Wiley: London.
13. Bohensky, M, et al., *Data linkage: A powerful research tool with potential problems*. *BMC Health Serv Res*, 2010. **10**(1): 346-352.
14. Lariscy, JT, *Differential record linkage by hispanic ethnicity and age in linked mortality studies*. *J Aging Health*, 2011. **23**(8): 1263-1284.
15. O'Reilly, D, M Rosato, and S Connolly, *Unlinked vital events in census-based longitudinal studies can bias subsequent analysis*. *J Clin Epidemiol*, 2008. **61**(4): 380-385.
16. Zingmond, D, et al., *Linking hospital discharge and death records--accuracy and sources of bias*. *J Clin Microbiol*, 2004. **57**(1): 21-29.
17. Cole, SR, et al., *Illustrating bias due to conditioning on a collider*. *Int J Epidemiol*, 2010. **39**(2): 417-420.
18. Hernán, MA, S Hernández-Díaz, and JM Robins, *A structural approach to selection bias*. *Epidemiology*, 2004. **15**(5): 615-625.
19. VanderWeele, TJ, *Commentary: Resolutions of the birthweight paradox: competing explanations and analytical insights*. *Int J Epidemiol*, 2014. **43**(5): 1368-1373.
20. Sayers, A, et al., *Probabilistic record linkage*. *Int J Epidemiol*, 2015. **45**(3): 954-964.
21. Goldstein, H, K Harron, and M Cortina-Borja, *A scaling approach to record linkage*. *Stat Med*, 2017. **36**: 2514-21.
22. Abbott, O, P Jones, and M Ralphs, *Chapter 8: Large-scale linkage for total populations in official statistics*, in *Methodological Developments in Data Linkage*, K. Harron, C. Dibben, and H. Goldstein, Editors. 2015, Wiley: London.
23. Herbert, A, et al., *Data Resource Profile: Hospital Episode Statistics Admitted Patient Care (HES APC)*. *Int J Epidemiol*, 2017. **Epub**: doi: 10.1093/ije/dyx015.

17 February 2017

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
24. Harron, K, et al., *Linking data for mothers and babies in de-identified electronic health data*. PLoS One, 2016. **11**(10): e0164667.
  25. Winglee, M, R Valliant, and F Scheuren, *A case study in record linkage*. Surv Methodol, 2005. **31**(1): 3-11.
  26. Aldridge, RW, et al., *Accuracy of Probabilistic Linkage Using the Enhanced Matching System for Public Health and Epidemiological Studies*. PLoS ONE, 2015. **10**(8): e0136179.
  27. Knight, HE, et al., *Birth "Out-of-Hours": An Evaluation of Obstetric Practice and Outcome According to the Presence of Senior Obstetricians on the Labour Ward*. PLoS Med, 2016. **13**(4): e1002000.
  28. Ferrante, A and J Boyd, *A transparent and transportable methodology for evaluating Data Linkage software*. J Biomed Inform, 2012. **45**(1): 165 - 172.
  29. Austin, PC, *Balance diagnostics for comparing the distribution of baseline covariates between treatment groups in propensity-score matched samples*. Stat Med, 2009. **28**(25): 3083-107.
  30. Bayoumi, AM, *STDDIFF: Stata module to compute Standardized differences for continuous and categorical variables*. 2016: Boston College Department of Economics.
  31. Cohen, J, *Statistical power analysis for the behavioral sciences Lawrence Earlbaum Associates*. Hillsdale, NJ, 1988: 20-26.
  32. Ford JB, Roberts CL, and Taylor LK, *Characteristics of unmatched maternal and baby records in linked birth records and hospital discharge data*. Paediatr Perinat Epidemiol, 2006. **20**(4): 329-337.
  33. Stata, *Stata Statistical Software: Release 14*. 2015: College Station, TX: StataCorp LP
  34. Boyd, JH, et al., *A Simple Sampling Method for Estimating the Accuracy of Large Scale Record Linkage Projects*. Methods of Information in Medicine, 2016. **55**(3): 276-283.
  35. Hagger-Johnson, G, et al., *Identifying false matches in anonymised hospital administrative data without patient identifiers* Health Serv Res, 2014. **50**(4): 1162-78.
  36. Blakely, T and C Salmond, *Probabilistic record linkage and a method to calculate the positive predictive value*. Int J Epidemiol, 2002. **31**(6): 1246-52.
  37. Paixao E, et al., *Quality of record linkage of large administrative databases in a middle income country: still births and dengue during pregnancy in Brazil*. BMC Med Inform Decis Mak, 2017. **in press**.
  38. Harron, K, et al., *Evaluating bias due to data linkage error in electronic healthcare records*. BMC Med Res Methodol, 2014. **14**(1): 36.
  39. Goldstein, H and K Harron, *Chapter 6: Record linkage: a missing data problem*, in *Methodological Developments in Data Linkage*, K. Harron, C. Dibben, and H. Goldstein, Editors. 2015, Wiley: London.
  40. Goldstein, H, K Harron, and A Wade, *The analysis of record-linked data using multiple imputation with data value priors*. Stat Med, 2012. **31**(28): 3481-93.
  41. Schneeweiss, S, *Sensitivity analysis and external adjustment for unmeasured confounders in epidemiologic database studies of therapeutics*. Pharmacoepidemiol drug saf, 2006. **15**(5): 291-303.
  42. Lash, TL, MP Fox, and AK Fink, *Applying quantitative bias analysis to epidemiologic data*. 2011: Springer Science & Business Media.
  43. Harron, K, C Dibben, and H Goldstein, *Methodological developments in data linkage*. 2015: Wiley.
  44. Benchimol, EI, et al., *The REporting of studies Conducted using Observational Routinely-collected health Data (RECORD) Statement*. PLoS Med, 2015. **12**(10): e1001885.

Box 1: Summary of approaches to evaluating linkage quality

	<b>Using a gold-standard dataset to quantify false matches and missed matches</b>	<b>Comparing characteristics of linked and unlinked data to identify potential sources of bias</b>	<b>Sensitivity analyses to evaluate how sensitive results are to changes in linkage procedure</b>
<i>Purpose</i>	To quantify errors occurred (missed matches and false matches)	To identify subgroups of records that are more prone to linkage error and are potential sources of bias	Assesses the extent to which results of interest may vary depending on different levels of error, and the direction of likely bias
<i>Strengths</i>	Easily interpretable; allows linkage error to be fully measured	Straightforward to implement and easily interpretable	Straightforward to implement
<i>Limitations</i>	Representative gold-standard data are rarely readily available	Cannot be applied if systematic differences are expected between linked unlinked records (e.g. if linking to death register)	Results may be difficult to interpret as false matches and missed matches may impact on results in opposing or compounding ways
<i>Technical requirements</i>	A representative group of people for which links are known; data linker capacity to perform evaluation (researchers rarely have access to gold-standard data)	A linkage design where all records in at least one file are expected to link; provision of record-level or aggregate characteristics of unlinked records	Provision of information on the strength of the match (e.g. deterministic rule or probabilistic match weight)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1: Characteristics of records in the HES mother-baby cohort according to linkage status derived from gold-standard data.** St. diff = standardised differences; 0.2, 0.5, and 0.8 can be considered as small, medium, and large effect sizes respectively.

	True matches (N=71,884)		False matches (N=636)		St. diff	Missed matches (N=297)		St. diff	
	N	%	N	%		N	%		
Still birth	325	0.5	6	0.9	0.1	9	3.0	0.2	
Survival to postnatal discharge	71,384	99.1	627	98.1	0.1	286	96.3	0.2	
Delivery risk factor <sup>1</sup>	6738	9.4	105	16.4	0.2	49	16.5	0.2	
Female infant	34,967	48.7	321	50.2	0.0	140	47.1	0.0	
Multiple birth	1961	2.7	126	19.7	<b>0.6</b>	31	10.4	0.3	
Caesarean section	18,034	25.1	63	9.9	0.4	10	3.4	<b>0.7</b>	
Pregnancy risk factor <sup>2</sup>	7388	10.3	16	2.5	0.3	1	0.3	0.5	
Neonatal medical condition <sup>3</sup>	6281	8.7	91	14.2	0.2	90	30.3	<b>0.6</b>	
Neonatal ICU	8461	11.8	32	5.0	0.2	33	11.1	0.0	
Parity: nulliparous	27,125	37.7	335	52.4	0.3	192	64.5	<b>0.6</b>	
Gestational age group	Full term (39+ wks)	45,611	72.3	102	44.4	<b>0.7</b>	27	44.3	<b>0.7</b>
	Early term (37-38 wks)	12,721	20.2	66	28.7		17	27.9	
	Late preterm (34-36 wks)	3280	5.2	39	17.0		6	9.8	
	Moderate / very preterm (<34 wks)	1494	2.4	23	10.0		11	18.0	
	Missing*	8775	12.2	409	64.0		236	79.5	
Birth weight	<1500	909	1.4	14	6.1	<b>0.7</b>	7	10.9	<b>0.7</b>
	1500-<2500	1798	6.0	45	19.7		12	18.8	
	2500-<4000	51,718	82.0	160	69.9		42	65.6	
	4000+	6687	10.6	10	4.4		3	4.7	
	Missing*	8769	12.2	410	64.2		233	78.5	
Size for gestation	Small (<10 <sup>th</sup> percentile)	5274	8.4	25	11.1	0.2	5	8.3	0.1
	Normal	54,367	81.6	187	93.1		51	85.0	
	Large (>10 <sup>th</sup> percentile)	6344	10.1	13	5.8		4	6.7	
	Missing*	8896	12.4	414	64.8		237	79.8	
Ethnicity	White	48,896	68.0	408	63.9	0.3	165	55.6	0.4
	Mixed	3410	4.7	24	3.8		14	4.7	
	Asian	7367	10.3	49	7.7		20	6.7	
	Black	4866	6.8	32	5.0		25	8.4	
	Other	4508	6.3	77	12.1		38	12.8	
	Unknown	2834	3.9	49	7.7		35	11.8	
Newborn length of stay (days)	<2	38,329	53.3	315	49.3	0.2	131	44.1	<b>0.7</b>
	2-6	28,946	40.3	244	38.2		74	24.9	
	7+	4599	6.4	80	12.5		92	31.0	
Maternal age (years)	<20	2859	4.0	21	3.3	0.1	13	4.4	0.2
	20-24	11,752	16.4	88	13.8		42	14.1	
	25-29	19,226	26.8	155	24.3		55	18.5	
	30-34	22,377	31.1	220	34.4		101	34.0	
	35-39	12,433	17.3	125	19.6		64	21.6	
	40+	3234	4.5	30	4.7		22	7.4	
Income/deprivation quintile <sup>4</sup>	Most deprived	27,042	37.7	206	32.3	0.1	97	32.9	0.2
	2	16,394	22.9	170	26.7		86	29.2	
	3	13,104	18.3	129	20.3		58	19.7	
	4	9,040	12.6	77	12.1		37	12.5	
	Most affluent	6,146	8.6	55	8.6		17	5.8	
	Missing*	155	0.2	2	0.3		2	0.7	

<sup>1</sup> Hypoxia, amniotic fluid embolism, placental-transfusion syndrome, umbilical cord prolapse, chorioamnionitis, fetal haemorrhage, birth trauma, complications of delivery, umbilical cord problem

<sup>2</sup> Eclampsia, gestational hypertension, diabetes, placental abruption or infarction

<sup>3</sup> Congenital anomaly, perinatal infection, neonatal abstinence syndrome, respiratory distress syndrome

<sup>4</sup> Quintiles of deprivation were derived from the Index of Multiple Deprivation (IMD) score based on patient postcode in HES.

\* percentage of records with missing data (excluded from other category percentages)



Table 2: Linkage success for a range of linkage criteria

	Original probabilistic linkage (threshold weight = 20)	High threshold probabilistic linkage (threshold weight = 45)	Deterministic linkage only
Linked records	72,520/72,817 99.6%	65,020/72,817 89.3%	35,324/72,817 48.5%
Missed match rate	297/72,817 0.4%	7797/72,817 10.7%	37,493/72,817 51.5%
False match rate	636/72,520 0.9%	212/65,020 0.3%	22/35,324 0.1%
Positive predictive value	71,884/72,520 99.1%	64,808/65,020 99.7%	35,302/35,324 99.9%

Table 3: Comparison of outcome measures for a range of linkage criteria

	Gold-standard	Original probabilistic linkage	High threshold probabilistic linkage	Deterministic linkage only
% Preterm births (95% CI)	7.65 (7.45-7.86)	7.64 (7.43-7.85)	7.31 (7.11-7.53)	7.43 (7.16-7.71)
% Stillbirths (95% CI)	0.47 (0.42-0.52)	0.46 (0.41-0.51)	0.44 (0.39-0.49)	0.45 (0.40-0.50)
Odds ratio (95% CI) for neonatal survival to discharge: mothers with delivery risk-factors vs. those without p=0.039	0.40 (0.17-0.95) p=0.039	0.42 (0.18-0.98) p=0.044	0.35 (0.15-0.79) p=0.011	0.52 (0.22-1.25) p=0.143
Odds ratio (95% CI) for delivery risk-factors: Black ethnicity vs. White ethnicity p=0.700	0.98 (0.88-1.09) p=0.700	0.97 (0.87-1.08) p=0.593	0.89 (0.79-1.01) p=0.067	0.80 (0.66-0.96) p=0.017




Figure 1: Creation of a gold-standard dataset for evaluating linkage quality in the HES mother-baby cohort.

<sup>1</sup> 83,195 records of births/deliveries from 15 English hospitals, April 2012-March 2013; <sup>2</sup> 672,955 records of births/deliveries from all NHS hospitals in England, April 2012-March 2013; <sup>3</sup> 72,817 records in the HES mother-baby validation cohort (gold-standard).

190x124mm (300 x 300 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 2: Number of linked records and percentage of missed matches and false matches for a range of linkage criteria. W=threshold used to classify links in probabilistic linkage.

190x130mm (300 x 300 DPI)

W Only