

Table 2. Age-standardized prevalence estimates in subjects aged 55 years or more, stratified by sex, time period and geographical area (prevalence (%) and 95 % confidence interval)

	Non-refractive visual impairment (WHO) (BCVA<20/60)	Non-refractive visual impairment (US) (BCVA<20/40)	Non-refractive blindness (BCVA<20/400)	
Sex				
Men	1.38 (0.72-2.03)	3.17 (1.98-4.36)	0.32	(0.12-0.52)
Women	1.81 (0.96-2.66)	4.24 (2.65-5.83)	0.39	(0.17-0.62)
Period				
1991-2006	2.22 (1.34-3.10)	4.68 (2.68-6.68)	0.53	(0.24-0.81)
2007-2012	0.92 (0.37-1.42)	2.86 (1.52-4.20)	0.13	(0.01-0.26)
Geographical area				
Northern countries ¹	1.64 (0.34-2.93)	3.90 (1.46-6.33)	0.38	(0.00-0.79)
Western countries ²	1.55 (0.70-2.41)	3.67 (1.49-5.85)	0.33	(0.10-0.56)
Southern countries ³	1.53 (0.65-2.42)	3.99 (2.79-5.19)	0.54	(0.08-1.00)

1: United Kingdom, Norway, Estonia

2: France, Germany, Netherlands

3: Greece, Italy, Portugal