

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Asferaw, M; Mekonen, SY; Woodruff, G; Gilbert, CE; Tesfaye, S; (2018) Outcome of paediatric cataract surgery in Northwest Ethiopia: a retrospective case series. The British journal of ophthalmology. ISSN 0007-1161 DOI: <https://doi.org/10.1136/bjophthalmol-2017-311513>

Downloaded from: <http://researchonline.lshtm.ac.uk/4647437/>

DOI: <https://doi.org/10.1136/bjophthalmol-2017-311513>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

Outcome of pediatric cataract surgery in Northwest Ethiopia: a retrospective case series

Journal:	<i>British Journal of Ophthalmology</i>
Manuscript ID	bjophthalmol-2017-311513.R2
Article Type:	Clinical science
Date Submitted by the Author:	06-Mar-2018
Complete List of Authors:	Asferaw, Mulusew; University of Gondar, Department of Ophthalmology; WGGA Institute & Eye centre, Pediatric Ophthalmology unit Mekonen, Sisay Yoseph ; University of Gondar , Department of Ophthalmology Woodruff, Geoffrey; University of Leicester , Department of Ophthalmology Gilbert, Clare; London School of Hygiene and Tropical Medicine, Clinical Research Unit, ITD Tesfaye, Samson; Orbis International-Ethiopia
Keywords:	Child health (paediatrics), Lens and zonules, Treatment Surgery

SCHOLARONE™
Manuscripts

Outcome of pediatric cataract surgery in Northwest Ethiopia: a retrospective case series**Corresponding author:**

Mulusew Asferaw, M.D

Department of Ophthalmology,
University of Gondar
Gondar, Ethiopia
E-mail: muasf@yahoo.com
Tel. +251 911806166

Co-authors

Sisay Yoseph Mekonen, B.Sc, MPH

Department of Ophthalmology,
University of Gondar
Gondar, Ethiopia
E-mail: sisayjoseph@ymail.com

Samson Tesfaye, M.D, MPH

Orbis International-Ethiopia
Addis Ababa, Ethiopia
E-Mail: samson.tesfaye@orbis.org, samitit@gmail.com

Geoffrey Woodruff, MBBS, FRCSEd, FRCOphth

Department of Ophthalmology,
University of Leicester
LE1 7RH, Leicester, UK
E-mail: geoff.woodruff@gmail.com

Clare Gilbert, FRCOphth, MD, MSc

International Centre for Eye Health,
Clinical Research Department
London School of Hygiene & Tropical Medicine
Keppel Street
London WC1E 7HT, UK
E-mail: clare.gilbert@lshtm.ac.uk

Word count: 3, 263

Number of tables: 4

Number of figures: 1

Conflict of interest declaration: None

Synopsis: Most eyes achieved 6/60 or better after surgery. In bilateral cataract cases poor outcomes were associated with nystagmus/strabismus; in traumatic cases trauma related complications were associated with poor outcomes.

Outcome of pediatric cataract surgery in Northwest Ethiopia: a retrospective case series

Mulusew Asferaw, Sisay Yoseph Mekonen, Geoffrey Woodruff, Clare Gilbert, Samson Tesfaye

ABSTRACT

Aim: To assess visual acuity outcomes, and factors associated with the outcome, of pediatric cataract surgery at the Child Eye Health Tertiary Facility, Gondar, Northwest Ethiopia.

Methods: The medical records of children aged below 16 years who underwent cataract surgery between September 2010 and August 2014 were reviewed for preoperative, surgical and postoperative data.

Results: One hundred seventy six eyes of 142 children (mean age 7.9 years \pm 4.2 [SD], 66% males) who had cataract surgery were included. Twenty-five percent (35/142) of children had bilateral cataract, 18 (13%) had unilateral non-traumatic and 89 (63%) had unilateral traumatic cataracts. An intraocular lens was implanted in 93% of eyes. Visual acuities at last follow up: bilateral cases in the better eye: good ($\geq 6/18$ or fix and follow) in 21/34 eyes (62%), borderline ($< 6/18$ - $6/60$) in 4 (12%) eyes and poor ($< 6/60$) in 9 (26%) eyes. In unilateral non-traumatic cases: good 6 eyes (33%); borderline 3 (17%) and poor 9 (50%). In unilateral traumatic cases: good 36 eyes (40%); borderline 20 (23%) and poor 33 (37%) eyes. In bilateral cataract worse outcomes were associated with preoperative nystagmus/strabismus. In traumatic cases, worse outcomes were associated with the preoperative trauma related complications.

Conclusions: Visual acuity improved significantly after surgery, with better outcomes in bilateral cases. Early detection and surgery by a trained surgeon with good follow up and postoperative rehabilitation can lead to better visual outcomes.

Key words: Pediatric cataract, visual outcome, congenital cataract, Ethiopia

INTRODUCTION

Up to a third of blindness in children in sub-Saharan Africa is due to congenital or developmental cataract, with approximately 19,000 new cases each year.[1] With the reduction of corneal blindness associated with measles and vitamin A deficiency, cataract is becoming an increasingly important cause of avoidable blindness in children in low income settings. Well documented challenges in achieving good outcomes after surgery in Africa are late presentation, surgical complications, poor follow up for refractive correction and management of visual axis opacification. In some regions there are gender differences in access.[2] However, good results can be achieved by early meticulous surgery with aggressive management of the posterior capsule, intensive treatment of post-operative uveitis and good visual rehabilitation and follow up.[3,4]

The prevalence of blindness in children in Ethiopia is estimated to be approximately 1 in 1000 children.[5] Cataract accounted for 10%[6,7] of the causes of blindness in two studies from Ethiopian schools for the blind, while a population based key informant study found cataract to be the commonest cause of blindness (33%) in Southern Ethiopia.[8] Only a few studies have reported visual acuity outcomes in children following cataract surgery in Africa[9-14] with only one study of 73 cases from Ethiopia.[15]

The World Health Organization (WHO) recommends that paediatric cataract surgery be undertaken in tertiary level eye care facilities with one facility per 10 million populations. With support from Orbis International, Gondar University Hospital established a tertiary paediatric eye unit in 2012 with a trained paediatric ophthalmologist and support team. The purpose of this study was to assess the outcome of paediatric cataract surgery at this centre. The findings will guide improvements in the management of paediatric cataract in the region and more widely.

PATIENTS AND METHODS

Study setting

The study was conducted at the University of Gondar Referral hospital, Northwest Ethiopia, which is the only tertiary eye care and referral centre for an estimated population of 20 million, almost half of whom are children. Before 2012 the standard paediatric cataract surgical procedure was extra capsular cataract extraction (ECCE), mainly in the form of manual small incision cataract surgery without primary posterior capsulotomy, performed by senior non-paediatric ophthalmologists. Since 2012 the currently recommended paediatric cataract surgery of lens aspiration with primary posterior capsulotomy (PPC) and anterior vitrectomy (AV)[16] has been performed by an ophthalmologist trained in paediatric ophthalmology. Biometry has been performed routinely. Subconjunctival steroids are given to all eyes at the end of surgery. Cataract surgery is usually performed under general anaesthesia, with local anaesthesia for older cooperative children. Most bilateral cases have both eyes operated under general anaesthesia at the same session following strict procedures (i.e., new surgical set and re-scrubbing between eyes). This approach is adopted to reduce the risk of a second anaesthesia, to ensure that second eyes are operated on without delay, and to reduce costs for parents. Children are usually seen on day 1 (at discharge), and at one week and one month after surgery.

Study design: Retrospective case series

All children aged 0-15 years undergoing cataract surgery between September 2010 and August 2014 were identified from the operating theatre logbook and their medical records were reviewed for demographic, pre-operative, intra-operative and post-operative information by the principal investigator (MA, a paediatric ophthalmologist).

Preoperative data included address, age at surgery, gender, pre-operative visual acuity (VA) and the method of measurement, type and laterality of cataract, interval between the recognition of cataract by the parents and admission to hospital for surgery, the presence of ocular and systemic comorbidities, and method of intraocular lens (IOL) power calculation. Surgical data included date of surgery, surgical technique and method of anaesthesia. Details of the surgical techniques included method of anterior capsulotomy, whether PPC and AV were performed, whether an intraocular lens (IOL) was inserted and site of placement, intra-operative complications, and surgeon profile. Postoperative complications and interventions, duration and total number of follow-up visits, and presenting VA at the last visit were also documented.

Definitions

Visual outcomes are reported as presenting VA measured at the last post-operative follow up visit and defined as good ($VA \geq 6/18$ or fix and follow (FF)), borderline ($VA < 6/18-6/60$), or poor outcome ($VA < 6/60$ or poor fixation pattern). We defined congenital cataract as a cataract manifested in the first year of life. Developmental cataract was defined as the cataract that developed after the first year of life. Postoperative complications which occurred within four weeks of surgery were defined as early, and complications after four weeks were defined as late.

Data analysis

Data were entered into Epi info version 3.3.2, and exported into SPSS version 16 for analysis. Multivariate binary logistic regression analysis was undertaken to determine predictors of postoperative VA outcomes for bilateral and traumatic cases separately. Unilateral non-traumatic cases were not included in the regression analysis because of the small number in this group. Visual outcome was defined as a dichotomous variable: Good/Borderline Outcome ($VA \geq 6/60$ or FF) or Poor outcome ($VA < 6/60$). In bilateral cases, analysis was based on the visual acuity outcome in the better-seeing eye. One child who had only one eye operated was excluded from this analysis. Variables included for multivariate regression analysis for both groups (bilateral and traumatic cataract groups) were only those with P value < 0.2 in univariate analysis, and all variables with P value < 0.2 were entered simultaneously. A p value < 0.05 was considered statistically significant.

Ethical considerations

The protocol of the study was approved by the institutional review board of the University of Gondar.

RESULTS

A total of 160 children with cataract were operated over the four year period. Complete data were available for 142 (89%) children of whom 35 (25%) had bilateral, 18 had non-traumatic unilateral,

and 89 had unilateral traumatic cataract. The mean age at surgery was 7.9 (range 0.2-15) years. Children having surgery for bilateral cataracts were younger (mean 5.8, range 0.3-15 years) than the other two groups: unilateral non-traumatic cases (mean 7.0, range 0.2-14 years) and traumatic cases (mean 9.8, range 4-15 years) years. There was a preponderance of boys (66%) over girls, particularly for traumatic cataract (71%). The children were referred from 41 districts of the four regional states. The majority of children (94%) came from Amhara regional state and most (56%) lived more than 100km from the hospital (Table 1).

Table 1: Demographic Characteristics of Children with Cataract (N= 142 children)

	Bilateral(N=35), n (%)	Unilateral non-traumatic(N=18), n (%)	Unilateral traumatic (N=89), n (%)
Age at surgery (year)			
<5	17 (48.6)	6 (33.3)	2 (2.2)
5-10	10 (28.6)	10 (55.6)	49 (55.1)
11-15	8 (22.8)	2 (11.1)	38 (42.7)
Sex			
Male	20 (57.1)	11 (61.1)	63 (70.8)
Female	15 (42.9)	7 (38.9)	26 (29.2)
Address by Region			
Amhara	32 (91.4)	15 (83.3)	87 (97.6)
Tigray	1 (2.8)	3 (16.7)	2 (2.2)
Others*	2(5.7)	-	-
Approximate distance fromhospital			
≤100 kms	18 (51.4)	6 (33.3)	39 (43.8)
>100 kms	17 (48.6)	12 (66.7)	50 (56.2)

*Southern Nations, Nationalities and People Region (1), BenshangulGumuz (1)

Visual acuity(VA)was assessed using Snellen acuity charts, Cardiff cards, LEA symbols and by assessing the fixation pattern in younger children. Among 176 eyes which had surgery, it was possible to record Snellen VA in 142 (81%) eyes pre-operatively and 148(84.1 %)eyes postoperatively. Before surgery 13 of the 34 better seeing eyes of bilateral cases could fix and follow (7 eyes), had poor fixation (6 eyes) or acuity was not recorded (1 eye); after surgery 12 of these eyes could fix and follow. Among the 18 eyes with unilateral non traumatic cataract, four had poor fixation before surgery; after surgery one eye had poor fixation and two could fix and follow. Among the 89 traumatic cases in two the visual acuity was not recorded preoperatively; after surgery 1 eye

could fix and follow. Best corrected visual acuity was not recorded in the medical records for 73 (41.5%) eyes. Overall, among the eyes recorded with Snellen VA, before surgery 5/142 eyes (4%) had a presenting VA of 6/6-6/18 and 131 (92%) had <6/60. Postoperatively 55/148 eyes (37%) had a presenting VA of 6/6-6/18 and 64 (43%) had <6/60.

For each type of cataract the proportion of children with a presenting VA of <3/60 after surgery was lower than the proportion before surgery (Figure 1). The proportion with a VA 6/6-6/18 or FF increased following surgery in all types of cataract, with better outcomes in bilateral cases compared to the other types (58% bilateral; 40% traumatic; 33% unilateral non traumatic).

Bilateral cataract

Sixty-nine eyes of 35 children with bilateral cataract were operated; one child had surgery to only one eye. Most bilateral cases (86%, 30/35 children) were operated at a single session. The interval between recognition of a problem in the eye and surgery ranged from 3 weeks to 156 months (median 25 months). Before surgery 58% of these eyes were blind (VA <3/60). Nystagmus was present in 24 (35%) eyes and strabismus in 13 (19%) eyes. The commonest type of surgery under the age of 10 years was lens aspiration with posterior capsulotomy and AV (55%). An IOL was implanted in 97% of operated eyes, mostly acrylic foldable IOLs (61%) (Table 2). The most common intraoperative complications were radial anterior capsulotomy tears (7 eyes, 10%) and posterior capsular rupture with or without vitreous loss (5 eyes, 7%). The most common early post-operative complication was fibrinous uveitis (4 eyes, 6%) whereas visual axis opacification was the commonest late post-operative complication which developed in 16 eyes (23%). Surgical capsulotomy was required in 12 eyes and YAG capsulotomy in two eyes. Glaucoma occurred in one eye which was controlled medically (Table 3).

Table 2: Pre-and intra-operative characteristics in operated eyes (N=176)

	Bilateral (N=69), n (%)	Unilateral non-traumatic (N=18), n (%)	Unilateral traumatic (N=89), n (%)
Delay to surgery (months)[†]			
Median (minimum, maximum)	25 (<1-156)	18 (<1 - 91)	3 weeks (<1-91)
≤6	9 (13.0)	5 (27.8)	72 (80.9)
>6 – 12	7 (10.1)	1 (5.6)	2 (2.2)
12 -18	18 (26.1)	3 (16.6)	8 (9.0)
>18	35 (50.7)	9 (50.0)	7 (7.9)
Preoperative presenting VA			
≥6/18 or fixes and follows	18 (26.1)	-	-
<6/18 - 6/60	4 (5.8)	1 (5.6)	1 (1.1)
<6/60 or poor fixation pattern [§]	47 (68.1)	17 (94.4)	88 (98.9)
Associated ocular co-morbidity			
Nystagmus	24 (34.8)	1 (5.6)	-
Strabismus	13 (18.8)	7 (38.9)	2 (2.2)
Corneal scar/opacity	-	-	26 (29.2)
Posterior/anterior synechiae	-	-	14 (15.7)
Corneal tear	-	-	14 (15.7)
Iris sphincter tear	-	-	3 (3.4)

Traumatic mydriasis	-	-	3 (3.4)
Iridodialysis	-	-	3 (3.4)
Dislocated/subluxated lens	-	1 (5.6)	2 (2.2)
Corneal foreign body	-	-	1 (1.1)
Retinal detachment	-	-	1 (1.1)
Surgical procedure			
LA+PPC+AV+PC IOL	38 (55.1)	9 (50.0)	28 (31.5)
LA+PC IOL	27 (39.1)	7 (38.9)	51 (57.3)
LA+PPC+PC IOL	2 (2.9)	-	1 (1.1)
Lensectomy with no IOL	2 (2.9)	2 (11.1)	2 (2.2)
LA with no IOL	-	-	6 (6.7)
LA +AC IOL	-	-	1 (1.1)
Type of anterior capsulotomy			
Vitrectorhexis	51 (73.9)	13 (72.2)	44 (49.4)
V-capsulotomy	16 (23.2)	5 (27.8)	44 (49.4)
CCC	2 (2.9)	-	1 (1.1)
Type of IOL			
Foldable	42 (60.9)	9 (50.0)	35 (39.3)
PMMA	25 (36.2)	7 (38.9)	45 (50.6)
No IOL	2 (2.9)	2 (11.1)	9 (10.1)
Site of IOL implanted			
In the bag	55 (79.7)	7 (38.9)	36 (40.4)
Sulcus	7 (10.1)	7 (38.9)	21 (23.6)
AC	-	-	1 (1.1)
No IOL	2 (2.9)	2 (11.1)	9 (10.1)
Unknown	5 (7.2)	2 (11.1)	22 (24.7)

[†]Interval between recognition of cataract in the eye by care givers and admission to hospital for surgery; [§]Four eyes with undetermined vision (2 in bilateral, 2 in unilateral traumatic) were included under VA<6/60 or poor fixation category.

VA: Visual Acuity; LA: Lens Aspiration; PPC: Primary Posterior Capsulotomy, AV: Anterior Vitrectomy; PC IOL: Posterior Chamber IntraOcular Lens.; ACIOL: Anterior Chamber Intra Ocular Lens; PC:Posterior Capsule; PMMA: polymethylmethacrylate; CCC: continuous curvilinear capsulorhexis

Table 3: Surgical complications, postoperative interventions and number of follow-up visits

	Bilateral (N=69 eyes) n (%)	Unilateral non- traumatic (N=18 eyes) n (%)	Unilateral traumatic (N=89 eyes), n (%)
Intra operative complication			
Radial tear of capsulotomy	7 (10.1)	1 (5.6)	10 (11.2)
PC rupture without vitreous loss	5 (7.2)	2 (11.3)	1 (1.1)
Vitreous loss only	1 (1.4)	-	1 (1.1)
PC rupture with vitreous loss	-	4 (22.2)	4 (4.5)
Early postoperative complication (up to four weeks)			
Fibrinous uveitis	4 (5.8)	2 (11.1)	19 (21.3)
Striate keratopathy/corneal oedema	2 (2.9)	2 (11.1)	16 (18.0)
Lens matter remnant	1 (1.4)	-	7 (7.9)
Pupillary capture of IOL	1 (1.4)	1 (5.6)	-
IOL decentration	-	-	5 (5.6)
Endophthalmitis	-	-	3 (3.4)
Iris prolapsed	-	-	2 (2.2)
Shallow anterior chamber	-	-	1 (1.1)
Late postoperative complication (after four weeks)			
Visual axis opacification	16 (23.2)	1 (5.6)	13 (14.6)
Pupillary capture of IOL	4 (5.8)	-	2 (2.2)
Decentration of IOL	2 (2.9)	1 (5.6)	1 (1.1)
Glaucoma	1 (1.4)	-	1 (1.1)
Secondary membrane	-	-	1 (1.1)
Vitreous haemorrhage	-	-	1 (1.1)
Postoperative intervention			
<u>Surgical</u>			
Surgical capsulotomy/membranectomy	12 (17.4)	2 (11.1)	10 (4.2)
IOL repositioning	5 (7.2)	2 (11.1)	3 (3.4)
YAG capsulotomy	2 (2.9)	-	4 (4.5)
Excision of limbal conjunctival cyst	1 (1.4)	-	-
Aspiration of lens matter	-	-	2 (2.2)
IOL replacement	-	-	1 (1.1)
<u>Medical</u>			
Spectacle prescribed	28 (40.6)	6 (33.3)	14 (15.7)
Patching therapy	23 (33.8)	9 (50.0)	8 (9.0)
Systemic steroid	2 (2.9)	1 (5.6)	12 (13.5)
Total number of follow-up visits			
1	8 (11.6)	2 (11.1)	16 (18.0)
2	20 (29.0)	6 (33.3)	26 (29.2)
3	20 (29.0)	5 (27.8)	28 (31.5)
4	7 (10.1)	2 (11.1)	9 (10.1)
5	4 (5.8)	2 (11.1)	5 (5.6)
6	8 (11.6)	1 (5.6)	2 (2.2)
7	2 (2.9)	-	3 (3.4)

PC:Posterior capsule; IOL: Intra Ocular Lens

1
2
3
4
5 Follow up after surgery ranged from day 1 to 33 months (median 2.8 months) and 41 (59%) had at
6 least three follow up visits. Among 34 cases with bilateral cataract where better seeing eyes were
7 analyzed, 17 (50%) children had follow up for 3 months or more and 17 (50%) had less than 3
8 months of follow up.

9 Visual outcome

11 **Total operated eyes:** At their last recorded follow up, among 45 eyes with quantitative Snellen
12 acuity, 16/45 eyes (36%) had a presenting VA of 6/6 - 6/18, 6 eyes (13%) had 6/24-6/60, and 23 eyes
13 (51%) had <6/60. In 24/69 (35%) bilateral cataract eyes, a Snellen acuity could not be measured post
14 operatively, but the eye was able to fix and follow. If these eyes are categorized as having a good
15 outcome, the number of bilateral cataract eyes achieving a good outcome is 40 (58%), borderline 6
16 (8%), and poor outcome 23 (33%). There was no statistically significant difference in postoperative
17 visual acuity between cataracts classified as developmental or congenital ($p=0.62$).

19 **Better eye:** Among 34 children with both eyes operated, visual outcome in the better-seeing eye was
20 analyzed: good outcome ($\geq 6/18$ or FF) in 21 (62%) of children, borderline in 4(12%), and poor
21 outcome in 9 (26%).
22
23
24

25 **Unilateral non-traumatic cataract**

26
27 Before surgery 94% (17/18) of operated eyes were blind ($VA < 3/60$). Strabismus was present in
28 seven eyes (39%) and nystagmus in one eye. The delay to surgery ranged from 1 week to 91 months
29 (median 18 months). The commonest type of surgery in this group was lens aspiration with PPV and
30 AV in nine eyes (50%) followed by lens aspiration without posterior capsulotomy. An IOL was
31 implanted in 89 % eyes. The commonest intraoperative complication was posterior capsular rupture
32 with vitreous loss, which occurred in 4 eyes (22%). Fibrinous uveitis and striate keratopathy
33 occurred in two eyes each. Visual axis opacification was a rare (6%) complication in this group
34 (Table 3).
35
36

37 Follow up after surgery ranged from 1 day to 6 months (median 1 month), 14 (77.8%) had less than 3
38 month follow up, only 4 (22.2%) had follow up for 3 or more months and 56 % of eyes had at least
39 three follow up visits. At their last follow up, four eyes (22%) achieved 6/6-6/18, two (11%) fix and
40 follow, one (5.6%), three (17%) 6/24-6/60 and 8 eyes (44%) had <6/60. In summary, six eyes (33%)
41 had good, three (17%) borderline, and nine eyes (50%) poor visual outcome in this group.
42
43

44 **Traumatic cataract**

45 A corneal tear repair was required at the time of cataract surgery in 13/89(14.7%) eyes. There was
46 multiple other trauma related ocular pathology including corneal scar or opacity in 26 (29.2%) eyes
47 and synechiae in 14 eyes (15.7%). Preoperatively, 85 eyes (95.5%) were blind ($VA < 3/60$). The
48 commonest surgical procedure was lens aspiration without vitrectomy in 51 eyes (57.3%) and an IOL
49 was not implanted in nine eyes (10.1%) (Table 2). The commonest intraoperative complication was
50 radial tear of anterior capsulotomy (10 eyes, 11.2%) whereas the most frequent postoperative
51 complications were fibrinous Uveitis (21.3%), corneal edema (18.0%) and visual axis opacification
52 (14.6%). Endophthalmitis occurred in three eyes (3.4%) (Table 3).
53
54
55
56
57
58
59
60

Follow up after surgery ranged from 1 day to 12.4 months (median 1.2 month), 63 (70.8%) of children had less than 3 month follow up, 26 (29.2%) had follow up for 3 or more months, and 52.8% of the eyes had three or more follow ups. After surgery, visual outcome was 6/6-6/18 in 35 eyes (39%), fix and follow in one eye (1%), 6/24-6/60 in 20 eyes (22%), and <6/60 in 33 eyes (37%).

Factors associated with visual outcome

Bilateral cases:

Regression analysis was carried out to determine factors associated with poor visual outcome in the better seeing eye in the 34 children with bilateral cataract. The variables that were analyzed are shown on table 4. In univariate and multivariate analysis, only the presence of ocular co-morbidities i.e., strabismus or nystagmus, was significantly associated with poor outcome (<6/60) than those without co-morbidities (Odds ratio (OR) 10.63 (95% CI, 1.22 -92.34, p=0.03).

Table 4: Factors associated with poor visual outcome in children with bilateral developmental or congenital cataract based on the better-seeing eye (N=34).

	Good or Borderline Outcome (≥6/60/FF)	Poor Outcome (<6/60)	Crude OR (95% CI)	P- value	Adjusted OR (95% CI)	P- value
Age at surgery (year) Mean=5.7, SD=4.6	-	-	1.01 (0.85 – 1.19)	0.94	-	-
Gender						
Male	13	6	0.93 (0.24 – 3.50)	0.91	-	-
Female	12	3	Reference			
Type of cataract						
Developmental	10	5	1.88 (0.40 – 8.74)	0.42	-	-
Congenital	15	4	Reference			
Delay to Surgery (month)[†] Mean=3.4, SD=3.4	-	-	1.02 (0.99 – 1.05)	0.09	1.03 (0.99 – 1.07)	0.09
Ocular co-morbidities(nystagmus/strabismus)						
Present	6	7	11.08 (1.80 – 68.40)	0.01	10.63 (1.22 – 92.34)	0.03
Absent	19	2	Reference		Reference	
PPC						
Not performed	12	1	0.50 (0.05 – 4.98)	0.08	0.07 (0.002 – 1.99)	0.12
Performed	13	8	Reference		Reference	
Intra-operative complications						
Present	5	1	0.07 (0.002 – 1.99)	0.55	-	-

Absent	20	8	Reference		
Postoperative complications					
Present	10	1	0.19 (0.02 – 1.74)	0.14	0.42 (0.03 – 5.75)
Absent	15	8	Reference		Reference
Duration of follow-up (months)			0.80 (0.55 – 1.16)	0.24	-
Mean=5.8, SD=9.2					

CI: Confidence Interval; PPC: Primary Posterior Capsulotomy; AV: Anterior Vitrectomy

*indicates interval between recognition of cataract in the eye by care givers and admission to hospital for surgery

Traumatic cases:

Only pre-existing trauma related ocular co-morbidity was significantly associated with poor outcome in univariate and multivariate analysis in traumatic cataracts (OR = 6.23; 95% CI 2.01 – 19.25, p=0.001).

DISCUSSION

Over half of eyes of better seeing eyes in bilateral cases in our series achieved a VA $\geq 6/18$, or could fix and follow. It is difficult to compare our study with others because of the difficulties in recording VA in children, particularly in young children and this is reflected in the large proportion of children with bilateral cataract in whom a Snellen acuity was not recorded. Children recorded as being able to fix and follow but without a Snellen VA were included in our good visual outcome category but the imprecision of this acuity assessment is a limitation of our study. It has been suggested that younger children may have better results because they may have had less delay between the onset of cataract and treatment, or alternatively, that they may have worse results because of the onset of cataract at a younger age. The limitations of our data prevent us from definitively resolving this issue, and the classifications of cases as congenital or developmental were based on parental recall. We used presenting VA to assess visual outcome as best corrected VA was not recorded in 41.5% of eyes implying that refraction may not have been undertaken at every visit. In addition, our intention was to include whether surgery was performed by a trained pediatric ophthalmologist or a qualified ophthalmologist experienced in adult surgery in the regression analysis but this was not possible as some cells contained no data.

Our results for good outcome in the better eye (62%) among bilateral cases are comparable to studies in Tanzania (62%),^[10] Nepal and Northern India (43%),^[4] Kenya (44%),^[9] South India (39.5%),^[17] UK (40.6%),^[18] and USA (46.6%),^[19] but much better than in studies from Zambia (29.7%)^[13] and South Africa (24.7%).^[14] Good visual outcome was achieved more often in bilateral cases compared to unilateral non-traumatic cases.

Only 33% of eyes in unilateral non-traumatic cataract achieved good outcome ($\geq 6/18$) and with half of the eyes having a poor visual outcome ($<6/60$) with median follow up period of one month. This finding is similar to the study in India where best corrected visual acuity was 6/18 or better in 25% and $<6/60$ in 66.7% of unilateral congenital and developmental cataracts at mean follow up of 4.4 years.^[20] The poor outcome in a significant number of eyes in unilateral cases can be explained by deep deprivational amblyopia.

1
2
3 Thirty-five eyes (39%) with traumatic cataract had a VA of 6/18 or better. This is slightly lower than
4 in studies in Zambia (68%),[13] and the previous study in Southern Ethiopia (64.7%).[21]
5

6 There was a substantial improvement in VA for each type of cataract, with the greatest improvement
7 being in traumatic cases, no doubt reflecting the lower risk of amblyopia in children who developed
8 cataract later in childhood. The finding of a relatively good acuity in a few children with non-
9 traumatic cataract may be because of undetected trauma, or later onset of cataract.
10

11 In this study, the mean age at surgery was 5.8 years and 7 years in bilateral and unilateral non-
12 traumatic cases, respectively. Children with traumatic cataract were older (mean 9.8 years). This is
13 similar to the findings in the previous study in Ethiopia(mean 7.1yr)[15] and other studies in
14 developing countries.[3, 10-12] However, the age at surgery for non-traumatic cases in our study was
15 higher than a report from developed countries.[18] A study from Kenya also showed a younger age
16 at surgery compared to our study.[9]
17
18

19 To achieve better visual outcome in children, earlier cataract surgery is recommended. However, the
20 finding in this study showed late presentation with a median delay to surgery of 25 months and 18
21 months in bilateral and unilateral non-traumatic cases, respectively. This delay will result in
22 significant deprivational amblyopia, reflected in the high rates of nystagmus and strabismus which
23 were associated with poorer outcomes. A mean delay of almost 3 years between first recognition of
24 an abnormality by parents/caregivers and surgery for congenital and developmental cataract was
25 reported in a study from Tanzania.[22] Delay in presentation was associated with developmental
26 cataract, living far from the hospital and low socio-educational status of the mother. Complex
27 interactions between socio-cultural barriers at family and community level as well as socio-
28 organizational barriers within the health care system also contributed to delay in presentation.[23]
29
30

31 However, our study showed that delay between presentation and surgery did not greatly affect
32 outcomes. This is in line with two studies from India[24,25] but unlike results from China.[26]
33

34 The high proportion of traumatic cataract in this series is worth noting, being higher than studies
35 from Southern Ethiopia (36.3%),[15] Madagascar (25.6%),[11] and Nepal(23.9%).[3] In contrast, in
36 high income settings, trauma accounts for only 10-29% of cases.[27] The high proportion of
37 traumatic cataract in our series may be because in Ethiopia children are often involved in agriculture,
38 and the sudden pain and loss of vision will stimulate different health seeking behavior by parents
39 compared to that for children with longstanding, painless loss of vision for which parents may have
40 no explanation or attribute to spiritual causes. The finding from the recent key informant study that
41 bilateral cataract was the leading cause of blindness in children indicates the need for active
42 community-based case finding with referral for assessment, surgery and counseling.[8]
43
44

45 In this series follow up after cataract surgery was inadequate. Before surgery parents need to be
46 counseled that surgery is only the first stage, and that postoperative visual rehabilitation that includes
47 amblyopia therapy, correction of refractive error and low vision therapy are all important. Barriers to
48 follow up have been reported from Tanzania and included being a girl with bilateral cataract, poorly
49 educated mothers, a large number of siblings and delay in initial presentation for surgery.[28]
50 Multiple strategies were used to improve follow up, including reimbursement of transport costs, a
51 dedicated childhood blindness coordinator, tracking children after surgery, good counseling of
52 parents, and use of cell phones reminders.[29]
53
54
55
56
57
58
59
60

1
2
3 In this study limitations included poor documentation of relevant details in medical records, lack of
4 standard definitions, methods of measuring VA, and difficulty in distinguishing congenital from
5 developmental cataract. Selection bias is also likely, as severely injured eyes might have developed
6 endophthalmitis or not be operated on if the prognosis was poor. Visual acuity findings have to be
7 interpreted cautiously, given the relatively short follow up. Electronic medical records may improve
8 documentation.
9

10 11 12 **CONCLUSIONS**

13 The short term visual outcome after cataract surgery was fair as the majority achieved a VA of 6/60
14 or better. There was a significant delay in presentation particularly in non-traumatic cases. The
15 presence of nystagmus/strabismus in bilateral cases and trauma related complications in traumatic
16 cases were predictors of poor visual outcome. To address cataract blindness active case finding
17 combined with counseling parents and clear referral pathways are needed. Surgery should be
18 undertaken by a pediatric ophthalmologist in a fully equipped unit, with trained support staff, and
19 multiple strategies are likely to be required to improve follow up after surgery. Prospective studies
20 are being set up to address some of the limitations identified in this case series.
21
22
23

24 **Contributors** MA, SY, and ST conceived of the study. MA collected the data. SY did statistical
25 analysis of the data. MA, GW and CG further analyzed the data. All the authors contributed to the
26 manuscript and approved the final draft for submission.
27

28 **Competing interests** None.
29

30 **Data sharing statement** No additional data are available.
31

32 **Ethical approval** Institutional review board of the University of Gondar.
33

34 **Funding** This research received no specific grant from any funding agency in the public, commercial
35 or not-for-profit sectors.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Courtright P. Childhood cataract in sub-Saharan Africa. *Saudi J Ophthalmol.* 2012;26: 3–6.
2. Gilbert CE, Lepvrier-Chomette N. Gender Inequalities in Surgery for Bilateral Cataract among Children in Low-Income Countries. *Ophthalmology* 2016;123:1245-51.
3. Wilson ME, Hennig A, Trivedi RH, et al. Clinical characteristics and early postoperative outcomes of pediatric cataract surgery with IOL implantation from Lahan, Nepal. *J Pediatr Ophthalmol Strabismus.* 2011; 48:286-91.
4. Hennig A, Schroeder B, Gilbert C. Bilateral Pediatric Cataract Surgery: Outcomes of 390 Children from Nepal and Northern India. *J Pediatr Ophthalmol Strabismus* 2013; 50(5):312- 19.
5. Berhane Y, Worku A, Bejiga A, et al. Prevalence and causes of blindness and low vision in Ethiopia. *Ethiop J Health Dev* 2007;21:204–10.
6. Kello AB, Gilbert C. Causes of severe visual impairment and blindness in children in schools for the blind in Ethiopia. *Br J Ophthalmol.* 2003;87(5):526-30.
7. Asferaw M, Woodruff G, Gilbert C. Causes of severe visual impairment and blindness in students in schools for the blind in Northwest Ethiopia. *BMJ Glob Health* 2017;2:e000264. doi:10.1136/bmjgh-2016-000264.
8. Berhan S. Demissie. Magnitude and causes of childhood blindness and severe visual impairment in Sekoru District, Southwest Ethiopia: a survey using the key informant method. *Trans R Soc Trop Med Hyg.* 2011;105:507– 11.
9. Yorston D, Wood M, Foster A. Results of cataract surgery in young children in East Africa. *Br J Ophthalmol.* 2001;85(3):267-71.
10. Bowman RJ, Kabiru J, Negretti G, Wood ML. Outcomes of bilateral cataract surgery in Tanzanian children. *Ophthalmology.* 2007;114(12):2287-92.
11. Randrianotahina HCL, Nkumbe HE. Pediatric cataract surgery in Madagascar. *Nigerian J clinical practice.* 2014; 17(1):14-7.
12. Umar M, Abubakar A, Achi I, Alhassan MB, et al. Pediatric cataract surgery in National Eye centre Kaduna, Nigeria: Outcome & challenges. *Middle East Afr J Ophthalmol.* 2015; 22(1): 92-6.
13. Mboni C, Gogate PM, Phiri A, et al. Outcomes of Pediatric Cataract Surgery in the Copperbelt Province of Zambia. *J Pediatr Ophthalmol Strabismus.* 2016;53(5):311-7.
14. Gogate P, Parbhoo D, Ramson P, et al. Surgery for sight: outcomes of congenital and developmental cataracts operated in Durban, South Africa. *Eye.* 2016; 30(3): 406–12.
15. Tomkins O, Itay BZ, Moore DB, Helveston EE. Outcomes of Pediatric Cataract Surgery at a Tertiary Care Center in Rural Southern Ethiopia. *Arch Ophthalmol.* 2011;129 (10):1293-7.

16. Wilson ME, Pandey SK, Thakur J. Paediatric cataract blindness in the developing world: surgical techniques and intraocular lenses in the new millennium. *Br J Ophthalmol*. 2003;87(1):14-9.
17. Khanna R, Foster A, Krishnaiah S, et al. Visual outcomes of bilateral congenital and developmental cataracts in young children in south India and causes of poor outcome. *Indian J Ophthalmol*. 2013;61:65-70.
18. Chak M, Wade A, Rahi JS. British Congenital Cataract Interest Group. Long-term visual acuity and its predictors after surgery for congenital cataract: findings of the British Congenital cataract Study. *Invest Ophthalmol Vis Sci*. 2006;47:4262-9.
19. Ledoux DM, Trivedi RH, Wilson ME, Payne JF. Pediatric cataract extraction with intraocular lens implantation: visual acuity outcome when measured at age four years and older. *J AAPOS*. 2007;11:218-24.
20. Gogate P, Patil S, Kulkarni A, Sahasrabudhe M, et al. Unilateral Congenital and Developmental Cataracts in Children in India: How Useful Were Long-Term Outcomes of Surgery? Miraj Pediatric Cataract Study 4. *Asia Pac J Ophthalmol (Phila)*. 2015;4(6):376-80.
21. Kinori M, Tomkins-Netzer O, Wagnanski-Jaffe T, Ben-zion I. Traumatic pediatric cataract in southern Ethiopia-results of 49 cases. *J AAPOS*. 2013;17:512-15.
22. Mwende J, Bronsard A, Mosha M, Bowman R, Geneau R, Courtright P. Delay in presentation to hospital for surgery for congenital and developmental cataract in Tanzania. *Br J Ophthalmol*. 2005;89(11):1478-1482.
23. Bronsard A, Geneau R, Shirima S, Courtright P et al. Why are Children Brought Late for Cataract Surgery? Qualitative Findings from Tanzania. *Ophthalmic Epidemiology*, 2008; 15:383-388.
24. Gogate p, Khandekar R, Shrishrimal M, et al. Delayed Presentation of Cataracts in Children: Are they Worth Operating Upon? *Ophthalmic Epidemiology*. 2010;17(1) :25-33.
25. Gogate PM, Sahasrabudhe M, Shah M, et al. Long term outcomes of bilateral congenital and developmental cataracts operated in Maharashtra, India. Miraj pediatric cataract study III. *Indian J Ophthalmol*. 2014 ;62(2):186-95.
26. You C, Wu X, Zhang Y, Dai Y, Huang Y, Xie L. Visual impairment and delay in presentation for surgery in Chinese pediatric patients with cataract. *Ophthalmology*. 2011;118:17-23.
27. Wilson ME Jr, Trivedi RH, Pandey SK. Traumatic cataracts in children. In: Wilson ME Jr, Trivedi RH, Pandey SK, eds. *Pediatric Cataract Surgery*. Philadelphia, PA: Lippincott Williams & Wilkins; 2005.
28. Eriksen JR, Bronsard A, Mosha M, Carmichael D. Predictors of Poor Follow-up in Children that had Cataract Surgery. *Ophthalmic Epidemiology*. 2006;13:237-243.

- 1
2
3 29. Kishiki E, Shirima S, Lewallen S, Courtright P. Improving postoperative follow-up of children
4 receiving surgery for congenital or developmental cataracts in Africa. *J AAPOS* 2009;13:280 -2.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential: For Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Percent of eyes with each category of VA before and after cataract surgery.

203x143mm (300 x 300 DPI)

Review Only