

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Ranke, J; Evans, JR; Gilbert, CE; (2018) Reducing inequity of cataract blindness and vision impairment is a global priority, but where is the evidence? The British journal of ophthalmology. ISSN 0007-1161 DOI: <https://doi.org/10.1136/bjophthalmol-2018-311985>

Downloaded from: <http://researchonline.lshtm.ac.uk/4647928/>

DOI: <https://doi.org/10.1136/bjophthalmol-2018-311985>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

**Reducing inequity of cataract blindness and vision
impairment is a global priority, but where is the evidence?**

Journal:	<i>British Journal of Ophthalmology</i>
Manuscript ID	bjophthalmol-2018-311985.R1
Article Type:	Global issues
Date Submitted by the Author:	n/a
Complete List of Authors:	Ramke, Jacqueline; University of Auckland, Epidemiology and Biostatistics Evans, Jennifer; London School of Hygiene and Tropical Medicine, International Centre for Eye Health Gilbert, Clare; London School of Hygiene and Tropical Medicine, Clinical Research Unit, ITD
Keywords:	Public health

SCHOLARONE™
Manuscripts

Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Reducing inequity of cataract blindness and vision impairment is a global priority, but where is the evidence?

Authors

Jacqueline **Ramke**^{1,2}
Jennifer R. **Evans**²
Clare E. **Gilbert**²

- 1. University of Auckland, School of Population Health, Department of Epidemiology & Biostatistics, Auckland, New Zealand
- 2. London School of Hygiene and Tropical Medicine, Clinical Research Unit, Department of Infectious & Tropical Diseases, London, UK

Corresponding author

Jacqueline Ramke jramke@gmail.com
Dept of Epidemiology & Biostatistics, Faculty of Medical and Health Sciences, The University of Auckland.
Private Bag 92019 Auckland Mail Centre
Auckland 1142, New Zealand
Phone +6421 029 58543

Word count: 1279

For Review Only

SUMMARY

Throughout the world, people who are socially or economically disadvantaged disproportionately experience blindness and vision impairment caused by cataract. Reducing vision loss from cataract and its unequal distribution must be a priority if the World Health Organization's aim of 'universal eye health' is to be realised. To help achieve this, decision-makers and service planners need evidence on which strategies improve access to cataract services among disadvantaged populations, and under what circumstances. Unfortunately, despite many strategies to improve cataract services being implemented in recent decades, evidence of what works, for who, and in what circumstances is not readily available. This paper summarises the extent of the evidence on interventions to reduce inequity of vision loss from cataract, and makes suggestions for how the evidence-base can be strengthened.

MAIN TEXT***Inequity in access to cataract services***

Cataract is the leading cause of blindness globally, and a major cause of moderate and severe vision impairment, affecting an estimated 65 million people in 2015.¹ Reducing cataract blindness and vision impairment is, therefore, a priority in the *Universal Eye Health: Global Action Plan 2014–2019* endorsed at the 66th World Health Assembly in 2013.²

Vision loss from cataract is unequally distributed throughout the world. For example, in 2015 among adults 50 years and above, the age standardized prevalence of cataract blindness ranged from 0.08% (80% uncertainty interval [UI] 0.03–0.19%) in high income countries of the Asia Pacific region to 2.35% (80% UI 0.72–5.04%) in West sub-Saharan Africa—almost a 30-fold difference.¹ Inequality (i.e., measurable differences between population subgroups) is also evident within countries, with a higher prevalence of cataract blindness among socially disadvantaged groups such as women, rural dwellers, and people who are not literate.³ These inequalities are avoidable and unjust and are referred to as inequities.⁴

Cataract surgery is usually highly effective, with good outcomes. Inequity in cataract blindness is, therefore, primarily due to differential access to quality eye care. To reduce this inequity, the provision of good quality cataract surgery must increase among socially disadvantaged people at a faster rate than among the more advantaged. To help achieve this, decision-makers and service planners need to understand which strategies improve access to cataract services among disadvantaged populations, and under what circumstances.

Reducing inequity: what is the evidence?

Cochrane Eyes and Vision recently published a systematic review that summarises the available evidence on strategies to improve access to cataract services in low- and middle-income countries (LMICs) and their impact on inequity.⁵ The purpose of the review was to explore the effectiveness of interventions that targeted disadvantaged populations, or reported the distribution of effects across population subgroups.

Exhaustive searching for publications with relevant study designs (i.e., randomised and quasi-randomised controlled trials, controlled before-and-after or interrupted time series studies) identified

1
2
3 only two publications. Both were cluster-randomised controlled trials conducted in rural China, and
4 both recruited adults with vision loss from cataract. In one study the intervention was additional
5 information and counselling about cataract and its treatment, compared with usual care.⁶ The other
6 study explored the impact of providing free cataract surgery and help with transport costs compared
7 with low-cost cataract surgery and no support for transport.⁷ The evidence was judged to be of low
8 certainty because the effect estimates were imprecise, and because findings from rural China may not
9 be applicable to other settings. Bearing this in mind, the findings suggest that offering more
10 information or counselling may not improve uptake of referral or surgery, and that offering free
11 cataract surgery may increase uptake. Neither study measured the effect of the interventions on the
12 prevalence of vision impairment from cataract in the community, nor whether effects were equitable.⁵
13
14
15
16
17
18
19
20
21
22

23 ***Strengthening the evidence***

24 Ideally decision-makers could draw on evidence from a range of contexts when planning services for
25 cataract. However, despite the implementation of strategies to improve cataract services in recent
26 decades, evidence of what works, for who, and in what circumstances is not readily available. This
27 problem is not unique to eye health—the recognised mismatch between the large volume of research
28 describing health inequity compared to the number of studies reporting interventions to reduce
29 inequity⁸ has partly been attributed to the evaluation of such interventions being more
30 methodologically difficult.⁹
31
32
33
34
35
36
37

38 While only two studies met the inclusion criteria for the systematic review, the search identified
39 additional studies describing strategies to increase uptake of cataract services which were excluded
40 as they did not meet the strict criteria for inclusion e.g. they had insufficient data collection time-points
41 before and after the intervention to be included as an interrupted time series study.⁵ Some of these
42 publications described promising strategies that warrant more rigorous evaluation and some
43 suggestions for strengthening the available evidence are offered in Table 1.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Opportunities to strengthen the evidence on reducing inequality in vision loss from cataract in low- and middle-income countries

Study component	Key opportunities
Population	<ul style="list-style-type: none"> • When planning studies, include subgroup analysis to assess equity implications of interventions across relevant PROGRESS axes* (and calculate sample sizes accordingly) • Use the Equity Tool (www.equitytool.org) or similar to assess outcomes by socioeconomic status
Intervention	<ul style="list-style-type: none"> • Test supply- as well as demand-side interventions (individually and in combination) • Use qualitative methods to identify scalable community-led strategies to test (e.g. local volunteer escorts; local motivators who have had successful cataract surgery; allow cataract patients to share transport with those travelling to market; establish a group of local philanthropists to cover costs; establish networks of case finders/escorts)
Comparison	<ul style="list-style-type: none"> • Use step-wedge designs and make comparisons over time—phase in interventions so that ultimately all participants receive the intervention by the last study step • Undertake sequential trials with embedded evaluations to iteratively refine implementation interventions¹⁰
Outcome	<ul style="list-style-type: none"> • Extend outcomes beyond uptake of service: <ul style="list-style-type: none"> - Plan longer-term studies (e.g. between two prevalence surveys) to assess policy-level interventions, and enable assessment of the prevalence of cataract blindness as an outcome - Include post-operative visual outcome (as a measure of intervention quality) whenever possible
Study Design	<ul style="list-style-type: none"> • Use quasi-experimental studies alongside experimental studies • Establish 'implementation laboratories'¹⁰ to make head-to-head comparisons between different settings • As more primary studies become available, synthesize evidence in systematic reviews
Setting	<ul style="list-style-type: none"> • Undertake primary studies in more contexts • Adopt more comprehensive ways to describe context

*PROGRESS: **P**lace of residence; **R**ace/ethnicity/ culture/ language; **O**ccupation; **G**ender/sex; **R**eligion; **E**ducation; **S**ocioeconomic status; **S**ocial capital/networks.¹¹

In particular, more implementation research has been called for,^{9 12} which aims to understand "what, why, and how interventions work in real world settings and test approaches to improve them".¹³

Quasi-experimental studies, such as difference-in-differences or interrupted time series,¹⁴ warrant particular consideration. In these studies participants are not randomly allocated to different interventions, and outcomes are either compared between groups after a period of time to measure

1
2
3 the difference in differences, or change is measured within the same group over time (i.e., before and
4 after the intervention).¹⁵
5
6

7 When testing strategies to address inequity in access to cataract surgery, quasi-experimental studies
8 can generate knowledge on what works and what does not in 'real world' settings. Quasi-experiments
9 do not alter the context in which data are generated, producing externally valid results, and can be
10 used when rolling out large scale programs. As these types of studies can use routinely collected
11 outcome data they can be faster and lower-cost than randomised controlled trials.¹⁶
12
13
14

15 Quasi-experimental methods could also be used within 'implementation laboratories', which have
16 been proposed as a way to assess the design and delivery of interventions on a large scale, through
17 collaboration between health service implementers and researchers.¹⁰ Ideally studies on strategies to
18 reduce inequity in vision loss from cataract would occur between two blindness prevalence surveys
19 that disaggregate outcomes by wealth quintiles and gender, so the social distribution of cataract
20 blindness in the target population is known at the two time points. Sequential trials of promising
21 strategies can then be tested—such as different models of financial support, variation in the location
22 and frequency of outreach, and community-led solutions (see Table 1)—with outcome data
23 disaggregated by wealth quintile and gender to assess equity impacts. Embedded process
24 evaluations of the interventions would provide knowledge on how the context, intervention design and
25 delivery influence effectiveness.¹⁰ If implementation laboratories were established in different settings,
26 knowledge could be generated on how health system characteristics and cultural and socioeconomic
27 factors modify the effects of the intervention.¹⁰
28
29
30
31
32
33
34
35
36
37
38
39

40 The evidence gap identified in the Cochrane review⁵ means resources may be wasted on ineffective
41 strategies, and opportunities are being missed to scale-up effective strategies. The *Universal Eye*
42 *Health: Global Action Plan 2014–2019* calls for more evidence to be generated and emphasis is given
43 to epidemiological studies such as cross-sectional prevalence surveys.² While we agree that
44 epidemiological evidence is essential to plan and monitor services, we urge researchers and funders
45 to also generate evidence on how best to implement services to reduce vision loss from cataract and
46 its unequal distribution.
47
48
49
50
51
52

53 Researchers must overcome a number of barriers to successfully generate evidence that can be used
54 to reduce inequity in vision loss from cataract. These barriers include weak health information
55
56
57
58
59
60

1
2
3 systems, a lack of routinely collected eye health data, and limited capacity to undertake and interpret
4 implementation research at the local level. Taking a collaborative approach can help address capacity
5 shortfalls. Indeed, one of the most important factors to promote the use of evidence are relationship-
6 and skills-building between researchers, implementers and policy-makers.¹⁷ An example of such a
7 collaboration in Mozambique recently demonstrated how health information system data can be used
8 in quasi-experiments, and shows what might be possible if similar efforts were attempted with
9 routinely generated eye health information.¹⁸

10
11
12 To realise Universal Eye Health more and better evidence is needed. Major innovation is required,
13 and researchers have a responsibility to learn and apply the most relevant methods to generate the
14 required evidence, which includes quasi-experimental methods¹⁴ and implementation research.
15
16 Governmental and non-governmental eye care service providers can improve the quality of data they
17 collect, and make it more readily available to researchers.¹⁴ Crucially, effective collaborations
18 between researchers and health system partners are required to produce generalisable evidence on
19 how to reduce inequity in vision loss from cataract.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Completing interests: None of the authors has any conflict of interest to disclose

Financial support: This manuscript received no specific funding.

References

1. Flaxman S, Bourne R, Resnikoff S, et al. Global Causes of Distance Vision Loss: 1990-2015 and projections to 2020. *Lancet Glob Health* 2017;5(12):e1221-e34.
2. World Health Organization. Universal Eye Health: A global action plan 2014-2019. Geneva: World Health Organization, 2013.
3. Ramke J, Zwi AB, Lee AC, et al. Inequality in cataract blindness and services: moving beyond unidimensional analyses of social position. *British Journal of Ophthalmology* 2017;101(4):395-400.
4. Whitehead M. The concepts and principles of equity and health. *International Journal of Health Services* 1992;22(3):429-45. [published Online First: 1992/01/01]
5. Ramke J, Petkovic J, Welch V, et al. Interventions to improve access to cataract surgical services and their impact on equity in low- and middle-income countries. *Cochrane Database of Systematic Reviews* 2017; Issue 11. Art. No.: CD011307
6. Liu T, Congdon N, Yan X, et al. A randomized, controlled trial of an intervention promoting cataract surgery acceptance in rural China: the Guangzhou Uptake of Surgery Trial (GUSTO). *Investigative Ophthalmology & Visual Science* 2012;53(9):5271-78.
7. Zhang XJ, Liang YB, Liu YP, et al. Implementation of a free cataract surgery program in rural China: a community-based randomized interventional study. *Ophthalmology* 2013;120(2):260-5.
8. Östlin P, Schrecker T, Sadana R, et al. Priorities for research on equity and health: towards an equity-focused health research agenda. *PLoS Medicine* 2011;8(11):e1001115.
9. Rasanathan K, Diaz T. Research on health equity in the SDG era: the urgent need for greater focus on implementation. *International Journal for Equity in Health* 2016;15(1):202.
10. Ivers NM, Grimshaw JM. Reducing research waste with implementation laboratories. *The Lancet* 2016;388(10044):547-48.
11. O'Neill J, Tabish H, Welch V, et al. Applying an equity lens to interventions: using PROGRESS ensures consideration of socially stratifying factors to illuminate inequities in health. *Journal of Clinical Epidemiology* 2014;67(1):56-64.
12. Ramke J, Qureshi B, Gilbert CE. To realize universal eye health we must strengthen implementation research. *Middle East African Journal of Ophthalmology* 2017;24(2):65-66.
13. Peters DH, Adam T, Alonge O, et al. Implementation research: what it is and how to do it. *BMJ* 2013;347:f6753.
14. Rockers PC, Tugwell P, Røttingen J-A, et al. Quasi-Experimental Study Designs Series-Paper 13: Realizing the Full Potential of Quasi-Experiments for Health Research. *Journal of Clinical Epidemiology* 2017;89(2017):106-10.
15. Bärnighausen T, Oldenburg C, Tugwell P, et al. Quasi-experimental study designs series—paper 7: assessing the assumptions. *Journal of Clinical Epidemiology* 2017;89:53-66.
16. Geldsetzer P, Fawzi W. Quasi-experimental study designs series-paper 2: complementary approaches to advancing global health knowledge. *Journal of Clinical Epidemiology* 2017;89(2017):12-16.
17. Oliver K, Innvar S, Lorenc T, et al. A systematic review of barriers to and facilitators of the use of evidence by policymakers. *BMC Health Services Research* 2014;14(1):2.
18. Wagenaar BH, Sherr K, Fernandes Q, et al. Using routine health information systems for well-designed health evaluations in low-and middle-income countries. *Health Policy and Planning* 2015;31(1):129-35.