

## Title

Voices from the frontline: findings from a thematic analysis of a rapid online global survey of maternal and newborn health professionals facing the COVID-19 pandemic

## Authors

Name	Affiliation
Aline Semaan*	Department of Public Health, Institute of Tropical Medicine, Antwerp Belgium Center for Research on Population and Health, Faculty of Health Sciences, American University of Beirut, Beirut, Lebanon
Constance Audet	Department of Public Health, Institute of Tropical Medicine, Antwerp Belgium
Elise Huysmans	Department of Public Health, Institute of Tropical Medicine, Antwerp Belgium
Bosede B Afolabi	Department of Obstetrics and Gynaecology, College of Medicine, University of Lagos, Lagos, Nigeria
Bouchra Assarag	National School of Public Health, Ministry of Health, Morocco
Aduragbemi Banke-Thomas	LSE Health, London School of Economics and Political Sciences, London, United Kingdom
Hannah Blencowe	Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London United Kingdom
Severine Caluwaerts	Department of Public Health, Institute of Tropical Medicine, Antwerp Belgium
Oona M R Campbell	Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London United Kingdom
Francesca L Cavallaro	Institute of Child Health, University College London, London United Kingdom
Leonardo Chavane	Department of Community Health, Faculty of Medicine, Eduardo Mondlane University, Maputo, Mozambique
Louise Tina Day	Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London United Kingdom
Alexandre Delamou	Africa Centre of Excellence for Prevention and Control of Transmissible Diseases (CEA-PCMT), University Gamal Abdel Nasser, Conakry, Guinea
Therese Delvaux	Department of Public Health, Institute of Tropical Medicine, Antwerp Belgium
Wendy Graham	Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London United Kingdom
Giorgia Gon	Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London United Kingdom
Peter Kascak	Trencin University Hospital, Trencin Slovakia
Mitsuaki Matsui	Department of Global Health, Nagasaki University School of Tropical Medicine and Global Health, Nagasaki, Japan
Sarah G Moxon	Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London United Kingdom
Annetee Nakimuli	Department of Obstetrics and Gynaecology, Makerere University and Mulago Specialized Women and Neonatal Hospital, Kampala Uganda
Andrea B Pembe	Department of Obstetrics and Gynaecology, Muhimbili University of Health and Allied Sciences, Dar es Salaam Tanzania
Emma Radovich	Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London United Kingdom
Thomas van den Akker	Department of Obstetrics and Gynaecology, Leiden University Medical Centre and Athena Institute, Vrije Universiteit Amsterdam, Netherlands
Lenka Benova	Department of Public Health, Institute of Tropical Medicine, Antwerp Belgium

\*Corresponding author

## **Abstract**

### **Objective**

To prospectively document experiences of frontline maternal and newborn healthcare providers during the COVID-19 pandemic.

### **Design**

Cross-sectional study via an online survey disseminated through professional networks and social media in 12 languages. We analysed responses using descriptive statistics and qualitative thematic analysis disaggregating by low- and middle-income countries (LMICs) and high-income countries (HICs).

### **Setting**

81 countries, between March 24 and April 10, 2020.

### **Participants**

714 maternal and newborn healthcare providers.

### **Main outcome measures**

Preparedness for and response to COVID-19, experiences of health workers providing care to women and newborns, and adaptations to 17 outpatient and inpatient care processes during the pandemic.

### **Results**

Only one third of respondents received training on COVID-19 from their health facility and nearly all searched for information themselves. Half of respondents in LMICs received updated guidelines for care provision compared with 82% in HICs. Overall, only 47% of participants in LMICs, and 69% in HICs felt mostly or completely knowledgeable in how to care for COVID-19 maternity patients. Facility-level responses to COVID-19 (signage, screening, testing, and isolation rooms) were more common in HICs than LMICs. Globally, 90% of respondents reported somewhat or substantially higher levels of stress. There was a widespread perception of reduced use of routine maternity care services, and of modification in care processes, some of which were not evidence-based.

### **Conclusions**

Substantial knowledge gaps exist in guidance on management of maternity cases with or without COVID-19. Formal information sharing channels for providers must be established and mental health support provided. Surveys of maternity care providers can help track the situation, capture innovations, and support rapid development of effective responses.

## Key Messages

### *What is already known*

- In addition to lack of healthcare worker protection, staffing shortages, heightened risk of nosocomial transmission and decreased healthcare use described in previous infectious disease outbreaks, maternal and newborn care during the COVID-19 pandemic has also been affected by large-scale lockdowns/curfews.
- The two studies assessing the indirect effects of COVID-19 on maternal and child health have used models to estimate mortality impacts.
- Experiences of frontline health professionals providing maternal and newborn care during the COVID-19 pandemic have not been empirically documented to date.

### *What this study adds*

- Respondents in high-income countries more commonly reported available/updated guidelines, access to COVID-19 testing, and dedicated isolation rooms for confirmed/suspected COVID-19 maternity patients.
- Levels of stress increased among health professionals globally, including due to changed working hours, difficulties in reaching health facilities, and staff shortages.
- Healthcare providers were worried about the impact of rapidly changing care practices on health outcomes: reduced access to antenatal care, fewer outpatient visits, shorter length-of-stay in facilities after birth, banning birth companions, separating newborns from COVID-19 positive mothers, and postponing routine immunisations.
- COVID-19 illustrates the susceptibility of maternity care services to emergencies, including by reversing hard-won gains in healthcare utilisation and use of evidence-based practices. These rapid findings can inform countries of the main issues emerging and help develop effective responses.

## Introduction

Coronavirus disease (COVID-19) is a respiratory tract infection caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which was first recognized in December 2019 in Wuhan, China.<sup>1</sup> COVID-19 is a highly infectious disease with two main routes of transmission: directly via close contacts with an infected person and indirectly via contact with contaminated surfaces. While evidence gathering continues, concerns are emerging regarding a possible vertical transmission (antenatally or intrapartum).<sup>2,3</sup> The effect of COVID-19 infection during the 1<sup>st</sup> and 2<sup>nd</sup> trimester of pregnancy remains to be clarified, highlighting the need for a good surveillance system to register adverse outcomes arising from infection in early pregnancy. At present, the virus has not been detected in breast milk of mothers with confirmed (or suspected) COVID-19 infection, so transmission via breastfeeding is then considered unlikely. Direct transmission from mother to child may occur via close contact, but breastfeeding continues to be encouraged with appropriate hygiene measures, including wearing face masks.<sup>4,5</sup> As of May 4<sup>th</sup> 2020, 3.3 million cases of COVID-19 have been confirmed and more than 230,000 deaths reported globally.<sup>6</sup> Based on data collected by the Chinese Centre for Disease Control and Prevention, an infection with COVID-19 can cause a range of illness severity: from mild to moderate (>80%), to severe (14%) and critical (5%). The overall case fatality rate is estimated at 2.3%.<sup>7</sup> Among the risk factors associated with an increased case fatality rate are older age, male sex, and comorbidities, particularly cardiovascular diseases and diabetes.<sup>8</sup>

Studies are currently ongoing to evaluate whether pregnant women have an increased susceptibility to infection with COVID-19 and if they present a greater risk of severe illness or mortality. However, the available limited evidence suggests that pregnant women have risks of infection comparable to the general population.<sup>9</sup> The disease severity in pregnant women does not appear significantly higher than in non-pregnant women.<sup>10,11</sup> A meta-analysis conducted by Di Mascio et al. of 41 pregnant women hospitalised in a context for a COVID-19 infection showed an increased risk of preterm birth, preeclampsia, and caesarean section.<sup>12</sup> Symptoms in newborns suspected or confirmed with COVID-19 seem to be mild, with good outcomes.<sup>13</sup> though one study reported a higher risk of perinatal death.<sup>12</sup> Amoroux et al. draw attention to the possible delay in the development of visible hypoxemic lesions in newborns of COVID-19 positive mothers, and advise their close follow-up after birth.<sup>3</sup> However, the limited size of samples in these studies calls for caution and more data need to be collected to draw definitive conclusions.<sup>14</sup> Considering the increased risk of infection with other respiratory viruses such as influenza, and the increased mortality linked with H1N1, it is important for pregnant women to be protected from illnesses.<sup>12</sup> However, for pregnant women, irrespective of the risks, symptoms, and severity, the main recommendations to avoid infection remain the same as for the general public.<sup>15,16</sup> Some countries, such as the United Kingdom, issued stricter measures for pregnant women, categorising them as part of a vulnerable group applying the precautionary principles and recommending self-isolation.<sup>9</sup>

The effects of COVID-19 are likely to go beyond the direct provision of care to women and newborns with suspected or confirmed COVID-19 infection. Previous infectious disease outbreaks severely reduced the capacity of health systems to provide critical reproductive, maternal, and newborn healthcare, with negative impacts on their health outcomes.<sup>16-18</sup> Studies of recent outbreaks of Ebola virus disease (EBV), SARS and MERS, highlighted several challenges in countries' preparedness to face outbreaks, amplified by a weak existing systems. This includes lack of protection and safety for healthcare workers leading to disruption in staffing, heightened risk of nosocomial transmission, and elevated stress levels among service providers.<sup>19,20</sup> A qualitative study by Qian Liu et al. also shows that healthcare providers in China were stressed during the ongoing outbreak because of an added workload and fear of contracting and transmitting the infection.<sup>21</sup> Other indirect consequences of previous infectious disease outbreaks include less healthcare utilisation and limited capacity for public health surveillance.<sup>22-24</sup> These impacts can persist long after the disease outbreak is contained.<sup>25</sup> However, much of the evidence available about these impacts on maternal and newborn health is either modelled or is from studies using secondary data such as population-based surveys and routine health management information system analysis.<sup>22,26,27</sup> Additionally, large disruption to health-seeking behaviour and healthcare provision is caused by the unprecedented measures countries implement to contain the pandemic (e.g., lockdowns, curfews, restrictions on public transport).

To date, studies assessing the potential indirect effects of the COVID-19 pandemic on sexual, reproductive, maternal and child health have used modelling approaches. Robertson and colleagues modelled three scenarios projecting a decrease in the coverage of basic life-saving interventions, varying the extent and duration.<sup>28</sup> They estimated an increase in maternal deaths between 12,190 and 56,700, and 253,500-1,157,000 additional deaths of children under five years of age. Similar conclusions have been drawn by Riley et al. who projected that a modest decline in the use of sexual and reproductive healthcare services in 132 low- and middle-income countries (LMICs) will result, over the course of a year, in 48 million additional women with unmet need for modern contraceptives, 15 million additional unwanted pregnancies, and over 3 million additional unsafe abortions will occur.<sup>29</sup> It is therefore critical that the precise nature of the direct and indirect impacts of COVID-19, and the adaptations and innovations tested to reduce its impact are captured and described, and that this be done prospectively.<sup>30</sup>

Levels of preparedness and response in maternal and newborn care services differ markedly between institutions and countries. Health personnel, including those defined by the World Health Organization as “competent maternal and newborn health professionals educated, trained and regulated to national and international standards”<sup>31</sup> are at the frontline of providing care for pregnant women and their newborns during the COVID-19 pandemic. They are expected to be sufficiently competent to: (i) provide and promote evidence-based, human-rights based, quality, socio-culturally sensitive and dignified care to women and newborns; (ii) facilitate physiological processes during labour and delivery to ensure a clean and positive childbirth experience; and (iii) identify and manage or refer women and/or newborns with complications. They are the ones that serve as the interface between the governments working to address the pandemic and pregnant women experiencing the consequences, as such they are uniquely placed to be able to describe the status of care provision during such times.

The objective of this paper is to synthesise the key themes identified in the first round of a global online survey of health professionals working in maternal and newborn health along four dimensions: preparedness for COVID-19, response to COVID-19, personal experience in the workplace, and changes in provision of care and care processes. This online survey is part of a larger study which seeks to 1) understand how health professionals and health facilities prepare and respond to COVID-19 in regard to the care provided to women and their babies during antenatal, intrapartum and postnatal care; and 2) document and analyse the effect of the COVID-19 pandemic on the services available to pregnant, labouring and postpartum women and their newborns, including as a result of increasing pressures on the healthcare system.

## **Methods**

### Study design

This is a cross-sectional study of health professionals providing maternal and newborn healthcare services. In the future, we plan to collect repeated rounds of the online survey to track the preparedness for, response to and effects of the COVID-19 pandemic over time and follow-up qualitative individual interviews from selected respondents to gain additional insights.

### Population and sampling

The target population for this survey was health professionals directly providing maternal (antenatal, intrapartum and/or postnatal) or newborn care. We included cadres such as midwives, nurses, obstetricians, gynaecologists, neonatologists, paediatricians, anaesthetists, general practitioners, medical officers, clinical officers, community health workers, lactation counsellors, paramedics, health technicians, and others, including health professionals in training. Due to the unavailability of a global sampling frame for this study population, the survey sampling was non-random and not intended to generate generalisable nationally representative results of either health professionals or health facilities. Rather, our intention was to collect and synthesise the voices and experiences of maternal and newborn health professionals from a range of countries, contexts, services and facility types at the early stage of the COVID-19 pandemic. An invitation to complete the survey was distributed using personal networks of the multi-country research team members, maternal/newborn platforms, and social media (e.g. Facebook, Twitter, WhatsApp). Respondents were encouraged to share the survey with other colleagues in an attempt to snowball the

sample population. Respondents provided informed consent online by checking a box affirming that they understood the consent form and voluntarily agreed to participate in the survey.

### Questionnaire development

A questionnaire was developed by an international team of collaborators including health professionals, experts in health systems, infectious diseases, infection prevention and control, maternal health epidemiologists, and public health researchers from various global settings. The questionnaire was prepared in English, and translated into 11 languages (French, Arabic, Italian, Portuguese, Spanish, Chinese, Japanese, Russian, German, Swahili, and Dutch), by native speakers with medical training (minimum two translators per language). It was piloted by asking five maternal/newborn health professionals from different global settings to complete the questionnaire and provide feedback. We used this feedback to assess face validity and refine the wording of the questions and the format and wording of response options. We collected data on the respondents' background (country and region, qualification and work responsibilities, gender, and basic characteristics of the health facility in which the respondents worked, if any). To avoid concerns over confidentiality, we did not collect names of health facilities. The questionnaire included three core modules focusing on preparedness for COVID-19, response to COVID-19, and health workers' own experience of work during the COVID-19 pandemic. In the fourth, optional module, we asked respondents to elaborate on adaptations to 17 care processes (timing, frequency, modality of contact with patients during various types of outpatient and inpatient care) and to comment on whether they perceived that the uptake of care by the population they serve has changed and, if it had, how. The full questionnaire is provided in Supplementary File 1.

### Data processing and analysis

In this paper, we use responses collected between the day the survey was launched (March 24, 2020) and April 10, 2020. First, we cleaned the 798 responses received by removing duplicate submissions ( $n=49$ ) and those who did not agree to the consent statement ( $n=14$ ), and submissions made by those not directly providing maternal or newborn care (such as lecturers and public health officials;  $n=10$ ). Quantitative analysis involved production of descriptive statistics including frequencies and percentages using Stata/SE version 14, and responses were stratified by country income levels (according to World Bank classification).<sup>32</sup> We conducted a qualitative thematic analysis of free-text answers to derive common themes related to respondents' concerns and reported changes in the work environment and care process by country income levels. When possible, we triangulated qualitative and quantitative results to validate emerging themes.

### Missing data

From the 725 remaining responses, we dropped 11 responses with missing answers on more than 90% of the survey questions from analysis. The proportion of missing answers to multiple choice questions ranged between 0.5 and 6.5%, and that to open-ended questions from 16 to 28% of respondents. Missing answers to the "Country" question were recorded based on the "Region" answer for 93 responses; for example, a respondent with a missing response for country but region reported as Maharashtra was coded as from India.

## **Results**

### Respondents' characteristics

The analysed sample included a total of 714 healthcare professionals caring for women and newborns, 59% of whom agreed to participate in the optional module of the survey ( $n=397$ ). Table 1 summarises respondents' characteristics. Participants were based in 81 countries and more than half (63%) were from high-income countries (HICs). A map showing respondents' geographic distribution is available in Supplementary File 2. Obstetricians/gynaecologists and midwives constituted the majority of respondents (38% and 35%, respectively), followed by nurse-midwives and nurses. Around one third worked in referral hospitals and 60% were employed in public sector facilities. Most facilities where participants worked provided caesarean sections (81%), accepted referrals from other facilities (71%), and included maternal intensive care units (ICU, 64%) and newborn intensive care units (NICU, 59%). Nearly half of respondents from HICs (49%) reported that their facilities had seen maternity patients with confirmed or

suspected COVID-19 infection, compared to 13% of respondents from low-and-middle income countries (LMICs).

Table 1 - Survey (n=714\*) and optional module (n=397) respondent characteristics

	Survey (%)	Optional module (%)
<b>Country income level (World Bank classification)</b>		
Low and Middle income countries	263 (37)	136 (35)
High income countries	444 (63)	256 (65)
<b>Region</b>		
East Asia & Pacific	82 (12)	51 (13)
Europe & Central Asia	249 (35)	131 (33)
Latin America & Caribbean	43 (6)	30 (8)
Middle East & North Africa	53 (7)	29 (7)
North America	87 (12)	53 (14)
South Asia	83 (12)	37 (9)
Sub-Saharan Africa	110 (16)	61 (16)
<b>Cadre</b>		
Midwife	248 (35)	135 (34)
Nurse-midwife	83 (12)	48 (12)
Nurse	22 (3)	14 (4)
Obstetrician/gynaecologist	269 (38)	148 (38)
Neonatologist	6 (1)	3 (1)
Paediatrician	4 (1)	4 (1)
General practitioner	10 (1)	5 (1)
Medical doctor (no specialization)	15 (2)	10 (3)
Medical student/intern/resident	13 (2)	6 (2)
Community health worker/Outreach worker	12 (2)	6 (2)
Other	29 (4)	16 (4)
<b>Position</b>		
Head of facility	60 (9)	34 (9)
Head of department or ward	71 (10)	41 (11)
Head of team	94 (13)	54 (14)
Team member	346 (50)	195 (50)
Locum or interim member	22 (3)	10 (3)
Other <sup>a</sup>	101 (15)	53 (13)
<b>Type of care provided (multiple responses allowed)</b>		
Outpatient ANC	438 (61)	244 (62)
Home-based childbirth care	77 (11)	47 (12)
Outpatient PNC	316 (44)	176 (45)
Outpatient Breastfeeding support	217 (30)	121 (31)
Inpatient ANC	374 (52)	218 (56)
Inpatient childbirth care	437 (61)	249 (64)
Inpatient PNC	350 (49)	193 (50)
Surgical care	213 (30)	115 (29)
Neonatal care (small and sick newborns)	85 (12)	47 (12)
Home visits	131 (18)	78 (20)
Community outreach	105 (15)	69 (18)
Abortion care	157 (22)	86 (22)
Post-abortion care	179 (25)	104 (27)
Other	84 (12)	42 (11)
<b>Health facility level</b>		
Referral hospital	250 (36)	144 (37)
District/regional hospital	154 (22)	77 (20)

Health center	76 (11)	46 (12)
Polyclinic	6 (1)	6 (2)
Clinic	66 (10)	36 (9)
Health post/unit or dispensary	16 (2)	9 (2)
Other <sup>b</sup>	116 (17)	68 (18)
<b>Health facility sector</b>		
Public (national)	183 (27)	86 (22)
Public (university or teaching)	138 (20)	81 (21)
Public (district level or below)	80 (12)	61 (16)
Social security	7 (1)	3 (1)
Health insurance or HMO	10 (2)	7 (2)
Private university	25 (4)	10 (3)
Private for profit	95 (14)	57 (15)
Non-governmental	61 (9)	29 (8)
Faith-based or mission	23 (3)	15 (4)
Other	56 (8)	34 (9)
<b>Type of area</b>		
Large city (more than 1 million inhabitants)	273 (40)	151 (39)
Small city (100,000 to 1 million inhabitants)	220 (32)	125 (32)
Town (fewer than 100,000 inhabitants)	106 (16)	61 (16)
Village/Rural area	64 (9)	38 (10)
Refugee/displaced persons camp	8 (1)	2 (1)
Other	9 (1)	9 (2)
<b>Facility characteristics</b>		
Caesarean-section provision	535 (81)	301 (81)
Accept referrals from other facilities	476 (71)	269 (71)
ICU available	429 (64)	236 (62)
NICU available	398 (59)	226 (59)

\*Differential number of missing values across variables

<sup>a</sup> Mainly self-practicing midwives

<sup>b</sup> Mainly birth centres and private practice

Abbreviations: Antenatal care (ANC); Intensive care unit (ICU); Neonatal intensive care unit (NICU); Postnatal care (PNC)

### Knowledge on the provision of maternal and newborn healthcare and COVID-19

Most respondents (90%) reported that the health facilities where they work provided them with information on preparing for the COVID-19 outbreak (Table 2). This included general guidance about the disease (definition, transmission mode and treatment options), prevention measures (e.g. hand hygiene, disinfecting surfaces and equipment, personal protective equipment [PPE] use, social distancing and isolation), patient screening, case reporting, and updated policies and guidelines. However, only one third of respondents reported receiving hands-on training/drills on the response to COVID-19 (Table 2). Several raised concerns about this lack of access to training activities and perceived this as a necessity that would have made them “*feel better prepared*” to respond to the needs of women and their babies during the outbreak.

Among respondents from LMICs (n=263), half reported receiving updated guidelines for the provision of maternal and newborn care reflecting measures for the COVID-19 outbreak, compared to 82% of those from HICs (Table 2). Some LMIC-based respondents (particularly from Tanzania, Rwanda, Uganda, and India) expressed their concern over the lack of updated guidelines and protocols. An obstetrician/gynaecologist from Uganda remarked: “*I am worried that no national guidelines [are] rolled out yet in regards to care for pregnant women and newborns.*” Some midwives working in HICs requested clearer guidelines on the provision of midwifery care during home visits. Nearly all respondents reported having searched personally for information on COVID-19 (92%), and received informal guidance from colleagues (90%, Table 2). Nonetheless, some participants in LMICs were worried about the lack of access to/availability of evidence on the effects of COVID-19 during pregnancy and the possibility of in-utero transmission, and transmission through breast milk to newborns. More than half (61%) of respondents

perceived that patients' questions regarding COVID-19 were being adequately answered by healthcare providers in their respective facilities. However, only 19% of participants felt that they were completely knowledgeable of the measures that should be taken to provide care to COVID-19 maternity patients (Table 2). Almost half of respondents indicated that their facilities had shared materials with maternity patients on COVID-19, and these materials were mainly disseminated through health facility websites, leaflets/fliers, posters, and on social media.

Table 2 – Preparedness for COVID-19 among maternal and newborn health professionals, by country income category

	LMIC N=263 (%)	HIC N=444 (%)	Total* N=714 (%)
Institution provided information on how to prepare for COVID-19	217 (86)	403 (93)	620 (90)
Institution provided training on COVID-19	94 (37)	143 (34)	237 (35)
Received updated guidelines for MNH care provision because of COVID-19	118 (47)	347 (82)	465 (69)
Personally searched for guidance and information to prepare for COVID-19	242 (95)	387 (90)	629 (92)
Received information related to COVID-19 informally through colleagues	230 (91)	386 (89)	616 (90)
Facility published materials covering COVID-19 targeted toward pregnant, labouring, or postnatal women	98 (38)	194 (46)	292 (43)
Perception that patients' questions were adequately answered at facility	145 (57)	267 (64)	412 (61)
<b>Level of knowing how to provide care for a woman with COVID-19</b>			
Not at all clear	16 (6)	6 (1)	22 (3)
Some points clear, but not confident in what to do	59 (24)	43 (10)	102 (15)
Somewhat clear but major issues remain	56 (23)	84 (20)	140 (21)
Mostly clear but some areas of concern remain	80 (32)	203 (48)	283 (42)
Very clear	38 (15)	90 (21)	128 (19)

\*Differential number of missing values across variables

Abbreviations: High income countries (HIC); Low and middle income countries (LMIC)

Some participants from LMICs such as India, Bangladesh, Bolivia, and Syria expressed concerns regarding patients' degree of application of instructions, particularly those related to social/physical distancing and hygiene measures. An obstetrician/gynaecologist from India mentioned worrying about "*patients and relatives not following instructions given by staff members*". As described by a nurse from Syria, some respondents attributed this to a "*lack of awareness and knowledge, and indifference among beneficiaries*". Furthermore, the implication of the local communities and sharing responsibilities in terms of application of the hygiene and social/physical distancing measures were mentioned by some respondents. A midwife from Bolivia worried "*that not enough is being done on a personal level by patients to keep themselves safe*".

#### Work environment adaptations in response to COVID-19

Three quarters of participants from HICs reported that their facilities had set up a well sign-posted general entrance and screening area for COVID-19 suspected cases, compared to 37% of respondents from LMICs. Among HIC respondents, 83% reported that their facilities reserved isolation rooms for suspected COVID-19 cases, compared to 57% of LMIC respondents (Table 3). The majority of respondents (62%) reported that their facilities have designated a COVID-19 liaison person or team. These teams were most commonly assigned at the level of the facility (56%), followed by both at the levels of facility and maternity ward (27%), and in the maternity ward alone (17%). Screening for COVID-19 symptoms among maternity patients was also more commonly reported by respondents working in HICs (76%) versus 47% in LMICs. For example, antenatal care (ANC) patients - both outpatients and inpatients - were screened either in person or over the phone before scheduling appointments. The ability to order COVID-19 tests for maternity patients was available for 61% of respondents in HICs; but limited in LMICs (23%), rural areas (9% in LMICs

and 28% in HICs) and completely unavailable to respondents working in refugee and/or displaced persons camps (n=6, data not shown).

Table 3 - Response to COVID-19 among maternal and newborn health professionals and their workplaces, by country income category

	LMIC N=263 (%)	HIC N=444 (%)	Total* N=714 (%)
<b>Sign-posted area for screening of COVID-19 suspected cases in facility</b>			
No	38 (15)	32 (8)	70 (11)
Some measures taken	106 (42)	65 (16)	171 (26)
Yes	95 (37)	298 (72)	393 (59)
<b>Reserved isolation rooms for suspected cases</b>	143 (57)	341 (83)	484 (73)
<b>Screening for COVID-19 symptoms among maternity patients</b>	117 (47)	320 (76)	437 (75)
<b>Possible to order a test for COVID-19 for maternity patients</b>	58 (23)	258 (61)	316 (47)
<b>Sufficient PPE items</b>			
Gloves	174 (70)	399 (92)	578 (84)
Masks	117 (47)	224 (52)	345 (50)
Aprons	88 (36)	260 (61)	352 (52)
All three types	79 (32)	188 (44)	267 (40)
<b>Respondents' work affected by COVID-19</b>	177 (71)	372 (86)	549 (81)
<b>Respondents' stress levels</b>			
Same as usual	21 (8)	47 (11)	68 (10)
Somewhat higher than usual	136 (54)	215 (50)	351 (52)
Substantially higher than usual	93 (37)	167 (39)	260 (38)

\*Differential number of missing values by variables

Abbreviations: High income countries (HIC); Low and middle income countries (LMIC); Personal Protective Equipment (PPE)

The lack of COVID-19 symptom screening and the inability to order tests constituted major concerns for respondents, who perceived these deficiencies as threats to the safety of the workforce and of other patients. A midwife from Canada wrote, *"I'm worried about being infected by someone who is asymptomatic, and then being a vector to others."* Compromising patients' and healthcare providers' safety was also viewed as stemming from deficiencies in availability of PPE, including face masks, gloves, and aprons. These findings were consistent across all settings but more prominent in LMICs (Table 3). A midwife from the United Kingdom pleaded, *"Let midwives who are in close contact with women wear masks. [...] Please let us use masks for all."* Additionally, respondents advocated for clear guidelines and unified protocols regarding the appropriate use of PPE. For example, a nurse-midwife from the United States wrote, *"[...] as of now we are not allowed to wear masks and goggles unless delivering a patient, we're told to « take the mask off or go home » that we're scaring the patients."* Despite the need to feel protected, wearing additional PPE can be burdensome. It was described as time-consuming and respondents worried about resulting delays in the provision of emergency care because of having to don and doff PPE. Additionally, they were concerned that PPE might reduce their ability to communicate clearly with patients, such as a midwife from Denmark who remarked that *"[i]t can be hard to connect with people through masks and [goggles] (facial expressions are harder to read)."*

The majority (81%) of respondents noted that their work had been affected by the COVID-19 outbreak and that their stress levels were either somewhat or substantially higher than usual (90%, Table 3). In the words of an obstetrician from Mozambique: *"My stress level at this point is immeasurable. Every time a pregnant woman with flu-like symptoms [visits the health facility], I feel almost completely lost and I end up only of [thinking about this] patient. I need to be equally protected and I don't feel any protection from whoever [is responsible for protecting me]."* A major challenge reported was the decrease in skill mix and shortages of qualified staff, either because of symptoms, self-isolation after potential exposure, or not being able to get to their workplace due to lockdowns and transport restrictions as described by a midwife in Uganda: *"[t]ransport to work is a big challenge due to lockdown; many staff live far away from the hospital. The staff who manage to come to work hurry to leave the hospital early to observe the curfew time of 7.00 p.m."* This

shortage has led to an increase in the workload and unexpected changes in work schedules. Anxiety and exhaustion levels have increased because of these rapid changes, and some respondents expressed the need for more support from management. In certain contexts, healthcare facilities were increasingly relying on locum workers and students to fill staffing shortages. An obstetrician/gynaecologist who headed a department in Uganda reported that “[t]here is no more clear work schedules as I get to attend many unscheduled/emergency meetings [...]. The staff are very anxious and panicky and need talking to all the time, which is exhausting.”

#### Changes to the care provided to women and newborns

We analysed responses from 397 health professionals who completed the optional module. Figure 1 shows the main reported changes in service provision and utilisation, care content and quality, and care process adaptations across the continuum of maternal and newborn care. In all settings and across the continuum of care, participants reported seeing fewer patients at healthcare facilities. This was described as the result of transportation difficulties in accessing health facilities or due to women’s fear of contracting COVID-19 at the facility. A nurse-midwife from Kenya wrote, “[a]ccessing inpatient antenatal care [is] minimal. Women fear to [get] infected with COVID-19 if [they are present] in hospitals. Most of them keep off from hospital even when they are sick”. Most respondents noted that their facilities have shortened visiting hours and reduced the number of visitors allowed, while others are screening visitors for symptoms, or have banned visits altogether. Importantly for the support of women during labour and childbirth, facilities are reportedly limiting the number of labour companions to one person designated as the single visitor allowed to stay with the mother after birth or banning birth companions altogether. This raised concerns among healthcare providers regarding the reduced support available to women, and increased workload on the staff. An obstetrician from the Czech Republic remarked that: “[the] Gynaecological and Obstetrical Society has recommended to ban partners and doulas from accompanying a woman at birth - outrageous!!!”

Some healthcare facilities were implementing social/physical distancing measures in the waiting areas of the outpatient departments, and in hospital rooms by reducing the number of beds. Yet, this recommendation is challenging to meet in facilities where resources are limited – an obstetrician/gynaecologist from India noted that “[it is] not practically possible [to place each patient in a separate birthing room] in our set up”. Non-essential services including elective gynaecological procedures and infertility treatments were being postponed or cancelled. In several settings, responding to the COVID-19 outbreak affected the delivery of routine ANC, which became restricted to the management of high-risk patients. A respondent from New York reported a “significant decrease in number of ANC visits”, whereby new policies recommend the reduction of the number of face-to-face visits during pregnancy “from 10-12 to four”. Other changes include eliminating the waiting area, spacing face-to-face appointments to reduce contact between patients, and cancelling all group activities such as health education sessions or group counselling.

Respondents also reported a shift to telemedicine for the provision of both antenatal and postnatal care (PNC), including breastfeeding counselling. Although telemedicine was considered a priority in certain LMICs where it is not implemented yet, participants acknowledged the challenges associated with this service provision modality. This includes lack of access to adequate communication infrastructure among women. Respondents from both LMICs and HICs noted that the demand for home births has increased and that new practices aimed to reduce induction of labour. In certain HICs, induction of labour was reported to be discouraged before 41 weeks of gestation. Changes in pain relief options for labouring women in HICs included decrease in the use of nitrous oxide to reduce the risk of infection transmission through aerosols, and suspended waterbirths. Across all settings, caesarean sections were reported as a commonly performed procedure among women who were diagnosed with COVID-19. Some respondents noted that their facilities have dedicated operating theatres specifically for this purpose. On the other hand, the numbers of elective caesarean sections have reportedly decreased among “healthy” maternity patients. However, this was not consistent in facilities where certain efforts were made to reduce the duration of labour and the time spent in the labour room by augmentation. As a result, respondents speculated about a potential rise in caesarean section rates in their facilities, as noted by an obstetrician/gynaecologist from India: “We will not allow as much time in second stage [of labour], [...] this is likely to push up our caesarean rate.”

Respondents frequently mentioned shortened length of stay in facilities after childbirth; for example a reduction “to 6-8 hours from 24 [or more hours]” (midwife from Canada). This was worrisome for some respondents as noted by a midwife from the UK: “[the] lack of time and staff will lead to mothers and babies going home with very little feeding support or knowledge which will have a short and long term impact on their health and ability to deal with infections.” Routine postnatal checks are being postponed in certain cases or substituted with telemedicine. A nurse-midwife from the United States reported that “[w]e are postponing the routine postpartum visit until 12 weeks postpartum, and are prescribing most contraceptives over the phone or [...] and breastfeeding support is all done virtually.” Changes to newborn postnatal care were infrequently reported, and mainly included monitoring and isolation of babies of mothers with confirmed COVID-19. Three respondents from India noted that the infant vaccination schedule was disrupted or postponed. Overall, respondents expressed their concern over the uncertain impact of reduced face-to-face interactions on the quality of care. A midwife from the UK wrote: “[w]hilst I completely see the need to restrict our face-to-face care to protect staff and patients, my heart just breaks for women and families who we won’t be able to offer the full range of midwifery support to... i.e. BF support, daily visits, and just generally our time”. Maternal and newborn health professionals feared that the changes to the standards of care would lead to poor health outcomes among women and newborns and, subsequently to the loss of progress achieved in certain indicators (e.g., stillbirth rates). “I am also worried about the implications of the policies that call for separating newborns from COVID-19 positive mothers immediately after birth, without allowing for skin-to-skin or delayed cord clamping,” wrote a nurse-midwife from the United States.

## Discussion

This paper uses a rapid collection of data from health professionals providing care to women and their babies globally. We describe preparedness for COVID-19, response to COVID-19, personal experience in the workplace, and changes in provision of care and care processes.

**Preparedness.** We found that respondents actively sought information related to COVID-19 through personal searches and existing informal networks. Studies show that healthcare providers commonly resort to such sources to fulfill information needs.<sup>33</sup> Knowledge gaps were generally related to the impact of COVID-19 on pregnancy and health outcomes for the mother and newborn, or to guidance on the management of COVID-19 maternity cases. There is a high possibility that unreliable information related to the outbreak might be accessed, particularly on social media platforms.<sup>34</sup> Facility-specific creation and distribution of guidelines for managing maternity patients is somewhat lagging behind despite frequent general updates published by Ministries of Health and professional associations.<sup>35–41</sup> Information sharing channels must be established to secure providers’ timely access to accurate information that empowers them to respond to patients’ needs.<sup>42–44</sup> Midwives supporting pregnant and labouring women during the pandemic,<sup>45,46</sup> and particularly those who are practicing independently, have voiced the need to access clear guidelines for providing care during home visits.<sup>47</sup>

**Response.** Our results highlight variability in the facility-level response to COVID-19 between HICs and LMICs, including sharp differences in updating guidelines, setting-up signage and patient/visitor screening, testing availability, and dedicating isolation rooms for maternity patients with confirmed or suspected COVID-19. These discrepancies could stem from the differential progression of the outbreak (whereby more respondents from HICs reported having provided care to COVID-19 confirmed or suspected maternity cases than those from LMICs). These differences could also be partly attributed to the limited capacities and resources of healthcare systems in some LMICs.<sup>48</sup> There is speculation that the outbreak in African countries might be attenuated, but equally possible that trends similar to those witnessed in Europe might be observed.<sup>49,50</sup> This indicates an urgent need to mobilise resources in resource-limited settings, improve testing capacities, and upgrade the responses, including at maternity facilities. The total absence of testing for suspected patients in refugee and/or displaced persons camps reported by all respondents with such experience raises concerns. Living conditions in these under-served settlements, such as overcrowding and lack of adequate water and sanitation, make the implementation of basic infection prevention and control measures nearly impossible.<sup>50–53</sup> Displaced women and their newborns face sub-optimal access to ANC, skilled attendance at birth, PNC, and vaccination, and subsequently experienced poor health outcomes even prior to the pandemic-induced disruptions of essential services.<sup>54–57</sup> Global and local efforts

must be established to ensure that displaced populations have access to appropriate infection prevention measures, testing and treatment, and to quality maternal and newborn services to halt anticipated exacerbations of negative health outcomes.<sup>29,52</sup>

**Personal Experiences.** Consistently with experiences from previous infectious disease outbreaks and emergencies, healthcare workers providing essential services to women and newborns during this pandemic experience increased levels of stress and anxiety.<sup>20,44</sup> Stress levels in LMICs were comparable to those in HICs even though countries were battling different stages of the outbreak. This might be due to uniformly reported changes in working hours, inability to reach health facilities, and shortages in skilled workforce (some of which were attributable to lockdowns and other blanket measures to combat the spread of COVID-19) leading to higher workloads, which can lead to staff burnout.<sup>48,58</sup> Wilson et al. compiled a list of measures that could prevent burnout among maternity care providers.<sup>44</sup> Other lessons learned from past epidemics include the provision of emotional, social and mental health support to care providers, and ensuring that adequate levels of support is available to them from facility management.<sup>20,59</sup> As our findings show, this can create additional burdens to management and special efforts should be placed to provide support to this group.<sup>44</sup> With the increasing reliance on students and trainees to compensate for staff shortage, Wilson et al. consider this group to be more exposed to stressors considering their lack of experience, and therefore senior colleagues should actively advocate for their wellbeing.<sup>44</sup> Future research should explore the availability of mental and social support to maternal and newborn healthcare providers during the pandemic, and its effectiveness.<sup>60</sup>

Another cause for increased stress levels among providers is the fear for their own/their relatives' safety, in addition to the safety of their patients, which is intensified by inadequate access to PPE. One reason for that is that in some facilities, PPE supplies are prioritised for departments treating COVID-19 cases and not reaching maternity wards, which is common in vulnerable settings. Workers who provide essential maternity care and their patients, could thus experience uneven risks of nosocomial infection during outbreaks.<sup>61,62</sup> In some countries, obstetricians/gynaecologists commonly work in multiple facilities across the public and private sectors, and their risk of exposure might be exacerbated by the higher number of contacts (other healthcare workers and patients) they experience in this dual practice.<sup>63</sup> Although PPE items are essential, the WHO issued guidelines that promote their rational use given universal shortage.<sup>64</sup> The application of these guidelines must be unified within healthcare facilities, and clearly communicated to maternity and newborn healthcare providers and explained to patients.<sup>44</sup> Health workers caring for women around the time of birth might be used to wearing PPE, however it can make them feel dehumanized, and the donning and doffing PPE is time-consuming and might delay the provision of emergency services.<sup>20,21</sup>

Healthcare providers also worry about the consequences of rapidly changing practices and the uncertainty of their impact on health outcomes. The perceived changes in healthcare seeking behaviours include fewer visits to the healthcare facilities, shortened lengths-of-stay after childbirth, less access to adequate ANC, and in certain cases, disrupted immunisation schedules. Our findings support the narratives told by healthcare providers regarding the discontinuity of basic services,<sup>65</sup> and align with disruptions witnessed during previous infectious disease outbreaks.<sup>58,66–68</sup> These disruptions have previously led to an increase in maternal and neonatal mortality,<sup>22,69</sup> and currently there are signs of similar trends reported from two maternity hospitals in Uganda.<sup>70</sup> Our knowledge of the impact of these changes is restricted to predictions resulting from modelling which strongly suggest a threat to achieved improvements in LMICs.<sup>28,29</sup> The actual impact is yet to be quantified<sup>28</sup> and the effect of these changes in HICs remains unclear. Prioritised measures depending on contextual needs must be put in place to mitigate these indirect consequences of the pandemic.<sup>28,30</sup>

Although some of the changes to care content and process are consistent with the updated guidelines on essential care provision,<sup>40,71–73</sup> other modifications diverge from available evidence, and could ultimately reverse achieved progress if proper action is not taken. These include the elimination of birth companions altogether,<sup>73,74</sup> reducing or banning visitors to maternity wards, performing caesarean section on all COVID-19 positive women,<sup>74,75</sup> augmenting labour or performing unindicated caesarean sections to gain control over timing of deliveries, separating newborns from COVID-19 positive mothers including not allowing breastfeeding,<sup>76</sup> and drastically reducing length-of-stay after facility birth with fewer home visit follow-ups.<sup>47</sup>

These practices could deny women's access to quality, respectful intrapartum and postpartum care, and jeopardize their wellbeing and that of their babies.<sup>77</sup> Unlike curative health services, maternity care focuses on providing holistic support to women going through a normal physiological process; both over and under-intervention can result in massive preventable burden. Another alarming adaptation to COVID-19 is freezing or postponing routine immunisation schedules.<sup>78</sup> Temporary disruptions to routine immunisation were only reported in India in our survey, however other LMICs have implemented similar measures.<sup>78</sup> This can result in overall declines in population coverage, and catch-up campaigns should be prioritised following the relaxation of preventive measures to ensure the sustainability of achievements.<sup>78,79</sup> The introduction of new models of care such as telehealth guidance<sup>29,71</sup> was described as a 'virtually perfect solution' to ensure sustained care provision during the pandemic.<sup>80</sup> Yet this model's feasibility is not universal to all healthcare services. Midwives dread this mode's disturbance of the quality of provided care,<sup>47</sup> and providers in LMICs consider this as an added barrier to achieving equitable access to essential services for women and families who lack the needed resources.

As several qualitative and ethnographic studies have shown,<sup>81,82</sup> healthcare seeking behaviours of patients and local communities relies, among others, on the provider-patient relationship and common cultural, economic and social understanding of health and hygiene.<sup>83,84</sup> Moreover, hierarchical issues such as racial and social discrimination may have a significant impact on the quality of maternal and newborn healthcare, as it has already been highlighted in, for example, West-African urban areas and Malagasy hospitals<sup>81,82,85</sup>, to only cite a few. Dynamics of mutual incomprehension between patients and providers about the attitudes toward and measures necessitated by the outbreak may be taken into account regarding the impact of COVID-19 on maternal and newborn healthcare; especially (but not only) concerning regions and countries where there is a strong medical and cultural pluralism towards the healthcare seeking behaviours. Although only a few respondents mentioned some resistance of patients and local communities to increased hygiene and physical distancing measures in facilities, we know from previous outbreaks such as Ebola that understanding social and cultural responses to the epidemics is essential to prevent healthcare disasters.<sup>86</sup> Furthermore, many respondents addressed the reduction of the number of visitors allowed during labor and childbirth, when such measures do not seem justified. Limiting social support during maternal and neonatal care can put mother and newborn at risk, more particularly in healthcare facilities with an ordinarily high lack of staff and irregular drug delivery where families and surrounding play a crucial role in limiting the impact of these shortcomings.<sup>82,87</sup> A local understanding of healthcare seeking behaviors and social maternal health organization must take place to avoid a top-down management of the outbreak guidelines that may miss the mark of these local pre-existing factors.<sup>88</sup>

## Limitations

We acknowledge that lack of representativeness and related sample bias are limitations of this sampling approach. Our sample might over-represent higher qualified cadres of health professionals in settings with limited use of technology among lower cadres of staff, and under-represent staff who are most overstretched in providing healthcare, or those with limited or no access to internet connection, as we received few responses from professionals working in lower-level facilities, particularly in LMICs. The representativeness of the sample is affected by the availability of the survey in three languages (English, French, and Arabic) for a longer time than the remaining nine languages. Additionally, some cadres, such as neonatologists and paediatricians, were less represented. The questionnaire asks about facilities where respondents work, which is not relevant to independently practicing professionals, especially midwives; this might have discouraged some of them from completing the survey. Finally, data were collected across countries that were going through different stages of the outbreak. As previously mentioned, this could account for country-level discrepancies in the response, and some of the differences seen between HIC and LMIC respondents. We intend to address some of these limitations in survey rounds.


## Conclusion

Despite the limitations, this is the first study attempting to describe the preparedness for, response to and effect of, the COVID-19 pandemic on the provision of maternal and newborn care. The multi-country aspect of the survey allows for a one-stop platform for lessons to be learnt and shared across systems. Our findings, ideally combined with an understanding of women's perspectives, hold enormous potential for creating a timely and evidence-based decision-making platform. Disseminating health workers' voices to

planners, programmers and policymakers is crucial to guide the development of global and contextual guidelines for practice and preparedness.

The COVID-19 pandemic illustrates the worldwide susceptibility to emergencies, which is not restricted to healthcare systems in LMICs. Responding to this crisis is proving to be challenging for health systems and providers, and affects access to basic health services across the globe. Preparedness for the global pandemic might have been equally inadequate for health systems in LMICs and HICs in some aspects, such as shortage in skilled staff, providing training and simulations, and PPE sufficiency. However, it is likely that HICs were able to respond more effectively, due to better health system resilience such as existing coordination systems to develop and implement changes to protocols.<sup>89</sup> Findings from this study will be useful in supporting the development of effective responses to the main issues identified, both during the COVID-19 pandemic and more broadly during future health system shocks.

Figure 1. Reported changes to service provision across the continuum of maternal and newborn care



Items in **bold** were commonly mentioned by respondents from a range of settings, including LMICs and HICs. \*Reported in high income countries. ^Reported in low- and middle-income countries.

## **Acknowledgments**

We would like to thank the study participants who took time to respond to this survey despite the difficult circumstances and increased workload. We acknowledge the Institutional Review Committee at the Institute of Tropical Medicine for providing helpful suggestions on this study protocol, and for the expedited review of this study. We thank all study collaborators and colleagues who distributed the invitation for this survey and provided suggestions on the questionnaire, including the co-authors of this paper, Dr Susannah Woodd and Dr Jean-Paul Dossou. We are immensely grateful to all those who volunteered to translate the survey, including Dr Francesca L Cavallaro, Dr Bouchra Assarag, Dr Karima Khalil, Dr Jose Penalvo, Dr Sonia Menon, Dr Claudia Nieto Sanchez, Virginia Castellano Pleguezelo, Dr Werner Soors, Verónica Blanco Gutiérrez, Dr Raffaella Ravinetto, Dr Manuela Straneo, Francesca Palestra, Dr Stacy Saha, Dr Shirajum Munira, Dr Mitsuaki Matsui, Wnurinham Silva, Dr Leonardo Chavane, Dr Enny Cruz, Min Dai, Dr Li Na, Dr Marc Snel, Dr Anna Larionova, Dr Vladimir Gordeev, Dr Christiane Schwarz, Evelyn Lesta, Elibariki Mkumbo, Dr Amani Kikula, Dr Elly Mertens, and Hilde Cortier.

## **Author contribution**

LB conceptualised the study and obtained funding. All authors contributed to the design of the study and development of the survey tool. AS analysed the data. CA, LB, EH and AS wrote the original draft of the manuscript. All authors contributed to the development of the manuscript, and read and approved the final version. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted. AS is the guarantor.

## **Funding**

This study was funded by the Institute of Tropical Medicine's COVID-19 Pump Priming fund supported by the Flemish Government, Science & Innovation. LB is funded in part by the Research Foundation - Flanders (FWO) as part of her Senior Postdoctoral Fellowship. The funders had no role in the study design, data collection, analysis, and interpretation of data or in writing the manuscript. Researchers are independent from funders and all authors, external and internal, had full access to all of the data (including statistical reports and tables) in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis is also required.

## **Competing interests**

All authors have completed the Unified Competing Interest form (available on request from the corresponding author) and declare: no support from any organisation for the submitted work, no financial relationships with any organisations that might have an interest in the submitted work in the previous three years, and no other relationships or activities that could appear to have influenced the submitted work.

## **Data sharing**

Anonymised data analysed during the current study will be made available from the corresponding author upon reasonable request.

## **Ethical approval**

This study was approved by the Institutional Review Committee at the Institute of Tropical Medicine (Antwerp, Belgium) on March 20, 2020 (approval reference 1372/20).

## **Patient and Public Involvement**

No patient or public involvement took place in the design or conduct of this study. We involved maternal and newborn health professionals, experts in health systems, infectious diseases, infection prevention and control, and maternal health epidemiologists, and public health researchers from various global settings in the design of this study and the survey tool.

**Dissemination to participants and related communities:**

The authors intend to disseminate this research through social media, press releases, and media departments and websites of authors' institutions.

**Transparency**

The guarantor affirms that this manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

## References

- 1 World Health Organization. Naming the coronavirus disease (COVID-19) and the virus that causes it. 2020. [https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-\(covid-2019\)-and-the-virus-that-causes-it](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it)
- 2 Zeng H, Xu C, Fan J, *et al.* Antibodies in Infants Born to Mothers With COVID-19 Pneumonia. *JAMA* Published Online First: March 2020. doi:10.1001/jama.2020.4861
- 3 Amoroux A, Attie-Bitach T, Martinovic J. Evidence for and against vertical transmission for SARS-CoV-2 (COVID-19). 2020;**2**:1–6.
- 4 World Health Organization. FAQ Breastfeeding and COVID-19. 2020. [https://www.who.int/docs/default-source/maternal-health/faqs-breastfeeding-and-covid-19.pdf?sfvrsn=d839e6c0\\_1](https://www.who.int/docs/default-source/maternal-health/faqs-breastfeeding-and-covid-19.pdf?sfvrsn=d839e6c0_1)
- 5 Salvatori G, Rose DU De, Concato C, *et al.* Managing COVID-19-Positive Maternal–Infant Dyads: An Italian Experience. 2020;**15**:3–4. doi:10.1089/bfm.2020.0095
- 6 World Health Organization. Coronavirus Disease (COVID-19) - situation report 104. 2020;;1–7. [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200503-covid-19-sitrep-104.pdf?sfvrsn=53328f46\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200503-covid-19-sitrep-104.pdf?sfvrsn=53328f46_2)
- 7 Wu Z, McGoogan JM. Characteristics of and Important Lessons from the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases from the Chinese Center for Disease Control and Prevention. *JAMA - J Am Med Assoc* 2020;**2019**. doi:10.1001/jama.2020.2648
- 8 Feng Z, Li Q. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *China CDC Wkly* 2020;**41**:145–51. doi:10.3760/cma.j.issn.0254-6450.2020.02.003
- 9 RCOG. Coronavirus ( COVID-19 ) Infection in Pregnancy. Version 8 - 17 April 2020. 2020. <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/coronavirus-pregnancy/>
- 10 Breslin N, Baptiste C, Gyamfi-Bannerman C, *et al.* COVID-19 infection among asymptomatic and symptomatic pregnant women: Two weeks of confirmed presentations to an affiliated pair of New York City hospitals. *Am J Obstet Gynecol MFM* 2020;;100118. doi:10.1016/j.ajogmf.2020.100118
- 11 Yu N, Li W, Kang Q, *et al.* Clinical features and obstetric and neonatal outcomes of pregnant patients with COVID-19 in Wuhan, China: a retrospective, single-centre, descriptive study. *Lancet Infect Dis* Published Online First: March 2020. doi:10.1016/S1473-3099(20)30176-6
- 12 Di Mascio D, Khalil A, Saccone G, *et al.* Outcome of Coronavirus spectrum infections (SARS, MERS, COVID 1 -19) during pregnancy: a systematic review and meta-analysis. *Am J Obstet Gynecol MFM* 2020;;100107. doi:10.1016/j.ajogmf.2020.100107
- 13 Zeng, Lingkong; Xia S. Neonatal Early-Onset Infection With SARS-CoV-2 in 33 Neonates Born to Mothers With COVID-19 in Wuhan, China. *JAMA* 2020;**23**:2–4.
- 14 Buekens P, Alger J, Bréart G. A call for action for COVID-19 surveillance and research during pregnancy. *Lancet Glob Heal* 2020;;2019–20. doi:10.1016/S2214-109X(20)30206-0
- 15 World Health Organization. Q&A on COVID-19, pregnancy and childbirth. 2020. <https://www.who.int/news-room/q-a-detail/q-a-on-covid-19-and-pregnancy-and-childbirth>

- 16 Delamou A, Ayadi AM El, Sidibe S, *et al.* Effect of Ebola virus disease on maternal and child health services in Guinea: a retrospective observational cohort study. *Lancet Glob Heal* 2017;**5**:e448–57. doi:10.1016/S2214-109X(17)30078-5
- 17 Leno NN, Delamou A, Koita Y, *et al.* Ebola virus disease outbreak in Guinea: what effects on prevention of mother-to-child transmission of HIV services? *Reprod Health* 2018;**15**:60. doi:10.1186/s12978-018-0502-y
- 18 World Health Organisation. Operational guidance for maintaining essential health services during an outbreak. 2020;:1–14. <https://www.who.int/publications-detail/covid-19-operational-guidance-for-maintaining-essential-health-services-during-an-outbreak>
- 19 Cipriano PF. 100 years on: the Spanish Flu, pandemics and keeping nurses safe. *Int Nurs Rev* 2018;**65**:305–6. doi:10.1111/inr.12483
- 20 McMahon SA, Ho LS, Brown H, *et al.* Healthcare providers on the frontlines: a qualitative investigation of the social and emotional impact of delivering health services during Sierra Leone's Ebola epidemic. *Health Policy Plan* 2016;**31**:1232–9.
- 21 Liu Q, Yang J, Liu Q, *et al.* The experiences of health-care providers during the COVID-19 crisis in China: a qualitative study. *Lancet Glob Heal* Published Online First: 29 April 2020. doi:10.1016/S2214-109X(20)30204-7
- 22 Sochas L, Channon AA, Nam S. Counting indirect crisis-related deaths in the context of a low-resilience health system: the case of maternal and neonatal health during the Ebola epidemic in Sierra Leone. *Health Policy Plan* 2017;**32**:iii32–9. doi:10.1093/heapol/czx108
- 23 Nuzzo JB, Meyer D, Snyder M, *et al.* What makes health systems resilient against infectious disease outbreaks and natural hazards? Results from a scoping review. *BMC Public Health* 2019;**19**:1–9. doi:10.1186/s12889-019-7707-z
- 24 Oppenheim B, Gallivan M, Madhav NK, *et al.* Assessing global preparedness for the next pandemic: Development and application of an Epidemic Preparedness Index. *BMJ Glob Heal* 2019;**4**:1–9. doi:10.1136/bmjgh-2018-001157
- 25 Camara BS, Delamou A, Diro E, *et al.* Effect of the 2014/2015 Ebola outbreak on reproductive health services in a rural district of Guinea: An ecological study. *Trans R Soc Trop Med Hyg* 2017;**111**:22–9. doi:10.1093/trstmh/trx009
- 26 Bietsch K, Williamson J, Reeves M. Family Planning During and After the West African Ebola Crisis. *Stud Fam Plann* 2020;**51**:71–86. doi:10.1111/sifp.12110
- 27 Stein D, Ward K, Cantelmo C. Estimating the Potential Impact of COVID-19 on Mothers and Newborns in Low- and Middle-Income Countries. *Health Policy Plus* 2020. <https://medium.com/@HealthPolicyPlus/estimating-the-potential-impact-of-covid-19-on-mothers-and-newborns-in-low-and-middle-income-3a7887e4a0ff>
- 28 Robertson T, Carter E, Chou V, *et al.* Early estimates of the indirect effects of the coronavirus pandemic on maternal and child mortality in low- and middle-income countries. *Lancet Glob Heal* 2020.
- 29 Riley T, Sully E, Ahmed Z, *et al.* Estimates of the Potential Impact of the COVID-19 Pandemic on Sexual and Reproductive Health In Low-and Middle-Income Countries. *Int Perspect Sex Reprod Health* 2020;**46**:73–6.

- 30 Graham W, Afolabi BB, Benova L, *et al.* Protecting hard won gains for mothers and newborns in low-and-middle-income countries in the face of COVID-19: call for a service safety net. *BMJ Glob Heal* 2020.
- 31 World Health Organization. Definition of skilled health personnel providing care during childbirth: the 2018 joint statement by WHO, UNFPA, UNICEF, ICM, ICN, FIGO and IPA. 2018;;1–4.
- 32 The World Bank. World Bank Country and Lending Groups. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>
- 33 Liverman CT, Ingalls CE, Fulco CE, *et al.* Understanding the information needs of health professionals. In: *Toxicology and environmental health information resources: The role of the national library of medicine*. National Academies Press (US) 1997.
- 34 Cinelli M, Quattrocioni W, Galeazzi A, *et al.* The COVID-19 Social Media Infodemic. 2020;;1–18. <http://arxiv.org/abs/2003.05004>
- 35 Ministry of Health - Kenya. GUIDELINES A Kenya Practical Guide for Continuity of Reproductive , Maternal , Newborn and Family Planning Care and Services in the Background of COVID19 Pandemic. 2020.
- 36 NHS. Clinical guide for the temporary reorganisation of intrapartum maternity care during the coronavirus pandemic. 2020. <https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/52/2020/04/C0241-specialty-guide-intrapartum-maternity-care-9-april-2020.pdf>
- 37 Chawla D, Chirala D, Dalwai S, *et al.* Perinatal-Neonatal Management of COVID-19 Infection - Guidelines of the Federation of Obstetric and Gynecological Societies of India (FOGSI), National Neonatology Forum of India (NNF), and Indian Academy of Pediatrics (IAP). *Indian Pediatr* 2020.
- 38 Federal Ministry of Health - Nigeria, Nigeria Centre for Disease Control. COVID-19 Guidelines for Pregnant Women and Nursing Mothers - Nigeria. 2020.<https://reliefweb.int/report/nigeria/covid-19-guidelines-pregnant-women-and-nursing-mothers> (accessed 6 May 2020).
- 39 New York State Department of Health. Health Advisory : COVID-19 Guidance for Hospital Operators Regarding Visitation Updated Guidance regarding Obstetrical and Pediatric Settings. 2020;19:2020. [https://northportny.gov/wp-content/uploads/2020/03/doh\\_covid19\\_obpedsvisitation\\_032720.pdf](https://northportny.gov/wp-content/uploads/2020/03/doh_covid19_obpedsvisitation_032720.pdf)
- 40 Royal College of Obstetricians & Gynaecologists, The Royal College of Midwives. Guidance for provision of midwife-led settings and home birth in the evolving coronavirus (COVID-19) pandemic. 2020.<https://www.rcog.org.uk/globalassets/documents/guidelines/2020-04-09-guidance-for-provision-of-midwife-led-settings-and-home-birth-in-the-evolving-coronavirus-covid-19-pandemic.pdf>
- 41 Royal College of Obstetricians & Gynaecologists. Guidance for deployment of obstetrics and gynaecology staff during the COVID-19 pandemic. 2020. <https://www.rcog.org.uk/globalassets/documents/guidelines/2020-05-01-guidance-for-deployment-of-obstetrics-and-gynaecology-staff-during-the-covid-19-pandemic.pdf>
- 42 World Health Organization. Coronavirus disease (COVID-19) outbreak: rights, roles and responsibilities of health workers, including key considerations for occupational safety and health, 18 March 2020. 2020. [https://www.who.int/publications-detail/coronavirus-disease-\(covid-19\)-outbreak-rights-roles-and-responsibilities-of-health-workers-including-key-considerations-for](https://www.who.int/publications-detail/coronavirus-disease-(covid-19)-outbreak-rights-roles-and-responsibilities-of-health-workers-including-key-considerations-for)

occupational-safety-and-health

- 43 Tran NT, Tappis H, Spilotros N, *et al.* Not a luxury: a call to maintain sexual and reproductive health in humanitarian and fragile settings during the COVID-19 pandemic. *Lancet Glob Heal* Published Online First: April 2020. doi:10.1016/S2214-109X(20)30190-X
- 44 Wilson AN, Ravaldi C, Scoullar MJL, *et al.* Caring for the carers: Ensuring the provision of quality maternity care during a global pandemic. *Women Birth* Published Online First: April 2020. doi:10.1016/j.wombi.2020.03.011
- 45 Brussels U. Midwives step up to support pregnant women during pandemic. 2020. <https://www.un.org/en/pregnant-women-are-worried-about-giving-birth-during-pandemic>
- 46 Ossola A. The challenges of supporting pregnant women during Covid-19, from a midwife. Published Online First: 2020. <https://qz.com/1847930/covid-19-is-transforming-the-role-of-midwives-during-childbirth/>
- 47 Audet C. Giving birth amid a pandemic in Belgium: the challenges faced by mothers and midwives. *BMJ Glob. Heal. blog.* 2020. <https://blogs.bmj.com/bmjgh/2020/04/28/giving-birth-amid-a-pandemic-in-belgium-the-challenges-faced-by-mothers-and-midwives/>
- 48 Hussein J. COVID-19: What implications for sexual and reproductive health and rights globally? *Sex. Reprod. Heal. matters.* 2020;**28**:1746065. doi:10.1080/26410397.2020.1746065
- 49 Martinez-Alvarez M, Jarde A, Usuf E, *et al.* COVID-19 pandemic in west Africa. *Lancet Glob Heal* 2020;**8**:e631–2.
- 50 Hopman J, Allegranzi B, Mehtar S. Managing COVID-19 in low-and middle-income countries. *Jama* 2020.
- 51 Gharibah M, Mehchy Z. COVID-19 pandemic: Syria's response and healthcare capacity. 2020.
- 52 San Lau L, Samari G, Moresky RT, *et al.* COVID-19 in humanitarian settings and lessons learned from past epidemics. *Nat Med* 2020;:1–2.
- 53 Vince G. The world's largest refugee camp prepares for covid-19. *Bmj* 2020;**368**.
- 54 DeJong J, Ghattas H, Bashour H, *et al.* Reproductive, maternal, neonatal and child health in conflict: a case study on Syria using Countdown indicators. *BMJ Glob Heal* 2017;**2**:e000302.
- 55 Islam MM, Nuzhath T. Health risks of Rohingya refugee population in Bangladesh: a call for global attention. *J Glob Health* 2018;**8**.
- 56 Hynes M, Sakani O, Spiegel P, *et al.* A study of refugee maternal mortality in 10 countries, 2008–2010. *Int Perspect Sex Reprod Health* 2012;:205–13.
- 57 Gibson-Helm M, Boyle J, Cheng I-H, *et al.* Maternal health and pregnancy outcomes among women of refugee background from Asian countries. *Int J Gynecol Obstet* 2015;**129**:146–51.
- 58 Elston JWT, Moosa AJ, Moses F, *et al.* Impact of the Ebola outbreak on health systems and population health in Sierra Leone. *J Public Health (Bangkok)* 2016;**38**:673–8.
- 59 World Health Organization. Mental health and psychosocial considerations during the COVID-19 outbreak . [https://www.who.int/docs/default-source/coronaviruse/mental-health-considerations.pdf?sfvrsn=6d3578af\\_2](https://www.who.int/docs/default-source/coronaviruse/mental-health-considerations.pdf?sfvrsn=6d3578af_2) (accessed 4 May 2020).

- 60 Kisely S, Warren N, McMahon L, *et al.* Occurrence, prevention, and management of the psychological effects of emerging virus outbreaks on healthcare workers: rapid review and meta-analysis. *BMJ* 2020;**369**:m1642. doi:10.1136/bmj.m1642
- 61 Strong AE. Causes and effects of occupational risk for healthcare workers on the maternity ward of a Tanzanian hospital. *Hum. Organ.* 2018;**77**:273–86. doi:10.17730/0018-7259.77.3.273
- 62 Dunn AC, Walker TA, Redd J, *et al.* Nosocomial transmission of Ebola virus disease on pediatric and maternity wards: Bombali and Tonkolili, Sierra Leone, 2014. *Am J Infect Control* 2016;**44**:269–72. doi:10.1016/j.ajic.2015.09.016
- 63 Moghri J, Rashidian A, Arab M, *et al.* Implications of dual practice among health workers: A systematic review. *Iran J Public Health* 2017;**46**:153–64.
- 64 World Health Organization. Rational use of personal protective equipment for coronavirus disease 2019 (COVID-19). 2020. [https://apps.who.int/iris/bitstream/handle/10665/331215/WHO-2019-nCov-IPCPPE\\_use-2020.1-eng.pdf](https://apps.who.int/iris/bitstream/handle/10665/331215/WHO-2019-nCov-IPCPPE_use-2020.1-eng.pdf)
- 65 Penn-Kekana L. COVID-19 and MNCH: Beyond the models, what are we hearing from countries? 2020. <https://www.harpnet.org/blog/covid-19-and-mch-beyond-the-models-what-are-we-hearing-from-countries/>
- 66 Elston JWT, Cartwright C, Ndumbi P, *et al.* The health impact of the 2014–15 Ebola outbreak. *Public Health* 2017;**143**:60–70.
- 67 Brolin Ribacke KJ, Saulnier DD, Eriksson A, *et al.* Effects of the West Africa Ebola virus disease on health-care utilization—a systematic review. *Front public Heal* 2016;**4**:222.
- 68 McQuilkin PA, Udhayashankar K, Niescierenko M, *et al.* Health-Care Access during the Ebola Virus Epidemic in Liberia. *Am J Trop Med Hyg* 2017;**97**:931–6.
- 69 Jones SA, Gopalakrishnan S, Ameh CA, *et al.* ‘Women and babies are dying but not of Ebola’: the effect of the Ebola virus epidemic on the availability, uptake and outcomes of maternal and newborn health services in Sierra Leone. *BMJ Glob Heal* 2016;**1**:e000065.
- 70 Wanduru P, Tetui M, Waiswa P. COVID-19 response in Uganda: notes and reflections. *BMJ Glob. Heal. blogs.* 2020.
- 71 UNFPA. COVID-19 Technical brief for maternity services. 2020. <https://www.unfpa.org/resources/covid-19-technical-brief-maternity-services>
- 72 International Confederation of Midwives. Women’s Rights in Childbirth Must be Upheld During the Coronavirus Pandemic. 2020.
- 73 Bobrow E. A Chaotic Week for Pregnant Women in New York City. *New Yorker* Published Online First: 2020. <https://www.newyorker.com/science/medical-dispatch/a-chaotic-week-for-pregnant-women-in-new-york-city>
- 74 Lowe NK. Reflections on COVID-19. *J Obstet Gynecol Neonatal Nurs* 2020.
- 75 Ferrazzi E, Frigerio L, Savasi V, *et al.* Vaginal delivery in SARS-CoV-2 infected pregnant women in Northern Italy: a retrospective analysis. *BJOG an Int J Obstet Gynaecol* 2020.
- 76 Petit G, Huet N. Coronavirus: Baby blues as some mothers in France separated from newborn amid COVID-19 fears | Euronews. 2020. <https://www.euronews.com/2020/03/31/coronavirus-baby->

- blues-as-some-mothers-in-france-separated-from-newborn-amid-covid-19-fear (accessed 6 May 2020).
- 77 Campbell OMR, Calvert C, Testa A, *et al.* The scale, scope, coverage, and capability of childbirth care. *Lancet (London, England)* 2016;**388**:2193–208. doi:10.1016/S0140-6736(16)31528-8
- 78 Nelson R. COVID-19 disrupts vaccine delivery. *Lancet Infect Dis* 2020.
- 79 World Health Organization. Guiding principles for immunization activities during the COVID-19 pandemic. 2020. <https://apps.who.int/iris/handle/10665/331590>
- 80 Hollander JE, Carr BG. Virtually perfect? Telemedicine for COVID-19. *N Engl J Med* 2020.
- 81 Jaffré Y, Olivier de Sardan J-P. Les difficiles relations entere soignants et soigne dans cinq capitales d’Afrique de l’Ouest. *Bull Amades* 2003;:1–466. doi:10.3917/kart.jaffr.2003.01
- 82 Pourette D. *Femmes, enfants et santé à Madagascar – Approches anthropologiques comparées*. Paris: : Éditions L’Harmattan 2018.
- 83 Nganga SW, Otieno NA, Adero M, *et al.* Patient and provider perspectives on how trust influences maternal vaccine acceptance among pregnant women in Kenya. *BMC Health Serv Res* 2019;**19**:1–13. doi:10.1186/s12913-019-4537-8
- 84 Gilson L. Trust in health care: Theoretical perspectives and research needs. *J Heal Organ Manag* 2006;**20**:359–75. doi:10.1108/14777260610701768
- 85 Mestre C. *Maladies et violences ordinaires dans un hôpital malgache*. Editions L. Paris: 2014.
- 86 Fairhead J. Understanding Social Resistance to the Ebola Response in the Forest Region of the Republic of Guinea: An Anthropological Perspective. *Afr Stud Rev* 2016;**59**:7–31. doi:10.1017/asr.2016.87
- 87 Huysmans E. *Quand la santé questionne les liens : anthropologie de la santé maternelle et infantile dans le Sud-Ouest de Madagascar*. 2019.
- 88 Ronse M. Biology and culture are inseparable – considerations for the “exit strategy” expert group from the field of medical anthropology. 2020. [https://07323a85-0336-4ddc-87e4-29e3b506f20c.filesusr.com/ugd/3cbdf6\\_f9ef1e533ce54fd4844a76a805aa084e.pdf](https://07323a85-0336-4ddc-87e4-29e3b506f20c.filesusr.com/ugd/3cbdf6_f9ef1e533ce54fd4844a76a805aa084e.pdf)
- 89 Kruk ME, Myers M, Varpilah ST, *et al.* What is a resilient health system? Lessons from Ebola. *Lancet* 2015;**385**:1910–2. doi:10.1016/S0140-6736(15)60755-3

## Supplementary File 1 - Online Questionnaire – Round 1

### Welcome message

Thank you for your interest in this survey. This research is being conducted by a group of leading maternal and newborn health researchers, and led by the Institute of Tropical Medicine (ITM) in Antwerp, Belgium.

This survey is aimed at healthcare workers providing care to women and their babies: antenatal, intrapartum and postnatal care. We would like to start by thanking you for the care you provide to women, babies and families at this difficult time. We hope that by contributing to this research your voice will be heard and your efforts will be better understood and acknowledged.

### Study Information & Online Consent Form

You are invited to participate in a research study on response to COVID-19 among maternity providers **globally**. The purpose of this study is to understand the range of actions taken to ensure care continues to be provided to women and their babies: antenatal, intrapartum and postnatal care during this pandemic. This survey will be distributed every 3-4 weeks to track the rapidly developing situation over time. You are free to take part on this survey once or multiple times.

We are aware that maternity care is provided very differently across many countries, and have provided space for your responses rather than restricting you to pre-defined options. We appreciate the time it takes to complete such questions. By doing so, you are helping us develop a more time-efficient survey for the next round. We appreciate that some questions are currently more relevant to some contexts/countries than others, and we thank you for your patience as you **answer or skip these questions depending on your context**. The situation is changing rapidly and we want to document how it develops over time.

This online survey will take approximately 15-30 minutes to complete.

Your decision to participate in this study is completely voluntary and you have the right to end your participation at any time by closing your browser window or mobile application. You may also skip any questions you do not wish to answer. Your participation in this research will be completely confidential (we will remove any identifiers) and data will be reported in aggregate.

*I agree to these terms:*

I have read and understand the above consent form,

I certify that I am a healthcare professional and, by clicking the button below to enter the survey, I indicate my willingness to take part in the study voluntarily.

I would like to receive an email when the next round of this surveys is available. I understand that I am under no obligation to respond in the future, and that my email will not be stored together with my responses to this survey, or used for any other purpose.

Email address: [text field]

We will be seeking to contact selected healthcare providers for individual interviews to understand their perspective and experience. If you give consent to being contacted, please provide an email address where we can reach you. This information will be treated confidentially.

I give consent to be contacted by the researchers for additional information

Email address: [text field]

### Researcher contact information

This study is coordinated by Associate Professor Lenka Benova from the Institute of Tropical Medicine in Antwerp, Belgium. If you have any concerns about this study, your confidentiality or data, please contact Dr Benova by email ([lbenova@itg.be](mailto:lbenova@itg.be)) or phone/Whatsapp (+31 61 26 999 64).

**Part 1. We would like to ask a few questions about your background**

Q#	Question	Response
1	In which <b>country</b> are you based (providing healthcare) at the moment?	[drop down menu of countries]
2	In which <b>region</b> of the country do you work? (please provide region, district, province, state, or governorate)	[free text]
3	What is your <b>job</b> ? (choose one)	<ul style="list-style-type: none"> <li>- Midwife</li> <li>- Nurse-midwife</li> <li>- Nurse</li> <li>- Obstetrician/gynecologist</li> <li>- Anaesthesiologist</li> <li>- Surgeon</li> <li>- Neonatologist</li> <li>- Pediatrician</li> <li>- General practitioner</li> <li>- Medical doctor (other/no specialisation)</li> <li>- Medical student/intern/resident</li> <li>- Medical officer</li> <li>- Clinical officer</li> <li>- Nurse or midwife in training</li> <li>- Ultrasound technician/ sonographer</li> <li>- Health technician</li> <li>- Paramedic</li> <li>- Community health worker/ Outreach worker</li> <li>- Lactation counsellor</li> <li>- Other: specify</li> </ul>
4	What is your <b>position</b> ? (choose one)	<ul style="list-style-type: none"> <li>- Head of facility (director, administrator)</li> <li>- Head of department or ward</li> <li>- Head of team</li> <li>- Team member</li> <li>- Locum or interim member</li> <li>- Other: specify</li> </ul>
5	What is your <b>gender</b> ? (choose one)	<ul style="list-style-type: none"> <li>- Female</li> <li>- Male</li> <li>- Other/Prefer not to say</li> </ul>
6	What <b>type of maternal and/or neonatal health care</b> do you currently provide as an individual? (mark all that apply)	<ul style="list-style-type: none"> <li>- Outpatient antenatal care</li> <li>- Outpatient (home-based) childbirth care</li> <li>- Outpatient postnatal care</li> <li>- Outpatient breastfeeding support</li> <li>- Inpatient antenatal care</li> <li>- Inpatient childbirth care</li> <li>- Inpatient postnatal care (mother and/or babies)</li> <li>- Surgical care</li> <li>- Neonatal care for small and sick newborns</li> <li>- Home visits</li> <li>- Community outreach, home visits, health education outside facility</li> <li>- Abortion care</li> <li>- Post-abortion care</li> <li>- Other: specify</li> </ul>

**Part 2. Setting: Can you tell us about the facility setting in which you work now**

Q#	Question	Response	Notes
1	In which <b>level</b> of health care institution do you primarily work? (if none of the response options fit well, please use the “Other” option and write what your facility type is called in your country)	<ul style="list-style-type: none"> <li>- Referral hospital</li> <li>- District/regional hospital</li> <li>- Health center</li> <li>- Polyclinic</li> <li>- Clinic</li> <li>- Health post/unit</li> <li>- Dispensary</li> <li>- Other: specify</li> </ul>	
2	What organisation <b>type</b> is your institution?	<ul style="list-style-type: none"> <li>- Public (national)</li> <li>- Public (university or teaching)</li> <li>- Public (district level or below)</li> <li>- Social security</li> <li>- Health insurance or HMO</li> <li>- Private university</li> <li>- Private for profit</li> <li>- Non-governmental</li> <li>- Faith-based or mission</li> <li>- Other: specify</li> </ul>	
3	In what type of <b>geographic area</b> is your facility located?	<ul style="list-style-type: none"> <li>- Large city (&gt;1 mil inhabitants)</li> <li>- Small city (100,000 to 1 mil inhabitants)</li> <li>- Town (&lt;100,000 inhabitants)</li> <li>- Village or rural area</li> <li>- Refugee or displaced persons camp</li> <li>- Other</li> </ul>	
4	How many <b>maternity beds</b> does your facility have (include antenatal, labour/childbirth and postnatal). Approximate number is ok	None Number: _____[free text]	
5	How many <b>births</b> took place in your facility in <b>2019</b> ? Approximate number is ok	None Number: _____[free text] Don't know	
6	Does your facility provide <b>caesarean sections</b> ?	Yes No	If no, skip to 8
7	If yes, what is the <b>csection rate</b> (% of births by csection) in your facility? Approximate number is ok	Number: _____[free text] Don't know	
8	Does your facility have an <b>Intensive Care Unit (ICU)</b> which can admit <b>women</b> with obstetric complications? (ICU is defined as a clinical area where ventilatory support can be provided)	Yes No Don't know	
9	Does your facility have a <b>neonatal intensive care unit (NICU)</b> ? (Neonatal intensive care is defined as a unit that provides invasive ventilatory support to small and sick newborns, not just CPAP)	Yes No Don't know	
10	Does your facility <b>receive maternity referrals</b> from other facilities, meaning that patients are sent to your facility from other health facilities?	Yes No Don't know	
11	Is running <b>water and soap</b> always available for hand hygiene on your ward <b>for the use of staff</b> ?	Yes No Don't know	

12	Is running <b>water and soap</b> always available on your ward for the use of <b>patients, visitors, companions</b> ?	Yes No Don't know	
13	Is there always <b>sufficient water and disinfectant for cleaning surfaces</b> ?	Yes No Don't know	

### Part 3. COVID-19 preparedness

Q#	Question	Response	Notes
1	Has your institution or ward provided you with <b>any information</b> on how to prepare for COVID-19?	Yes No	No – skip to 6
2	What did you learn from this information? Please list main areas or themes	[free text]	
	On a scale from 1 (poor) to 5 (excellent), how would you rate the following dimensions of this information:	1 – poor 2 – somewhat useful 3 – average 4 – good 5 – excellent Not applicable	
3	Clarity		
4	Helpfulness for your daily work		
5	Value in helping you feel safe		
6	Has your institution or ward provided you with any training on COVID-19, for example simulations or drills?	Yes No	
7	Have you received new or updated guidelines specifically for the provision of care to pregnant, labouring or postpartum women and their newborns because of COVID-19?	Yes No	If no, skip to 9
8	If yes, which guidelines? Please list all used (WHO, FIGO, ICM, RCOG, RCPCH, RCN, RCM, COINN, your country's Ministry of Health, Country professional organisations etc)	Free text	
9	Have you personally searched for source of guidance and sources of information to prepare for COVID-19 in your work?	Yes No	
10	Have you received information related to COVID-19 and your work informally through other colleagues (in your own facility or outside)?	Yes No	
11	Have you been a part of any self-organisation on the part of healthcare workers in response to the COVID-19 outbreak? (exchange of information, virtual discussion groups (Whatsapp, Facebook, etc)	Yes No	
12	Has your facility published or distributed any materials (brochure, flier, posters, etc) covering COVID-19 targeted toward pregnant, labouring, or postnatal women?	Yes No Don't know	If no or don't know - skip to 15

13	What kind of information is on these materials? (social distancing, symptoms, when/how to self-isolate, when/where to seek care etc.)	Free text	
14	In what form is it provided? (mark all that apply)	-Health talks -Leaflets/fliers -Posters -Counselling during consultations -Facility website -Phone line with advice -Other: specify	
15	In your facility, do you feel that patients' questions about COVID-19 are being addressed adequately by staff?	Yes No Don't know	
16	What are you worried about most at this time in regard to <b>being able to provide care to women and newborns</b> ?	Free text	
17	Has your facility set up a well sign-posted general entrance and screening area for COVID-19 suspected cases? (regardless whether for maternity patients or not)	-Yes -Some measures taken but not done well -No measures taken -Don't know	
18	Has your facility reserved <b>isolation rooms</b> for COVID-19 suspected cases?	Yes No Don't know	
19	Has <b>routine cleaning of the maternity ward</b> changed in response to COVID-19?	Yes, increased Yes, decreased Unchanged Don't know	

#### Part 4. Response to COVID-19 in your facility

Q#	Question	Response	Notes
1	Is your facility currently screening for COVID-19 symptoms among maternity patients?	Yes No Don't know Other: specify	
2	Is it possible to order a test for COVID-19 at the moment for maternity patients at your facility?	Yes No Don't know Other: specify	If no/DK: skip to 4
3	If yes, how long does it take to get a result? (note whether your response is in hours or days)	[free text]	
4	Are the testing criteria for COVID-19 clear to you? (the conditions/symptoms for which a test can be ordered)	Yes No	If no – go to 6
5	Can you describe these criteria	[free text]	
6	Have you had any maternity patients with COVID-19 in your facility so far?	Yes, suspected Yes, confirmed Yes, both confirmed and suspected cases No Don't know Other: specify	If no or DK – go to 8
7	If yes, approximate number	[free text]	
8	Is there a designated COVID-19 lead person / liaison or team in the maternity ward or the facility?	-Yes, in maternity -Yes, in facility as a whole -Yes, both in the maternity ward and in the facility as a whole -No, neither maternity nor facility -Don't know	
9	Were you aware that the WHO developed a pregnancy/postpartum module to be included in the current Case Report Forms (CRFs) recommended to report COVID-19? cases ( <a href="https://isaric.tghn.org/novel-coronavirus/">https://isaric.tghn.org/novel-coronavirus/</a> )	Yes No	If no, skip to 11
10	If yes, are you using this module or preparing to use this module in your facility?	Yes, already using Yes, preparing to start No Don't know Other: specify	
11	On a scale from 1 (not at all) to 5 (I am very clear), do you personally feel you know what you should do if a woman with COVID-19 symptoms arrives in your facility today?	1 – Not at all clear 2 – Some points are clear to me, but I am not confident in what to do 3 – Somewhat clear but major issues remain 4 – I am mostly clear but some questions / areas of concern remain 5 – I am very clear	
12	On a scale from 1 (not at all) to 5 (completely), do you feel that you are	1 – not at all 2 –minimal protection	

	sufficiently protected from infection with COVID-19 in your workplace?	3 – some protection 4 – well protected 5 – completely protected	
<b>Is a sufficient quantity of personal protective equipment (PPE) available to you?</b>			
13	Gloves	Yes - No	
14	Masks	Yes - No	
15	Aprons	Yes - No	

**Part 5. Your work and experience in light of the COVID-19 outbreak**

Q#	Question	Response	Notes
1	Has your work been affected by the COVID-19 outbreak?	- Yes - No	If no skip to 3
2	If yes, how has your work changed? Please feel free to describe changes you perceive as important for patients as well as yourself, your team and the institution. (example: work hours, overtime, types of work being done, collaboration between team members, collaboration between facilities, etc)	Free text	
3	On a scale of 1 (not at all) to 5 (completely), do you feel that your concerns about the response to COVID-19 have been addressed by your facility or ward?	1 – not at all 2 – minimally 3 – somewhat 4 – well 5 – completely	
4	How would you rate your own levels of stress at this time?	-Same as usual -Somewhat higher than usual -Substantially higher than usual	
6	Do you consider your personal role as a health worker in this COVID-19 outbreak is valued by the community you are serving?	Not at all Very little Somewhat Highly Unsure/don't know	
7	What is the one thing that could be done to support you more at this time of outbreak?	Free text	
8	Is there anything else you would like to share?	Free text	

You have now completed the main part of the questionnaire. If you are interested in accessing information and guidance on COVID-19, please click [here](#).

We value your time and experiences greatly. Thank you for your participation. If you have more time, we would like to ask some questions about how the provision of care in your facility has been affected by COVID-19. Please click [\[here\]](#) if you would like to continue to this additional last section of this survey.

**Additional module. Effect of COVID-19 on the provision of maternal and newborn care**

Can you describe how the COVID-19 outbreak has affected the provision of care to women and newborns in your facility and community? This includes changes made directly in response to the threat of COVID-19 and other indirect influences (for example, pressure on the health system).

Q#	Question	Response
1	Changes to provision of <b>outpatient antenatal</b> care (examples include staffing levels, location – in person or phone/internet, waiting times, patient flow, ability to conduct all routine tests and investigations, etc)	Yes - No If yes, please describe the changes
2	Changes to provision of <b>inpatient antenatal</b> care	Yes - No If yes, please describe the changes
3	Changes to capacity to provide <b>intrapartum care</b> (number of rooms or beds, availability of equipment, supplies and medications)	Yes - No If yes, please describe the changes
4	Changes to rules on number or type of <b>labour companions</b> (includes family members and professional doulas, whether able to stay overnight, etc)	Yes - No If yes, please describe the changes
5	Changes to <b>pain relief</b> options available to women in labour	Yes - No If yes, please describe the changes
6	Changes to <b>rules on induction</b> of labour	Yes - No If yes, please describe the changes
7	Changes to <b>caesarean section provision</b> (e.g. location of theatre, type of anaesthesia, guidelines for elective csections)	Yes - No If yes, please describe the changes
8	Changes to <b>visiting hours</b> or number/type of visitors, for mothers and newborns	Yes - No If yes, please describe the changes
9	Changes to <b>inpatient postnatal</b> care provision following vaginal births (for example, shorter length of stay, frequency of routine checks)	Yes - No If yes, please describe the changes
10	Changes to <b>inpatient postnatal</b> care provision following caesarean section births (for example, change in cleaning schedules, hand hygiene indications, hand hygiene supplies, delivery equipment decontamination and sterilisation)	Yes - No If yes, please describe the changes
11	Changes to <b>outpatient postnatal</b> care provision (ability to follow-up in women's homes, breastfeeding counselling, postpartum family planning etc)	Yes - No If yes, please describe the changes
12	Changes to provision of <b>routine newborn</b> care before discharge (screenings, vaccinations, etc)	Yes - No

		If yes, please describe the changes
13	Changes to provision of <b>newborn intensive</b> care (example: bed capacity, oxygen, equipment)	Yes - No If yes, please describe the changes
14	Changes to provision of <b>non-essential care</b> (such as cancellations of elective surgery, gynecological procedures, IVF provision, etc)	Yes – No If yes, please describe the changes
15	Changes to <b>staffing levels or team/shift composition</b> (for example, more people hired, such as cleaners, change in shift hours, staff being absent due to symptoms )	Yes - No If yes, please describe the changes
16	Changes to process of <b>referring patients to other facilities</b> (timing, destination, mode of transport, capacity, guidelines)	Yes - No If yes, please describe the changes
17	Changes to <b>process of receiving incoming maternity referrals</b> , including antenatal and emergency. (timing, destination, mode of transport, capacity, guidelines)	Yes - No If yes, please describe the changes
18	Do you feel that <b>women’s use of care</b> in your facility has been affected by COVID-19? For example, do you see fewer or more patients than usual? If so, why?	Yes - No If yes, please describe the changes

Supplementary File 2 Frequency distribution of respondents by country

