

Delivering health interventions to women, children, and adolescents in conflict settings: What have we learned from 10 country case studies?

Neha S. Singh PhD¹, Anushka Ataullahjan² PhD, Khadidiatou Ndiaye PhD³, Jai K. Das⁴, Prof. Paul H. Wise PhD⁵, Chiara Altare PhD⁶, Zahra Ahmed MSc⁷, Samira Sami PhD^{6,8}, Chaza Akik DrPH⁹, Hannah Tappis DrPH⁶, Shafiq Mirzazada MPH¹⁰, Prof. Isabel C. Garcés-Palacio DrPH¹¹, Hala Ghattas PhD⁹, Prof. Ana Langer MD¹², Prof. Ronald J. Waldman^{13,14}, Prof. Paul Spiegel PhD⁶, Prof. Zulfiqar A. Bhutta PhD^{2,4}, Prof. Karl Blanchet PhD^{1,15*}, and the BRANCH steering committee

¹ Health in Humanitarian Crises Centre, London School of Hygiene and Tropical Medicine, London, United Kingdom

² Center for Global Child Health, Hospital for Sick Children, Toronto, Canada

³ Milken Institute School of Public Health, George Washington University, Washington, DC, USA

⁴ Center of Excellence in Women and Child Health, The Aga Khan University, Karachi, Pakistan

⁵ Department of Pediatrics, Stanford University School of Medicine, Stanford, California, USA

⁶ Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

⁷ Somali Disaster Resilience Institute, Mogadishu, Somalia

⁸ CDC National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention
Division of HIV/AIDS Prevention, Atlanta, Georgia, USA

⁹ Center for Research on Population and Health, Faculty of Health Sciences, American University of Beirut, Beirut, Lebanon

¹⁰ Aga Khan University, Kabul, Afghanistan

¹¹ Epidemiology group, School of Public Health, Universidad de Antioquia, Medellín, Colombia

¹² Women and Health Initiative; Department of Global Health and Population, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

¹³ Doctors of the World USA, 222 Broadway, 19th Floor, New York, NY 10038, USA

¹⁴ Global Health Department, Milken Institute School of Public Health, George Washington University, 950 New Hampshire Ave. NW 4th Floor, Washington, DC 20052, USA.

¹⁵ The Geneva Centre of Humanitarian Studies, University of Geneva, Graduate Institute, Geneva 1211, Switzerland

***Corresponding author:**

Karl Blanchet, PhD; *Address:* The Geneva Centre of Humanitarian Studies, University of Geneva, Graduate Institute, Geneva 1211, Switzerland; *Email:* karl.blanchet@unige.ch

Word count: 4467

Summary

Armed conflict disproportionately affects women, newborns, children, and adolescents. Our study presents insights from a collection of ten country case studies aiming to assess the provision of sexual, reproductive, maternal, newborn, child and adolescent health and nutrition (abbreviated to women's and children's health, i.e WCH in this paper) interventions in conflict-affected settings in Afghanistan, Colombia, Democratic Republic of Congo, Mali, Nigeria, Pakistan, Somalia, South Sudan, Syria and Yemen. We found that despite large variations in contexts and decision-making processes, antenatal care, basic emergency obstetric and newborn care (BEmONC), comprehensive emergency obstetric and newborn care (CEmONC), immunisation, treatment of common childhood illnesses, infant and young child feeding (IYCF) and malnutrition treatment and screening were prioritised in these ten conflict settings. Many lifesaving WCH services, including the majority of reproductive, newborn and adolescent health services, are not reported as being delivered in the ten conflict settings, and interventions to address stillbirths are absent. International donors remain the primary drivers of influencing the what, where, and how of implementing WCH interventions. Interpretation of WCH outcomes in conflict settings are particularly context-dependent given the myriad of complex factors that constitute conflict and their interactions. Moreover, the comprehensiveness and quality of data remain limited in conflict settings. The dynamic nature of modern conflict and the expanding role of Non-State Armed groups in large geographic areas pose new challenges to delivering WCH services. However, the humanitarian system is creative and pluralistic and has developed some novel solutions to bring lifesaving WCH services closer to populations using new modes of delivery. These solutions, when rigorously evaluated, can represent concrete response to current implementation challenges to modern armed conflicts.

Key messages

1. Many lifesaving women's and children's health (WCH) services for key populations in conflict settings are not delivered everywhere.
2. Priorities of donors are the primary drivers of influencing the *what, where, and how* of implementing WCH interventions
3. Priority predefined packages of WCH services are not commonly agreed on and implemented in conflict settings.
4. Working within the political and governance systems in conflict settings is increasingly challenging compared to previous decades, given the dynamic nature of modern conflict and the expanding role of Non State Armed groups.
5. The humanitarian system is creative and has developed new solutions to bring lifesaving WCH services closer to populations in very challenging environments.
6. Recognising and valuing the primary role of local actors would improve timely and appropriate WCH care delivery.

Introduction

Armed conflicts have disastrous effects on civilian populations. It is estimated that more than half a million civilians have been killed by combat operations in Syria alone between 2011 and 2019,² a significant number of whom were civilians (1126230) and amongst them women (13173 casualties), children and adolescents (21605 casualties under the age of 18). The toll from the extended, indirect effects of conflict due to the destruction of food supplies, roads, electricity and water infrastructure, and health facilities has also been catastrophic.^{3,4} In 2017, 701 attacks were reported on health facilities, health care staff, patients and ambulances in 23 conflict-affected countries.⁵ Armed conflicts have also negatively affected the number of forcibly displaced people in the world, increasing each year in the last decade, with 70.8 million people displaced by December 2018.⁶ New estimates from Bendavid et al. in this Series of the number of women and children affected by conflict – at least 630 million in 2017, including over 50 million women and children displaced by conflict – is, at over 8% of the world's population, strikingly large.^{7,8} This paper complements other Series papers by presenting empirical insights from a collection of ten country case studies aiming to assess the provision of WCH services in contemporary conflicts in Afghanistan, Colombia, Democratic Republic of Congo (DRC), Mali, Nigeria, Pakistan, Somalia, South Sudan, Syria and Yemen.

1

2 The key research question asked today is whether traditional humanitarian assistance has had
3 a positive effect on saving lives and mitigating morbidity and mortality in modern armed
4 conflicts. The implicit question is whether the humanitarian community has been able to
5 adapt to the changing nature of armed conflicts and respond to women's, children's and
6 adolescents' health needs.

7

8 Country case study teams comprised of local and international research partners often
9 supported by relief agencies. Their work was guided by a common research protocol (Panel
10 1),¹ with desk review parameters, quantitative analysis of national datasets (when available),

11 primary qualitative data collection tools and fieldwork strategies adapted to examine factors
12 influencing planning and implementation of WCH services in each setting. Detailed results
13 from each country are published elsewhere.¹ The present paper presents the synthesised
14 results and implications from the analysis of the ten country case studies.
15

Panel 1. Case study selection criteria and methodology

This Series paper presents insights from a collection of ten country case studies aiming to assess the provision of women's and children's health (WCH) services in contemporary conflicts in Afghanistan, Colombia, Democratic Republic of Congo (DRC), Mali, Nigeria, Pakistan, Somalia, South Sudan, Syria and Yemen. Countries were selected to ensure representation across geographies and conflict stages (e.g. acute, protracted, post-conflict).

Country case study teams comprised of local and international research partners often supported by humanitarian agencies. The teams' work was guided by a common research protocol,¹ with desk review parameters, quantitative analysis of national datasets (where available), primary qualitative data collection tools and fieldwork strategies adapted to examine factors influencing planning and implementation of WCH services in each setting..

We used a framework analysis approach to describe the coverage and spectrum of WCH interventions delivered, and to assess explanatory variables affecting variation in health service delivery. Our analysis focused on decision-making processes, obstacles to implementation of proven WCH interventions, and adaptation of service delivery strategies to address health needs of women, newborns, children and adolescents in varied geographic, political, economic and environmental conditions.

16
17

Social determinants affecting the health of women, newborns, children and adolescents

18 The effects of armed conflict are the combination of a number of risks factors including the
19 nature and exposure to conflict, the social determinants of health and the level of risks and
20 vulnerabilities experienced by women, newborns, children and adolescents. The social
21 determinants affecting their health in conflict settings include: lack of safe water and
22 sanitation; poor quality housing; poor nutrition; and lack of timely access to quality health
23 services; which in turn influence the health, opportunities for social and intellectual
24 development and quality of life of children and newborns even more, as they grow up in an
25 environment in which their ability to exercise their basic rights have deteriorated due to
26 immediate threats to security during occupation, fighting, etc.; experience of traumatic
27 events; and lack of opportunity to play as a way of developing social and motor skills.
28 Additionally, conflict and its attendant trauma often require that women undertake new social
29 and economic roles. Alternatively, women may become more vulnerable, if they are isolated
30 and exposed to violence and lack of resources. Women and adolescent girls are more
31 commonly exposed to sexual and gender-based violence including rape, which is often used
32 as a weapon of war.
33

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61

The nature of contemporary armed conflicts: analysing security attributes of the ten case study countries

Humanitarian actors are confronted today by increasingly complex armed conflicts. As analysed by Wise *et al.* in this Series,⁹ each conflict possesses its own unique character and history, and the impact of each conflict on civilian populations is rooted in complex political, strategic, and military determinants. Derived and expanded from the conceptual framework presented in Wise *et al.*,⁹ **Table 1** presents selected attributes of the case study conflicts related to the nature of warfare and the strategies and tactics of the engaged state and non-state combatant groups.

All the country case study conflicts are both intra-state, often labelled “civil” wars, and also internationalised, inter-state wars or conflict initiated by Non State Armed Groups (NSAGs) operating internationally. While most of the studied conflict settings are primarily rural in nature, Syria and Yemen have experienced destructive urban sieges, with the large-scale use of high explosives, including from airstrikes and artillery, in densely populated areas.^{10,11} It is challenging to summarise the organisational structure, strategies and tactics utilised by the various combatant groups, as the number of these groups are large and vary over time. However, it is useful to distinguish generally between the strategic attempts of belligerents to gain political legitimacy among civilian populations or to coerce civilian compliance through direct attacks or the deprivation of essentials of life or access to humanitarian assistance, and more specifically to essential WCH services.

In all ten countries where the conflict is characterised by a multitude of NSAGs, the access to populations that humanitarian actors have achieved has mostly been a result of humanitarian negotiations with parties to the conflict. Faced with increasingly complex dynamics in armed conflict as shown in **Table 1** below, humanitarian actors have increased investments in guidance, skills and capacities to operate in high-risk and access-constrained environments.¹²

62 **Table 1.** Case study country classification according to attributes of conflict

	Conflict setting*	Total Population (World Bank, 2018¹³)	Population in need of humanitarian assistance (OCHA, 2020¹⁴)	Battle-related deaths 1989-2018 (Uppsala Conflict Data Program)	Combatants: Number of Non State Armed Groups⁹
Afghanistan	Rural/Urban	37.2 million	9.4 million	227,510	8+ Main parties include Afghan Armed Forces, North Atlantic Treaty Organization (NATO) Resolute Support, US Forces Afghanistan, Taliban; also have Al Qaeda, Islamic State of Iraq and Syria (ISIS), Haqqani network, militia groups
Colombia	Rural	49.6 million	5.1 million	27,617	8+ Main parties include Colombian military, Fuerzas Armadas Revolucionarias de Colombia (FARC), Ejército de Liberación Nacional (ELN); and Autodefensas Gaitanistas de Colombia (AGC), Rastrojos, Aguilas Negras, Puntilleros, Ejército Popular de Liberación (EPL)
Democratic Republic of Congo	Rural	84.1 million	15.9 million	112,327	100+ Main parties are Forces Armées de la République Démocratique du Congo (FARDC), Mouvement du 23 mars (M23), Forces démocratiques de libération du Rwanda (FDLR), Union pour la Réhabilitation de la Démocratie du Congo (URDC), Mission de l'Organisation des Nations Unies pour la stabilisation en République démocratique du Congo (MONUSCO), Nduma défense du Congo-Rénové (NDC-R), Union Paysanne pour le Développement Intégral (UPDI), Forces démocratiques alliées/National Army for the Liberation of Uganda (ADF/NALU), Lord's Resistance Army (LRA), Forces nationales de libération (FNL), Alliance des patriotes pour un Congo libre et souverain (ALPCS) Kamuina Nsapu, Bana Mura, Mai Mai Mazembe, Force de résistance patriotique d'Ituri (FRPI), other Mai Mai militias and ethnic factions, and Burundian antigovernment militias.
Mali	Rural	19.1 Million	3.6 million	5,886	10+ Main parties are Military of Mali, African-led International Support Mission to Mali (AFISMA), Mouvement National pour la Libération de l'Azawad (MNLA), Ganda Iso, Front de libération nationale de l'Azawad (FLNA), Mouvement pour

					le salut de l'Azawad (MSA), Groupe d'Autodéfense Tuareg Imghad et Alliés (GATIA), Al-Qaeda in the Islamic Maghreb (AQIM), ISIS, Boko Haram
Nigeria	Rural	195.9 million	7.7 million + 284,843 refugees in region	54,538	12+ Main parties include Northeast actors: Nigerian military, Multinational Joint Task Force (MNJTF), Civilian Joint Task Force (CJTF), Boko Haram, ISIS West Africa, hunters, dan banga, other militias, AQIM; Middle Belt: Fulani and Hausa herders, Tiv and Tarok farmers
Pakistan	Rural/Urban	212.2 million	2.9 million	41,562	14+ Main parties are Pakistani military/Inter-Services Intelligence (ISI), TalibanHaqqani network, al Qaeda, Lashkar-e-Taiba, Balochistan Liberation Army (BLA), Harkat-ul-Jihad al-Islami, Jaish-e-Mohammed, Hizbul Mujahideen, Harkat-ul-Mujahideen, Al-Badr, Islamic State of Iraq and the Levant – Khorasan Province (ISIL-KP), Islamic Movement of Uzbekistan, other Sunni militant factions
Somalia	Rural	15 million	5.2 million	48,009	7+ Main parties are Somali armed forces, Al-Shabaab, African Union Mission to Somalia (AMISOM), Kenyan troops, United States (US) counterterror operations, ISIS, Al Qaeda
South Sudan	Rural	10.9 million	7.5 million + 2.3 million refugees in region	11,104	40+ Main parties are Sudan People's Liberation Army (SPLA), United Nations Mission in South Sudan (UNMISS), Mathiang Anyoor, Maban Defence Force, outh Sudan Liberation Movement (SSLM), Justice and Equality Movement (JEM), Sudan People's Liberation Movement-North (SPLM-N), Ethiopian Unity Patriots Front (EUPF), Sudan People's Liberation Movement-in-Opposition (SPLM-IO), Nuer White Army, and many others as it is estimated that South Sudan has at least 40 armed groups. ¹⁵
Syria	Urban	16.9 million	11 million+ 5.6 million refugees in region	345,079	15+ Main parties are Syrian Arab Army, Russia, National Defense Force, Shabiha, Foreign Shia militias including Hezbollah, Free Syrian Army, Jabhat Tahrir Souriya, ISIS, Hayat Tahrir al-Sham (formerly Nusra), US-led coalition, Syrian Democratic Forces, People's Protection Units (YPG), Turkey, Israel
Yemen	Urban	28.5 million	24 million	26,230	11+

					Main parties are Saudi-led Coalition, Houthi forces, pro-Hadi security forces, Saleh loyalists, ISIS, Al-Qaeda in the Arabian Peninsula (AQAP), Southern Movement; supporting groups include Alliance of Yemeni Tribes, Pro-Houthi Popular Committees, Special Security Forces, Tihamah resistance, other alleged groups
--	--	--	--	--	--

63

64

***Legend:**

65

Rural = conflict setting is primarily rural in nature

66

Urban = conflict setting is primarily urban in nature

67 **When conflicts undermine primary health care delivery**

68 Our analyses found no clear patterns on WCH intervention delivery in conflict settings. In
69 Afghanistan and Pakistan, our analysis suggested a statistically significant difference in
70 coverage of various WCH interventions between severely/moderately and minimally
71 conflict-affected provinces/districts based on the battle-related deaths.^{16,17} In Colombia,
72 maternal mortality, antenatal care coverage, caesarean section rate, and fertility in 15 to 19
73 year-old adolescents were significantly different in municipalities with high versus low levels
74 of conflict (measured as victimisation rates), and no statistically significant difference was
75 found in vaccination coverage, neonatal, early neonatal and infant mortality rates between
76 high and low conflict quintiles.¹⁸ In Nigeria, although there were differences in various
77 indicators including vaccination coverage and neonatal and child mortality among the
78 conflict-affected, marginal and stable (non-conflict) areas, these differences were not
79 statistically significant. In DRC, insecurity (measured as conflict-related fatality rate) had a
80 significant effect on maternal death and stillbirth rates, while the impact on coverage of
81 selected WCH interventions was not significant.¹⁹

82

83 In many cases, health services may exist on paper but have ceased to be delivered to people
84 in the specified catchment area. For example, in Afghanistan, comprehensive emergency
85 obstetric and newborn care (CEmONC) is largely not delivered in most provinces. Similarly,
86 the introduction of user fees at public and private health facilities has become standard
87 practice, as has been seen in Afghanistan (at tertiary level) and southern Sudan. An
88 overriding concern in Yemen and Syria has been how to evacuate victims of violence from
89 the site of an incident to the nearest emergency medical centre, an issue that was regularly
90 raised in the media because of its huge humanitarian, social and political dimension (personal
91 communication). Health systems in conflict settings can, as in “normal” situations, support a
92 healthy life, or, by their absence or ineffectiveness, undermine it and perpetuate health
93 inequity.

94

95 Data from the case study countries show that the presence of armed conflict has attracted the
96 attention, and even the intervention, of the international community, with evidence of
97 mitigation of the adverse health impact of conflicts and sometimes even greater health and
98 healthcare improvements that lead to important improvements in population health beyond
99 the pre-crisis levels (e.g. DRC, Nigeria, Somalia, South Sudan).¹⁹⁻²¹ Thanks to the
100 humanitarian system, many maternal and child health and nutrition services are now offered
101 to a majority of those who can be reached in conflict settings.

102

103 **Pre-conflict capacity of health systems as a determinant of WCH priorities**

104 There were differences among countries in terms of what services were delivered and how.
105 These differences can be attributed both to the intensity and nature of the conflict (e.g.
106 whether active, protracted, cyclical) and the capacity of the health system prior to the
107 conflict. For example, Syria was a middle-income country prior to the conflict with a
108 functioning health system providing free-at-the-point-of-delivery primary health care
109 services. Participants in our study reported that during the war, there were certain gaps in
110 WCH service availability over time and by geographical locations. In areas where the health

111 system was affected the most by the conflict, there was a delay in re-establishing WCH
112 service delivery. Prior to the conflict, Colombia also had all the infrastructures in place to
113 deliver WCH services, which were all delivered even during the conflict. Since the conflict in
114 Colombia spanned about 50 years, the capacity of its health system (e.g. infrastructure and
115 personnel) was in part developed in the midst of the conflict. All the other case study
116 countries reported long standing limitations in their health system capacity, at minimum in
117 the area where the conflict was occurring and experienced a state of protracted conflict,
118 which limited the reconstruction of capacities of these countries to deliver the wide spectrum
119 of WCH interventions.

120

121 The period of armed conflict that we studied cannot be disconnected from either the status
122 antebellum or from what comes (or is to come) after. In situations like Syria, the destruction
123 caused by armed conflict devastated the ability of whichever authorities are in charge to
124 maintain the health status of the population at a high level. However, in some case study
125 countries (e.g. Nigeria, Somalia), the situation was very different prior to the eruption of
126 conflict. In Nigeria, for example, it would be difficult to claim that the exacerbation of
127 conflict in the Northeast in 2009 had a major impact on routine health service delivery when
128 the 2008 Demographic Health Survey recorded a vaccination coverage in Borno State of 8%,
129 which is far below the national average.²² Moreover, there are countries where central
130 Governments have been purposely neglecting the conflict-affected parts of the country, e.g.
131 the weaknesses in Nigeria may, in fact, contribute to some causes for the origin of the
132 conflict. Weak health systems in conflict-affected parts of the country are not just “innocent
133 bystanders,” but rather a symptom of longstanding prejudicial policies. This might not be the
134 case in Syria, where the national health system was generally strong, but in countries like
135 Mali, Nigeria and Afghanistan there has been longstanding neglect, which leads to a situation
136 where, with intervention from the humanitarian community, it is possible that the health
137 status of the population will counter-intuitively improve during the conflict, only to
138 deteriorate once again when there is either peace or when the humanitarian community is no
139 longer providing assistance at the same level.

140

141 **Prioritising among WCH interventions: who decides?**

142 *Priority setting: an unclear process*

143 As it stands, the prioritisation of WCH interventions is not very clear. Priority in all the case
144 study settings is given to a set of specific interventions: antenatal care, BEmONC and
145 CEmONC for pregnant women and immediate care for newborns; childhood immunisation,
146 treatment of common childhood illnesses, infant and young child feeding (IYCF) and
147 malnutrition screening and treatment through in-patient, outpatient and stabilisation centres,
148 as these were considered life-saving interventions. Immunisation is a clear priority in
149 humanitarian response albeit implementation barriers vary across case study settings,
150 including limited humanitarian access to populations, lack of infrastructure for cold chain
151 maintenance and community reluctance. In Afghanistan and Pakistan, we also found a
152 specific focus on polio campaigns, which has been prioritised due to funding opportunities
153 rather than based on life saving grounds.

154

155 On the other hand, there is a set of interventions that is neglected in most countries: abortion
156 and post-abortion care, as well as provision of contraception were not prioritised by the
157 implementing stakeholders, particularly in countries where religious and cultural practises
158 affected the acceptability of such services like in Afghanistan, Mali, Nigeria, Pakistan,
159 Somalia and Yemen. Policy and political environments also influenced the provision of these
160 services, e.g. in Colombia, family planning interventions were mainly restricted to urban
161 areas, whereas DRC continues to enforce a law that prohibits the sale and use of
162 contraceptive methods for young people and adolescents. Adolescent health was also another
163 area that was alarmingly largely ignored, with the majority of the case studies reporting no
164 evidence of such implementation, with the exception of international humanitarian
165 organisations-led reproductive health programmes - although with a very limited scope - for
166 adolescents in DRC and Somalia.

167

168 Ten years ago, the call from Hurst et al. for “accountability for reasonableness” in the
169 humanitarian sector²³ has not been fully heard. What we observed in the ten countries is that
170 decisions are rather the fruit of a negotiation process between the international organisations
171 and the national authorities but also between the humanitarian organisations themselves.
172 Beyond the politics of humanitarian aid, the driver for interventional implementation remains
173 the access to the right resources: financial resources from international donors and expertise
174 from national and international organisations. Respondents in most country case study
175 settings described that the priorities of donors are the primary drivers of influencing the what,
176 where, and how of implementing interventions. Although the central government in some
177 countries (e.g. Colombia, Nigeria, and Afghanistan) is more actively involved in managing
178 and overseeing services, often the involvement in healthcare delivery of government officials
179 shrinks as conflict escalates in terms of scale and intensity. We found that decisions and the
180 ability to implement humanitarian WCH interventions are not uniquely based on needs
181 assessments and security situation, but also on the availability of local and international
182 actors on the ground who can rapidly deploy, access population groups and monitor the
183 quality and coverage of WCH interventions. Going forward, a rational prioritization process
184 in conflict settings could follow the logic of determining the epidemiological burden of the
185 problem that the intervention seeks to allay; a consideration of the available cost-effective
186 interventions and actors to address those problems with a high epidemiological burden; and a
187 consideration of the context, including security risks encountered by communities to get
188 access to health services and health care workers to deliver health care, cultural factors, the
189 capacities of the health system, and issues outside of the health sector that might be of higher
190 priority than health sector issues. For example, this process was followed in 2019 in
191 Afghanistan to develop its Integrated Package of Essential Health Services.²⁴

192

193 To bring clarity to the situation, we have identified the existence of four different but not
194 mutually exclusive models of decision-making amongst governmental and humanitarian
195 actors, which all dictate the mode of relationship and operation (**Table 2**).

196

197 **Table 2.** Models of decision-making amongst humanitarian and governmental actors

Decision-making model	Description	Examples from case study countries
Centralised	The decision-making system on which interventions are prioritised is centralised at government level and often influenced by UN agencies and funders. As an illustration, participants in Pakistan report stringent government regulations to work in conflict areas and approvals for non-governmental organisations (NGOs) are particularly cumbersome, in particular to work in federally administered tribal areas (FATA).	Afghanistan, Colombia, Pakistan, Syria
Humanitarian actors-led	In some settings, UN agencies and other humanitarian actors lead the choice of interventions to be implemented and delivery of health services. However, we found in our case studies that the decision-making process within and across these UN and humanitarian agencies on which interventions to prioritise was variable and not following uniform guidelines.	Somalia, South Sudan, Yemen, Syria
Collaborative	Settings where decision-making is a collaborative endeavour shared between government and humanitarian actors for defining both which WCH interventions are delivered, and how. These collaborative relationships are not always fully balanced between actors in terms of power and technical capacity	Afghanistan, DRC, Nigeria, Mali, Syria
Gatekeeping	This model is characterised by other conflict settings where the ministry of health (MoH) and Non-State Armed Groups have little or no control or power over the technical content of interventions, but become gatekeepers to regulate who works in government and rebel-controlled areas and delivers which interventions.	Yemen

198

199 The four models are not mutually exclusive and vary over time and space. For example, in
200 the case of Syria, the decision-making model depends on which entity (i.e. government
201 versus non-governmental authorities) has the authority to deliver health services in a specific
202 geographic area, which is why two different models co-exist: the humanitarian actors-led
203 model in non-government-controlled areas; and collaborative model in government-
204 controlled areas. Our findings suggest that there are ongoing tensions between humanitarian
205 modes of delivery and national health systems.

206

207 **Tensions between the humanitarian system and the national health system**

208 The differences in terms of approach between different humanitarian actors or between
209 national authorities and humanitarian actors illustrate the unpredictability and uncertainty of
210 situations that require constant adaptability.

211

212 Many case study countries reported needing to frequently adapt their WCH interventions to
213 the escalation of insecurity in some parts of the country, the constant changes of the conflict
214 (e.g. nature, scale, movement of troops, nature and intention of belligerents) and the cost of
215 delivering WCH services in hard-to-reach locations. For example, air delivery is the only
216 means to resupply health facilities in some parts of South Sudan and Somalia. Humanitarian

217 actors reported making adaptations to rapidly respond to and anticipate situations that are
218 often unpredictable. These adaptations included pre-establishing partnerships between UN
219 agencies and NGOs with pre-defined roles and responsibilities to respond to population
220 movements in DRC; using mobile clinics in Afghanistan to access hard-to-reach populations
221 either due to conflict and/or terrain; and donors making emergency funds available in South
222 Sudan to pre-stock medical supplies to be able to rapidly respond to outbreaks (e.g. cholera)
223 or sudden escalation in violence. These modes of operations are based on agile management
224 mechanisms, which are most often not present in public service and create differences in the
225 mode of operation between mainstream health services and humanitarian services, that
226 sometimes create tensions.

227

228 The tensions between the mainstream health system and the humanitarian system are caused
229 not only by power imbalances and unequal access to resources between international and
230 national actors, but also by differences in intervention principles. For example, in DRC, there
231 are tensions between free health care promoted by humanitarian actors clashing with the user
232 fee policy of the MOH or in Yemen where health authorities expressed frustration at times by
233 impactful decisions unilaterally made by UN agencies in terms of implementation of key
234 interventions. There is no doubt that most national authorities and humanitarian agencies
235 have a common vision of providing care to the most vulnerable. However, their time horizon,
236 budget and scale vary between humanitarian agencies targeting special geographical areas
237 and national authorities managing the national space. They also vary by country, which the
238 four models of decision-making processes have identified.


239

240 **Strategies to deliver WCH services**

241 Humanitarian actors (local and international) and national authorities are confronted by
242 various obstacles to deliver WCH interventions, as explained earlier. For the purpose of this
243 study, we developed health²⁵ and humanitarian system building blocks (an adaptation of the
244 WHO health systems building blocks²⁵), and classified our findings according to the
245 following domains: leadership, governance and coordination; health financing, health
246 workforce; essential medicine and supplies; health service delivery; health information
247 systems and communication; community dynamics and sociocultural factors; and security.

248 **Figure 1** visualises the areas of bottleneck to delivering WCH interventions in the case study
249 countries, and **Table 3** provides solutions documented in our case studies.

250 **Figure 1. Areas of bottleneck to implementing sexual, reproductive, maternal, newborn, child and adolescent health and nutrition**
 251 **interventions in ten country case study settings**
 252



253

254 *Remote management*

255 Cooperation between different humanitarian actors and local authorities (including
256 government) took on a variety of forms including subcontracting to local organisations (e.g.
257 Afghanistan, Somalia, Yemen, Mali, Syria). For example, in Syria, multiple cooperation
258 strategies were used. Humanitarian actors worked with local NGOs while monitoring their
259 activity through telephone calls and cross-border visits from Syrian healthcare providers to
260 Gaziantep or Amman. Coordination between a variety of actors also led to improved data
261 collection in some cases as well as unconventional data collection e.g. e-health, telephones
262 and informants (e.g. Syria).

263

264 *The emergence of pooled funds*

265 Funding levels and conditions varied greatly between contexts. Unresponsive funding
266 mechanisms, political interference in services (e.g. the Mexico City Policy or “global gag
267 rule,” which blocks United States federal funding for NGOs that provide abortion counselling
268 or referrals, advocate to decriminalise abortion or expand abortion services), competition to
269 get access to funding, and delays in the release of funds contributed to gaps in funding.
270 Moreover, donors were reluctant to invest in infrastructure or operational costs (e.g. South
271 Sudan), and multi-year programs to address the root causes of insecurity (e.g. South Sudan,
272 DRC) and better respond to population needs during protracted crises. Corruption among
273 governments in some countries also affected how donors distributed funds and to whom,
274 which in many cases remains mainly concentrated in international organisations. Respondents
275 described several mechanisms to address funding shortages and being more responsive to
276 emergencies that included relying upon emergency pooled funds (e.g. Somalia), and other UN
277 organizations (e.g. Yemen).

278

279 *Local health workforce at the forefront*

280 As highlighted earlier, armed conflicts are usually an exacerbating factor of existing
281 weaknesses of national health systems but also an accelerating factor for health staff
282 displacement. Shortages of health care workers, in particular the limited availability of health
283 workers with certain qualifications and specific profiles (e.g. specialisation and gender) was a
284 key problem in all contexts. Specialists such as gynaecologists, obstetricians, surgeons,
285 paediatricians, and physiotherapists were often unavailable in most countries. Female
286 midwives and nurses were lacking in conflict-affected areas of countries like Afghanistan,
287 Pakistan, and Syria to enable women’s access to health services. Several strategies were used
288 to address the lack of health workers. These included increasing training programs for health
289 staff (e.g., Somalia, Yemen, South Sudan), task shifting or task sharing (e.g. DRC, Mali,
290 Somalia, Syria, Pakistan, Afghanistan, Yemen), and expanding the catchment area and
291 populations for which health workers were responsible (e.g. Pakistan). In Keich district in
292 Pakistan, for example, senior staff living centrally within the district would rotate trips to
293 remote areas every month for a week

294

295 Local hiring and partnership with local organisations was a strategy that was employed by
296 humanitarian agencies to address concerns surrounding health workforce, financing, and
297 security. Local health workers, given their connection to the communities, were more likely to

298 continue to work even when there were funding gaps and salary delays (e.g. Somalia,
299 Colombia). Local workforce also contributed to mitigate security threats (e.g. DRC,
300 Colombia), and their importance in communicating and establishing trust with the community
301 (e.g. Somalia, Colombia). Findings from Colombia, DRC and Somalia demonstrate the
302 importance of intrinsic motivation and a ‘sense of duty’ for health worker retention: national
303 health workers, particularly those who worked with children, felt a need to protect the future
304 of the country. Similarly, in South Sudan, health workers who stayed in insecure communities
305 did so out of a sense of duty to these communities.

306
307 Numerous forms of health delivery to address barriers included task shifting and task sharing;
308 hiring other types of health workers (e.g. community midwives, community health workers;
309 traditional healers, traditional birth attendants); using new modes of delivery (e.g. remote
310 management, technology such as WhatsApp or electronic clinical protocols, mobile clinics,
311 treatment posts, home visits); implementing packages of services (e.g. sexual and
312 reproductive health and family planning, gender-based violence centres providing delivery
313 care) or and addressing demand for services. For example in Syria, capacity building
314 programmes for midwives were implemented to address the population’s preference for home
315 births, in part due to reported feelings of insecurity inside hospitals which are often targeted in
316 attacks.

317 318 *Rebuilding trust in the community*

319 Culturally-situated beliefs and behaviours influenced acceptance of and access to health
320 services. The use of humanitarian assistance as a political tool can have resounding damaging
321 effects on the community’s perception of and trust in lifesaving WCH services. Specific
322 services such as immunisation and family planning encountered religious and cultural
323 oppositions in countries such as Pakistan, DRC, South Sudan, Nigeria and Yemen, while
324 gender rules about female mobility limited access to care (e.g. Pakistan, Afghanistan).
325 Moreover, the protracted nature of war and the politicisation of aid have fractured the
326 community’s trust in health service providers (e.g. in Afghanistan, DRC, Pakistan). For
327 example, the use of polio vaccinators to identify and target Osama Bin Laden in Pakistan has
328 created long-lasting resistance and reluctance from communities in regard to public health
329 campaigns in both Pakistan and Afghanistan. The recruitment of local staff and use of social
330 science methods in programmes to better understand community perceptions and expectations
331 (e.g. the role of women in the household, the decision bearer in a family regarding child
332 health or number of children a woman has, or the image of humanitarian organisations) and
333 shape humanitarian interventions has been valued in countries such as Afghanistan or
334 Pakistan.

335 336 *Insecurity: a key driver of WCH service delivery*

337 Facilities’ resource shortages already existing before the crisis were further exacerbated by
338 attacks, looting and lack of investments during the crisis (e.g. Mali, South Sudan, Yemen).
339 Vulnerable populations’ access to health care are further exacerbated by breaches of medical
340 neutrality, i.e. violations of the Fourth Geneva Convention, Article 18. Direct insecurity was a
341 huge disincentive for working in conflict-affected areas due to the elevated risk of targeted

342 threats, attacks on or kidnapping of health workers (e.g. Yemen, Afghanistan, Somalia, Syria,
 343 Colombia). The insecure situation forced health care workers to put in place contingency
 344 plans (e.g. reduced movement and presence of health staff, generating patient evacuation
 345 plans, remote management). In Colombia, health personnel in some zones were trained about
 346 duties and rights of their medical mission and security plans to reduce their personal security
 347 risk. In Syria, remote management from certain hubs (e.g. a cross-border one in Gaziantep),
 348 was used to improve accessibility to certain geographic areas when no physical access was
 349 possible.²⁶ Respondents cited the importance of local partners to provide intelligence about
 350 security threats (e.g. Mali, Colombia). Many international respondents highlighted that they
 351 relied more heavily on local and national staff and partners to deliver services (e.g. DRC,
 352 Somalia, Syria, Yemen).^{19,20,26,27} In some cases, insecurity necessitated the negotiation with
 353 non-state armed groups to protect their personnel (e.g. Mali, Somalia, South Sudan,
 354 Colombia).^{18,20,21,28} For respondents in Syria and Afghanistan, such negotiations with
 355 opposition parties allowed access to restricted geographies during a a polio campaign.^{16,26}
 356

357 **Table 3.** Solutions to deliver sexual, reproductive, maternal, newborn, child and adolescent
 358 helath and nutrition interventions in ten country case studies

Health and humanitarian building blocks	Solutions identified
Governance, leadership, coordination	<ul style="list-style-type: none"> - Political analysis on power balance between the various warring parties and the various humanitarian actors - Various roles taken by the Health Cluster based on level of engagement and capacity of national authorities in the humanitarian response - Decentralisation of operations by contracting local organisations
Health financing	<ul style="list-style-type: none"> - Creation of multi-year funding mechanisms to respond to protracted crises - Creation of emergency pooled funds to respond to emergencies such as outbreaks or sudden displacement of populations
Health workforce	<ul style="list-style-type: none"> - Task-shifting and task-sharing - Rotation of senior staff to remote areas - Hiring local staff to nurture trust with local communities and value the “sense of duty” of local staff vis-à-vis their country
Essential medicine and supplies	<ul style="list-style-type: none"> - Creation of electronic stock management and supply information system to automate the identification of shortage and need for resupply
Health service delivery	<ul style="list-style-type: none"> - Use of mobile clinics to deliver services to remote areas - Recruitment of lay workers who have good knowledge of their community - Promotion of community-based services to bring services closer to populations - Delivery of integrated packages of services at the point of care to avoid populations to move several times
Community dynamics and sociocultural factors	<ul style="list-style-type: none"> - Social research used as way of informing how to deliver humanitarian programmes
Insecurity	<ul style="list-style-type: none"> - Training of health staff on security measures - Remote management - Use of security intelligence to assess the situation and authorise staff movement - Contextually-driven negotiations with non-state armed groups (NSAGs) to have access to populations and protect populations and health staff

359

360 **Conclusion**

361 Working within the political and governance systems in modern conflict settings is
362 increasingly challenging given the dynamic nature of modern conflict and the expanding role
363 of Non-State Armed groups who are often in control of large geographic areas, which pose
364 new challenges to delivering services to women, children and adolescents.

365
366 Decision-making processes vary by government, organisation and context. We categorised
367 them into four different models (centralised; humanitarian actors-led; collaborative;
368 gatekeeping) that are not mutually exclusive. We also found that pre-conflict health system
369 capacity, and thus readiness to respond in unpredictable events, varied greatly by case study
370 setting. Important enabling factors included the income status of countries (e.g. middle-
371 income versus low-income), existing infrastructure and resilience of the health system.

372
373 Despite large variations in sociocultural and geographical contexts and decision-making
374 processes, there was consistency among prioritisation of key WCH interventions (i.e.
375 antenatal care, BEmONC, CEmONC, immunisation, treatment of common childhood
376 illnesses, IYCF and malnutrition treatment and screening) as well as the neglect of other such
377 interventions (sexual, reproductive, newborn and adolescent health, and those for stillbirths).
378 Given the dynamic nature of modern conflicts, proactively defining and agreeing upon an
379 evidence-based comprehensive WCH package of interventions, taking into account existing
380 evidence and guidelines (e.g. Inter-Agency Field Manual on Reproductive Health in Crises²⁹,
381 Newborn Health in Humanitarian Settings Field Guide³⁰) could be an effective strategy to
382 respond to sudden situations. The prioritisation process, in a context of scarce resources,
383 could be an effective approach to identify the key interventions to implement and decide on
384 the allocation of resources.

385
386 The humanitarian system is creative and has developed new solutions to bring lifesaving
387 WCH services closer to populations by hiring and training other types of health workers,
388 often from the affected community and using new modes of delivery. These solutions, when
389 rigorously evaluated, can represent a concrete, timely response to current implementation
390 challenges and remind health authorities of their responsibility to deliver basic health services
391 to the whole population.

392
393 **Contributors**

394 Neha S. Singh and Karl Blanchet conceived the analysis and all authors contributed to the
395 final structure and content of the paper. Neha S. Singh and Karl Blanchet led the overall data
396 analysis, and Anushka Ataullahjan, Zahra Ahmed, Chaza Akik, Chiara Altare, Jai K. Das,
397 Shafiq Mirzazada, Khadidiatou Ndiaye, Samira Sami, Paul Spiegel, Hannah Tappis, and Paul
398 H.Wise made contributions to specific sections. Neha S. Singh wrote the first draft of the
399 paper and all authors contributed to critical interpretation of the results and development of
400 the paper. All authors saw and approved the final version.

401
402 **Acknowledgments**

403 We would like to thank all study respondents who took part in the key informant interviews,
 404 and the BRANCH consortium members who contributed to specific country case studies: (1)
 405 Afghanistan (Wardah Ahmed, Zulfiqar A Bhutta, Jai K. Das, Malika Fatima, Zahra Feroz,
 406 Sultana Jabeen, Shafiq Mirzazada, Mohammed I. Zubair); (2) Colombia (Gladis Adriana
 407 Vélez Álvarez, Edison Bedoya Bedoya, Jewel Gausman, Isabel C. Garcés-Palacio, Sara
 408 Milena Ramos Jaraba, Marly Andrea García Jiménez, Ana Langer, Laura Ruiz Sánchez,
 409 María Ochoa Sierra, Natalia Quiceno Toro, Ivan Dario Arroyave Zuluaga); (3) DRC (Chiara
 410 Altare, Hamady Ba, Stéphane Muzindusi Bikoro, Ghislain Bisimwa, Ties Boerma,
 411 Christopher Hook, Espoir Bwenge Malembaka, Jerome Pfaffmann, Thea Scognamiglio, Paul
 412 Spiegel, Hannah Tappis, Maphie Tosha); (4) Mali (Anushka Ataullahjan, Diego Bassani,
 413 Samba Diarra, Seydou Doumbia, Michelle F. Gaffey, Moctar Tounkara); (5) Nigeria (Eskedar
 414 Dejene, Khadidiatou Ndiaye, Ume Nwanneamaka, Chinwenwo Weli, Christina Okere,
 415 Victory Inyang, Jennifer Tyndall, Ron Waldman); (6) Pakistan (Wardah Ahmed, Zulfiqar A.
 416 Bhutta, Jai K. Das, Malika Fatima, Zahra Feroz, Sultana Jabeen); (7) Somalia (Zahra Ahmed,
 417 Anushka Ataullahjan, Abdirizak Dalmar, Michelle Gaffey, Mohamed Osman, Chantal
 418 Umutoni); (8) South Sudan (Ties Boerma, Nicole Lightman, Augustino Mayai, Wilfred
 419 Ochan, Samira Sami, Grace Sheehy, Paul Spiegel, Hannah Tappis, James Wanyama, Hannah
 420 Wild); (9) Syria (Chaza Akik, Linda Shaker Berbari, Karl Blanchet, Joanne Constantin,
 421 Jocelyn DeJong, Abdulkarim Ekzayez, Hala Ghattas, Zeina Jamaluddine, Katherine Lopes,
 422 Ghada Saad-Haddad, Aline Semaan, Neha S. Singh); and (10) Yemen (Iman Ahmed Saleh
 423 Al-Gawfi, Eman Al-Kubati, Nagiba A. Abdulghani AlShawafi, Huda BaSaleem, Sarah
 424 Elaraby, Shatha Elnakib, Fouad Othman, Nuzhat Rafique, Fouzia Shafique, Paul Spiegel,
 425 Hannah Tappis).

426
 427 Funding for the country case studies was provided through a subgrant from the Centre for
 428 Global Child Health at the Hospital for Sick Children (SickKids), with travel and meeting
 429 costs supported directly by Aga Khan University and the Partnership for Maternal, Newborn
 430 and Child Health (PMNCH). As coordinator of the BRANCH Consortium (Bridging Research
 431 & Action in Conflict Settings for the Health of Women & Children), the SickKids Centre for
 432 Global Child Health has received funding for BRANCH research activities from the
 433 International Development Research Centre (IDRC) (108416-002 & 108640-001), the
 434 Norwegian Agency for Development Cooperation (Norad) (QZA-16/0395), the Bill &
 435 Melinda Gates Foundation (OPP1171560), and UNICEF (PCA 20181204). Aga Khan
 436 University received funding for BRANCH activities from the Family Larsson-Rosenquist
 437 Foundation.

438
 439 **BRANCH Consortium Steering Committee Members**
 440

Author order on Pubmed	First name and middle initials	Surnames
1	Zulfiqar A.	Bhutta
2	Robert E.	Black

3	Karl	Blanchet
4	Ties	Boerma
5	Michelle F.	Gaffey
6	Ana	Langer
7	Paul B.	Spiegel
8	Ronald J.	Waldman
9	Paul H.	Wise

441

442 Author details:

443 Prof. Zulfiqar A Bhutta

444 *Centre for Global Child Health, The Hospital for Sick Children, Toronto, Canada;*

445 *Dalla Lana School of Public Health, University of Toronto, Toronto, Canada;*

446 *Centre of Excellence in Women and Child Health, Aga Khan University, Karachi, Pakistan*

447

448 Prof. Robert E Black

449 *Department of International Health, Johns Hopkins Bloomberg School of Public Health,*

450 *Baltimore, USA*

451

452 Prof. Karl Blanchet

453 *The Geneva Centre of Humanitarian Studies, University of Geneva, Graduate Institute,*

454 *Geneva, Switzerland;*

455 *Health in Humanitarian Crises Centre, London School of Hygiene and Tropical Medicine,*

456 *London, UK*

457

458 Prof. Ties Boerma

459 *Centre for Global Public Health, University of Manitoba, Winnipeg, Canada*

460

461 Michelle F Gaffey

462 *Centre for Global Child Health, The Hospital for Sick Children, Toronto, Canada;*

463 *Dalla Lana School of Public Health, University of Toronto, Toronto, Canada*

464

465 Prof. Ana Langer

466 *Women and Health Initiative; Department of Global Health and Population, Harvard T.H.*

467 *Chan School of Public Health, Boston, Massachusetts, USA*

468

469 Prof. Paul B Spiegel

470 *Center for Humanitarian Health, Johns Hopkins University, Baltimore, USA*

471

472 Prof. Ronald J Waldman

473 *Milken Institute School of Public Health, George Washington University, Washington, DC,*

474 *USA;*

475 *Doctors of the World USA, New York, NY, USA*

476

477 Prof. Paul H Wise

478 *Department of Pediatrics, Stanford University School of Medicine, Stanford, California, USA*

479 **References**

- 480 1. Atallahjan A, Gaffey MF, Sami S, et al. Investigating the delivery of health and
481 nutrition interventions for women and children in conflict settings: a collection of case studies
482 from the BRANCH Consortium. *Conflict and Health* 2020; **14**(1): 1-4.
- 483 2. Syrian observatory for Human Rights. More than 570 thousand people were killed on
484 the Syrian territory within 8 years of revolution demanding freedom, democracy, justice, and
485 equality. 2019. <http://www.syriahr.com/en/?p=120851>. Accessed on 29 October 2019.
- 486 3. Wise PH. The epidemiologic challenge to the conduct of just war: confronting indirect
487 civilian casualties of war. *Daedalus* 2017; **146**(1): 139-54.
- 488 4. Alda E, Mc Evoy C. Beyond the Battlefield: Towards a Better Assessment of the
489 Human Cost of Armed Conflict: Small Arms Survey; 2017.
- 490 5. Rubenstein L, Bales C, Spitzer W, Wille C. Violence on the frontline: Attacks on
491 Health Care in 2017, 2019.
- 492 6. UNHCR. Mid-year trends 2018. 2019.
493 <https://www.unhcr.org/afr/statistics/unhcrstats/5c52ea084/mid-year-trends-2018.html>
494 (accessed 15 March 2019).
- 495 7. Bendavid E, Boerma T, Akseer N, et al. The health effects of armed conflict on
496 women, children and adolescents. *The Lancet* 2021 (in press).
- 497 8. Peterson K. From the field: gender issues in disaster response and recovery. *Natural*
498 *Hazards Observer, Special Issue on Women and Disasters* 1997; **21**(5).
- 499 9. Wise PH, Shiel A, Southard N, et al. The political and security dimensions of the
500 humanitarian health response to violent conflict. *The Lancet* 2021 (in press).
- 501 10. Guha-Sapir D, Schlüter B, Rodriguez-Llanes JM, Lillywhite L, Hicks MH-R. Patterns
502 of civilian and child deaths due to war-related violence in Syria: a comparative analysis from
503 the Violation Documentation Center dataset, 2011–16. *The Lancet Global Health* 2018; **6**(1):
504 e103-e10.
- 505 11. Shield R. The Saudi air war in Yemen: A case for coercive success through battlefield
506 denial. *Journal of Strategic Studies* 2018; **41**(3): 461-89.
- 507 12. Chaudhri S, Cordes K, Miller N. Humanitarian health programming and monitoring in
508 inaccessible conflict settings: a literature review. *Journal of International Humanitarian*
509 *Action* 2019; **4**(1): 9.
- 510 13. World Bank. Total population database.
511 <https://data.worldbank.org/indicator/sp.pop.totl> (accessed 30 April 2020).
- 512 14. United Nations Office for the Coordination of Humanitarian Affairs (OCHA). Global
513 Humanitarian Overview 2020. Accessed on 30 April 2020 2019: Available from:
514 <https://www.unocha.org/global-humanitarian-overview-2020>.
- 515 15. Irin News. Who's who: South Sudan's splintering war. 2018.
516 <https://www.irinnews.org/analysis/2018/04/12/who-s-who-south-sudan-s-splintering-civil-war>
517 (accessed 15 March 2019).
- 518 16. Mirzazada S, Padhani ZA, Jabeen S, et al. Impact of conflict on maternal and child
519 health service delivery: a country case study of Afghanistan. *Conflict and Health* 2020; **14**(1):
520 1-13.

- 521 17. Das JK, Padhani ZA, Jabeen S, et al. Impact of conflict on maternal and child health
522 service delivery—how and how not: a country case study of conflict affected areas of Pakistan.
523 *Conflict and Health* 2020; **14**: 1-16.
- 524 18. Ramos Jaraba SM, Quiceno Toro N, Ochoa Sierra M, et al. Health in conflict and
525 post-conflict settings: reproductive, maternal and child health in Colombia. *Conflict and*
526 *Health* 2020; **14**: 1-21.
- 527 19. Altare C, Malembaka EB, Tosha M, et al. Health services for women, children and
528 adolescents in conflict affected settings: experience from North and South Kivu, Democratic
529 Republic of Congo. *Conflict and health* 2020; **14**: 1-19.
- 530 20. Ahmed Z, Atallahjan A, Gaffey MF, et al. Understanding the factors affecting the
531 humanitarian health and nutrition response for women and children in Somalia since 2000: a
532 case study. *Conflict and Health* 2020; **14**: 1-15.
- 533 21. Sami S, Mayai A, Sheehy G, et al. Maternal and child health service delivery in
534 conflict-affected settings: a case study example from Upper Nile and Unity states, South
535 Sudan. *Conflict and Health* 2020; **14**: 1-12.
- 536 22. Abuja N. National Population Commission and ICF Macro; 2009. *National*
537 *Population Commission (NPC)[Nigeria] and ICF Macro Nigeria Demographic and Health*
538 *Survey* 2008.
- 539 23. Hurst SA, Mezger N, Mauron A. Allocating resources in humanitarian medicine.
540 *Public Health Ethics* 2009; **2**(1): 89-99.
- 541 24. Islamic Republic of Afghanistan Ministry of Public Health. Integrated Package of
542 Essential Health Services. Available at: [https://www.wslshmacuk/sites/default/files/2019-](https://www.wslshmacuk/sites/default/files/2019-09/IPHES_2019pdf)
543 [09/IPHES_2019pdf](https://www.wslshmacuk/sites/default/files/2019-09/IPHES_2019pdf) 2019; **Accessed on 22 November 2019**.
- 544 25. World Health Organization. Monitoring the Building Blocks of Health Systems: A
545 handbook of Indicators and their Measurement Strategies. Geneva: WHO, 2010.
- 546 26. Akik C, Semaan A, Shaker-Berbari L, et al. Responding to health needs of women,
547 children and adolescents within Syria during conflict: intervention coverage, challenges and
548 adaptations. *Conflict and Health* 2020; **14**(1): 1-19.
- 549 27. Tappis H, Elaraby S, Elnakib S, et al. Reproductive, maternal, newborn and child
550 health service delivery during conflict in Yemen: a case study. *Conflict and health* 2020; **14**:
551 1-16.
- 552 28. Atallahjan A, Gaffey MF, Tounkara M, et al. C'est vraiment compliqué: a case study
553 on the delivery of maternal and child health and nutrition interventions in the conflict-affected
554 regions of Mali. *Conflict and Health* 2020; **14**: 1-17.
- 555 29. Inter-Agency Working Group on Reproductive Health in Crises (IAWG). Inter-
556 Agency Field Manual on Reproductive Health In Humanitarian Settings. 2018.
- 557 30. Save the Children, WHO, UNICEF. Newborn Health in Humanitarian Settings: Field
558 guide. 2017.
559 [https://resourcecentre.savethechildren.net/sites/default/files/documents/newbornhealthbook-](https://resourcecentre.savethechildren.net/sites/default/files/documents/newbornhealthbook-production2017-v4b-press.pdf)
560 [production2017-v4b-press.pdf](https://resourcecentre.savethechildren.net/sites/default/files/documents/newbornhealthbook-production2017-v4b-press.pdf) (accessed 12 March 2019).

561
562