



Combining IP₃ affinity chromatography and bioinformatics reveals a novel protein-IP₃ binding site on *Plasmodium falciparum* MDR1 transporter

Eduardo Alves^a, Helder Nakaya^{b,c}, Euzébio Guimarães^d, Célia R.S. Garcia^{b,*}

^a Life Science Department, Imperial College London, London, United Kingdom

^b Department of Clinical and Toxicological Analyses of Pharmaceutical Sciences, University of São Paulo, São Paulo, Brazil

^c Computational Systems Biology Laboratory, INOVA, University of São Paulo, São Paulo, Brazil

^d Federal University of Rio Grande do Norte, Pharmacy Department, Health Science Center, Natal, Brazil

ARTICLE INFO

Keywords:

Plasmodium falciparum
IP₃ receptor
Malaria
Signaling
MDR transporter

ABSTRACT

Intracellular Ca²⁺ mobilization induced by second messenger IP₃ controls many cellular events in most of the eukaryotic groups. Despite the increasing evidence of IP₃-induced Ca²⁺ in apicomplexan parasites like *Plasmodium*, responsible for malaria infection, no protein with potential function as an IP₃-receptor has been identified. The use of bioinformatic analyses based on previously known sequences of IP₃-receptor failed to identify potential IP₃-receptor candidates in any *Apicomplexa*. In this work, we combine the biochemical approach of an IP₃ affinity chromatography column with bioinformatic meta-analyses to identify potential vital membrane proteins that present binding with IP₃ in *Plasmodium falciparum*. Our analyses reveal that PF3D7_0523000, a gene that codes a transport protein associated with multidrug resistance as a potential target for IP₃. This work provides a new insight for probing potential candidates for IP₃-receptor in *Apicomplexa*.

Introduction

The inositol 1,4,5-triphosphate (IP₃) is an important second messenger that regulates cytosolic Ca²⁺ in a variety of Eukaryotic organisms (Michell, 2011; Berridge, 2009). Briefly, the activation of phospholipase C (PLC) mediated by surface receptor breaks phosphatidylinositol 4,5-bisphosphate (PIP₂) into soluble short life second messenger IP₃ that binds into IP₃ receptor (IP₃R), culminating in intracellular Ca²⁺ release (Streb et al., 1983; Berridge and Irvine, 1984).

The phylum *Apicomplexa* includes unicellular eukaryotes parasites like *Plasmodium*, the etiology agent of malaria infection, and possesses the metabolic enzymes responsible for generation and degradation of IP₃, see review (Garcia et al., 2017). IP₃ can mobilize Ca²⁺ from intracellular stores in isolate and permeabilized blood-stage *P. chabaudi* (Passos and Garcia, 1998) and in intact *P. falciparum* within red blood cells (RBCs) (Alves et al., 2011). Within RBCs, parasites manage to maintain the Ca²⁺ stores full even under a low Ca²⁺ environment (Gazarini et al., 2003). An increasing number of reports supporting the existence of intracellular Ca²⁺ release induced by IP₃ in malaria parasites (Passos and Garcia, 1998; Alves et al., 2011; Beraldo et al., 2007; Martin et al., 1994; Enomoto et al., 2012; Raabe et al., 2011) suggest the existence of a Ca²⁺ channel sensitive to IP₃, the IP₃R.

The IP₃R is a well known described protein in vertebrates and contains around four to six transmembrane domains (TMDs), see review (Mikoshiba, 2007). Prole and Taylor (2011) used the sequence of mammal N-terminal IP₃R binding domain and the amino-terminal RIH (Ryanodine and IP₃R homology) domains to perform a BLAST (Basic Local Alignment Search Tool) on the genome of diverse parasites. However, this work failed to find any potential candidate for IP₃R in *Apicomplexa*. So far, no apicomplexan IP₃R candidate has been identified or suggested through bioinformatics approach. Moreover, there is no publication that attempted to use a biochemical approach like an IP₃ affinity chromatography column in *Apicomplexa* to identify proteins that might bind to IP₃.

Hirata and collaborators (Hirata et al., 1990) managed to enrich proteins from rat brain sample that has an affinity to IP₃, like IP₅-phosphatase and IP₃ 3-kinase using an analogous IP₃ affinity chromatography column 2-O-[4-(5-aminoethyl-2-hydroxyphenylazo) benzoyl]-1,4,5-tri-O-phosphono-myo-inositol trisodium salt-Sepharose 4B. Using a similar column, Kishigami and collaborators (Kishigami et al., 2001) managed to identify the PLC protein from octopus' eyes *Todarodes pacificus* and reported that squid rhodopsin also has an affinity to IP₃. Nevertheless, besides the potential of these columns to enrich proteins that bind to IP₃, no IP₃R has ever been identified using an

* Corresponding author.

E-mail address: cgarcia@usp.br (C.R.S. Garcia).

<https://doi.org/10.1016/j.crmicr.2022.100179>

IP₃-affinity column alone.

By adapting the protocol from Hirata/Kishigami (Hirata et al., 1990; Kishigami et al., 2001), we created a column containing IP₃ conjugated with biotin linked with a high-performance sepharose-streptavidin and challenged with proteins from asynchronous *P. falciparum* blood-stage. Using an IP₃-free column containing only sepharose-streptavidin as a reference, we selected only the candidates exclusive on the IP₃-column to undergo a series of bioinformatic meta-analyses. Our approach targeted candidates with at least one transmembrane domain, considered essential, conserved among most apicomplexan species, and with unknown or non-clear function. Finally, the candidate that fit all these criteria were used as targets for *in silico* molecular docking against IP₃.

Using this strategy, we identified the *P. falciparum* multidrug resistance protein 1 (PfMDR1) as a vital and conserved membrane protein that has the potential to bind to IP₃. This protein is located on the parasite food vacuole, a Ca²⁺ storage compartment (Biagini et al., 2003). Combined, our IP₃ affinity column and bioinformatic approach successfully narrow to provide the first small list of malaria proteins candidates with quintessential features expected from an IP₃R.

Material and methods

P. falciparum culture

P. falciparum (D37) parasites were maintained in culture as described (Trager and Jensen, 1976). Briefly, *P. falciparum* were cultured in RPMI media supplemented with 50 mg/L hypoxanthine; 40 mg/L gentamycin; 435 mg/L NaHCO₃; 2% hematocrit of A⁺ human red blood cells and 10% A⁺ human blood serum in an atmosphere of 5% CO₂; 3% O₂; 92% N₂ at 37 °C. Media was changed every 24 h and RBCs replaced every 48 h. Parasitemia and the development stage of cultures were determined by Giemsa-stained smears.

P. falciparum protein sample

Total *P. falciparum* protein extract was obtained from 2.5 L of unsynchronized culture, at 8% parasitemia. The culture was washed three times in PBS (300 g, 5 min) and parasites isolated from erythrocytes using 0.03% (w/v) saponin (Sigma) on PBS containing protease inhibitors: antipain, pepstatin, chymostatin, and leupeptin (Sigma) at concentrations of 20 µg/mL each and 500 µM benzamide (Sigma). Isolate parasites were centrifuge on 1300 g for 10 min at 4 °C and washed three times in PBS with protease inhibitors. The isolate parasite samples were resuspended in 50 mM TRIS-HCl buffer pH 7.4 containing 2 mM EDTA, 0.1% Triton X-100, protease inhibitors, and 1 mM PMSF. Samples were sonicated on SONIC (Vibracell) 50% potency for 20 s for 3 times on ice (10 s interval between each sonication) follow by a 1300 g centrifugation for 10 min at 4 °C for removal of the insoluble pellet. DNase and RNase (final concentration 200 ng/µL each) were added on soluble pellet and incubated for one hour at 37 °C. The samples were passed through a 0.45 µm filter. The amount of protein was quantitated using Pierce's BCA protein assay kit.

IP₃-affinity chromatography column

For the column, it was used a commercial high performance Sepharose substrate bound to streptavidin (GE Healthcare Life Science) and biotin-conjugated IP₃ (Echelom Biosciences). The streptavidin-sepharose column was equilibrated by washing once with 10x volume of ice-cold, 0.45 µm filter binding buffer (20 mM NaH₂PO₄, 150 mM NaCl, 20 mM LiCl and 2 mM EDTA, pH 7.5). The columns were mounted in a 15 mL sterile falcon tube. For each column it was used 1.25 ml of equilibrated Sepharose-streptavidin resuspended in binding buffer mixed with 20 µg of IP₃-biotin. The columns were left by constant stirring for 12 h at 4 °C in a dark environment and then centrifuged for 1 min, 300 g at 4 °C. The supernatant containing excess IP₃-biotin was

removed and columns were washed five times with 2 ml of ice-cold binding buffer to remove any free IP₃-biotin. Two distinct columns were assembled: one containing IP₃-biotin-sepharose-streptavidin and other containing only sepharose-streptavidin. In each column was loaded with 2.5 mg of *P. falciparum* protein extract and the volume was adjusted with ice-cold binding buffer with protease inhibitors until a final volume of 5 ml. The columns were incubated at 4 °C under gentle, steady shaking in a light-protected environment for 12 h and finally centrifuged for 1 min at 300 g at 4 °C to discard the supernatant. Each column was washed seven times with ice-bound binding buffer with protease inhibitors. To elute the proteins, 1 mL of an ice-cold elution buffer (8 M Guanidin-HCl, 20 mM LiCl, 2 mM EDTA, pH: 1.5 with protease inhibitors) was added on each column followed by constant stirring for 1 h at 4 °C. At the end of incubation, the columns were centrifuged for 1 min at 300 g, and the supernatant was collected in sterile low binding protein Eppendorf.

Mass spectrometry

The protein samples were applied on 8% polyacrylamide gel and run at low voltage (60 v) until the bands were discriminated. After the run, the gel was fixed and stained following the recommendations of the "Colloidal Blue Staining Kit" from Invitrogen. The gel sections containing visible bands were cut and sent for analysis on a mass spectrometer at Taplin Mass Spectrometry, Harvard Medical School (<https://taplin.med.harvard.edu/>) for protein identification. All identified proteins containing at least one exclusive peptide match were considered for analyses.

Transmembrane domain prediction

To detect a transmembrane domain's presence, the whole amino acid sequence from the protein identified at mass spectrometry was analysed using the public HMMTOP program version 2.0 (www.enzim.hu/hmmtop/). This program predicts the number of transmembrane helices and their position from the peptide/protein amino acid sequence.

Phenotype score, conservation, and function predictions

The phenotype score used the determinate gene essentiality of each candidate was obtained from the work of Zhang et al. (2018) available on PlasmoDB (<https://plasmodb.org/plasmo>). To identify orthologs candidates among the *Apicomplexa* group, we use the OrthoMCL database (<https://orthomcl.org/orthomcl>). For function prediction, we consulted the gene annotation information provided by PlasmoDB.

In silico docking with IP₃

The primary sequence of MDR1 (Gene - PF3D7_0523000, plasmodb.org) was used to build its probable 3D structure by homology modeling. The server SwissModel (Schwede et al., 2003) was employed to automatically create the models optimized to bind IP₃ at various locations inside MDR1 homology. Blind molecular docking simulations were carried out to obtain possible interactions for the intermembrane domain as predicted by the TMHMM Server (Krogh et al., 2001). The SwissDock (Grosdidier et al., 2011) server enabled the study of IP₃ intermembrane MDR1 domain binding poses. Additionally, the IP₃-Ion-MDR1 binding was further investigated using the multidrug transporter permeability (P)-glycoprotein is adenosine triphosphate (ATP)-binding cassette (PDB id: 6COV). The later ability to bind simultaneously ATP and a divalent cation at the intracellular domain was used to guide the inspect a hypothetical IP₃-Ion-MDR1 interaction. IP₃ was manually positioned inside the ATP cavity to mimic an IP₃-Mg²⁺ interaction. The binding conformation was optimized with molecular mechanics employing the UCSF (Pettersen et al., 2004) chimera minimize structure tools.

Protein-protein interaction network

Using *Plasmodium* interactome data (Hillier et al., 2019), we looked for the proteins that interact with the MDR1. The protein annotation and functions were also retrieved from the original publication. The network was generated using Cytoscape (Shannon et al., 2003).

Results

IP₃-affinity chromatography data

Adapting the protocol based on Hirata/Mishigami (Hirata et al., 1990; Kishigami et al., 2001), we use an IP₃ affinity chromatography column with protein homogenate from unsynchronized asexual blood stages of isolated *P. falciparum* as the first step to identified potential proteins that have a similar function to IP₃R receptor in a mammal (Fig. 1).

The access code of the brute data on mass spectrometry analyses from the eluate samples of the IP₃-affinity chromatography column can be found in Supplemental Material Table 1. At least 695 proteins from *P. falciparum* containing at least one exclusive peptide were detected from the IP₃-sepharose column. In comparison, 494 proteins were detected from the sepharose matrix alone (Fig. 1). All proteins exclusively present on IP₃-sepharose were selected (total 201 proteins) for the bioinformatic meta-analyses (Sup. Table 2).

Once the proteins exclusive for IP₃-column were identified (Sup. Table 2), the first bioinformatic approach aimed to select proteins that contain at least one transmembrane domain (TMD). The TMD is an important structure to anchor proteins through biological membranes by its physical properties like the length and hydrophilicity of the transmembrane span (Cosson et al., 2013), every IP₃R in vertebrates, invertebrates and single eukaryotes organism possess a TMDs, so we used this feature as the second step to select potential candidates for IP₃R. Fig. 1.

Table 1 summarizes 26 proteins exclusively found at IP₃-biotin-streptavidin-sepharose column containing at least one TMDs. Transfection of *P. falciparum* to constitutively express IP₃-sponge, a protein containing a modified IP₃ binding domain based on mouse IP₃R that sequesters cytosolic IP₃ (Usui-Aoki et al., 2005), did not result in viable parasites (Pecenin et al., 2018) suggesting a vital role of IP₃ signaling in *P. falciparum*. Accordingly, the next step to narrow the number of potential candidates that might act as IP₃R in malaria is to focus on essential genes. To deem whether a gene is essential, we considered only the candidates that scored lower than 0.5 on its mutagenic index of phenotype graphic (data provided by PlasmoDB). That decreases the number of candidates to 11 (Fig. 1, Table 1).

There is pharmacological evidence of the IP₃R in multiple

Table 1

The table I: The list of 26 proteins exclusively found at IP₃-biotin-streptavidin-sepharose column that contains at least one TMDs.

Gene code (PlasmoDB)	Predicted function/ annotation	Number TMDs	Essential gene
PF3D7_1001500	Early transcribed membrane protein 10.1	2	Yes
PF3D7_0501300	Skeleton-binding protein 1	1	No
PF3D7_1133400	Apical membrane antigen 1	1	No
PF3D7_0827900	Protein disulfide-isomerase	1	Yes
PF3D7_0918000	Glideosome-associated protein 50	2	No
PF3D7_1364100	6-cysteine protein P92	1	No
PF3D7_0523000	Multidrug resistance protein 1	11	Yes
PF3D7_0202500	Early transcribed membrane protein 2	1	No
PF3D7_0817500	Histidine triad nucleotide-binding protein 1	1	No
PF3D7_0402100	<i>Plasmodium</i> exported protein (PHISTb), unknown function	1	No
PF3D7_0501200	Parasite-infected erythrocyte surface protein	3	No
PF3D7_0501100	Heat shock protein 40, type II	1	yes
PF3D7_1252100	Rhoptry neck protein 3	3	Yes
PF3D7_1237700	Conserved protein, unknown function	5	Yes
PF3D7_0801800	Mannose-6-phosphate isomerase, putative	1	No
PF3D7_0731300	<i>Plasmodium</i> exported protein (PHISTb), unknown function	1	Yes
PF3D7_0702500	<i>Plasmodium</i> exported protein, unknown function	2	No
PF3D7_1344800	Aspartate carbamoyltransferase	1	Yes
PF3D7_1332600	DNA-(apurinic or apyrimidinic site) lyase 1	1	No
PF3D7_1105300	Conserved <i>Plasmodium</i> protein, unknow function	1	Yes
PF3D7_1038000.1	Antigen UB05	2	Yes
PF3D7_1016900	Early transcribed membrane protein 10.3	2	No
PF3D7_1002100	EMP1-trafficking protein	1	No
PF3D7_1476600	<i>Plasmodium</i> exported protein, unknown function	1	No
PF3D7_1458100	Protein PET117, putative	1	No
PF3D7_0508000	6-cysteine protein	1	Yes

apicomplexan parasites (Garcia et al., 2017), so in our analyses we considered only conserved genes among multiples species within the *Apicomplexa* phylum as the fourth step for candidate screening. Only four essential candidates with TMDs domains met this criterion: multidrug resistance protein 1 (MDR1); a heat shock protein 40, type II (HSP40); aspartate carbamoyltransferase (ATCase), and antigen UB05. PlasmoDB access code: PF3D7_0523000, PF3D7_0501100, PF3D7_1344800 and PF3D7_1038000 respectively. Among these 4




Fig. 1. Schematic approached to pinpoint potential candidates for IP₃R from isolates blood stage *P. falciparum*. 1° step: selection of proteins that are exclusively found on IP₃-Biotin-streptavidin-sepharose column. 2° step: selection of proteins that contains at least one transmembrane domain. 3° step: selection of proteins that are considered essential for malaria parasite during red blood stage development. 4° step: Selection of proteins that are conserve among most species within *Apicomplexa* group. 5° step: selection of protein with unknow or non-specific metabolic function. 6° step: candidates with positive in-silico docking against IP₃.

candidates, only MDR1 and antigen UB05 has unknown or unclear function. The HSP40 is a cochaperone protein with conserved J-domain that regulates other heat shock protein 70 (HSP70) (Walsh et al., 2004), and the ATCase is an enzyme important for the pyrimidine biosynthetic pathway (Simmmer et al., 1990). The MDR1 was the only candidate with information available to build a 3D structure by homology modeling to perform an in-silico binding with IP₃.

IP₃-MDR1 binding modeling and protein interactions network

The MDR1 model provided by the SwissModel server proved to be quite similar to the human P-glycoprotein ABCB1 receptor, protein data bank id: 7A69 (Nosol et al., 2020). The sequence alignment proved that a homology model could be built with fair quality with an identity of 29.7% and similarity of 48.2% (Pairwise Sequence Alignment EMBOSS Water server, <https://www.ebi.ac.uk/Tools/emboss/>). Two binding positions at the transmembrane domain of MDR1 and IP₃ binding was estimated by the SwissDock server (Fig. 2).

The pocket 1 (binding energy -15.7 kcal/mol) proved to be the best IP₃ docking position. The site is a lysine-rich domain able to form various hydrogen bonds with IP₃. The second-best binding pocket proved to be less favored as derived from the lower interaction energy (-11.4 kcal/mol). Another binding possibility investigated was the interaction with the same pocket ATP binding. The interaction involves the presence of a divalent cation (green spheres) like Mg²⁺ intercalating with IP₃. The MDR1 is an ATP-binding cassette (ABC) transporter family member associated with multidrug drug resistance due to translocating

amphiphilic compounds (Koenderink et al., 2010). The translocation of a substrate across the membrane by proteins like *P. falciparum* MDR1 requires an ATP binding on Q-loop site that causes a rearrangement of TM (Jones et al., 2009). The binding on IP₃-divalent cation on the MDR1 Q-loop site suggests a potential competition between ATP and IP₃. Interestingly, ATP is known to allosterically modulate the functional of mammal IP₃R (Ferris et al., 1990; Bezprozvanny and Ehrlich, 1993) including the inhibition of Ca²⁺ flux regulated by IP₃R under a high concentration of ATP (Bezprozvanny and Ehrlich, 1993).

To help uncover the cellular function of MDR1 protein, we searched for proteins that interact with MDR1 in the *Plasmodium* interactome data (Hillier et al., 2019) (Fig. 3). The data suggests that MDR1 interacts with activated C kinase receptors (RACK1, PF3D7_1148000). The *Pf*RACK1 can inhibit host IP₃-mediated Ca²⁺ signaling by direct interaction with IP₃R (Sartorello et al., 2009). The interaction with eukaryotic translation initiation factor 2 (EIF2, PF3D7_1410600), EIF2β (PF3D7_1010600), EIF2γ (PF3D7_1410600) and serine/threonine protein kinase (PF3D7_1148000) suggests that *Pf*MDR1 can associate or have similar functions to other receptors and nuclear factors that coordinate signaling events regulated by protein kinase.

Discussion and conclusion

Phylogenetic analyses and comparative genomic data revealed both unique and conserved proteins related to calcium signaling pathways on apicomplexan parasites (Prole and Taylor, 2011; Nagamune and Sibley, 2006; Ladenburger et al., 2009), nevertheless, the IP₃R still remains a




Fig. 2. Schematic representation of the Swiss-Dock most energetically favored binding poses of IP₃-MDR1. The cytosolic nucleotide-binding domain (upper part) display an IP₃ associated with divalent cation (green spheres) interacting on the same ATP binding pocket. The transmembrane domains (lower part) display two possible pocket sites on IP₃ interaction and their respective interaction energy values (kcal/mol). On the right side, details of the IP₃-MDR1 pocket 1 interaction, a region rich on lysine..




Fig. 3. Proteins that interact with MDR1. The network shows the proteins (gray nodes) that interact with MDR1 (red node) according to the *Plasmodium* interactome data (Hillier et al., 2019). Gene codes: 2270.T00246, translation initiation factor (PF3D7_0607000). SNRPD1, small nuclear ribonucleoprotein Sm D1 (PF3D7_1125500). ABCE1, ABC transporter E family member 1 (PF3D7_1,368,200). PRP2, pre-mRNA-splicing factor ATP-dependent RNA helicase PRP2 (PF3D7_1231600). PF11_0488, serine/threonine protein kinase (PF3D7_1148000). DDX1, ATP-dependent RNA helicase DDX1 (PF3D7_0521700). SEC13, protein transport protein SEC13 (PF3D7_1230700). SEC24A, transport protein Sec24A (PF3D7_1361100). EIF2GAMMA, eukaryotic translation initiation factor 2 subunit gamma (PF3D7_1410600). RON6, rhopty neck protein 6 (PF3D7_0214900). RACK1, receptor for activated C kinase (PF3D7_0826700). MDR2, multidrug resistance protein 2 (PF3D7_11447900). 2270.T00294, ATP-dependent RNA helicase MTR4, (PF3D7_0602100). DBP5, ATP-dependent RNA helicase DBP5 (PF3D7_1459000). SEC12, guanine nucleotide-exchange factor (PF3D7_1116400). WDR26, WD repeat-containing protein 26 (PF3D7_0518600). CBP20, nuclear cap-binding protein subunit 2 (PF3D7_0415500). PRPF19, pre-mRNA-processing factor 19 (PF3D7_0308600). EIF2BETA, eukaryotic translation initiation factor 2 subunit beta (PF3D7_1010600). 2270.T00096, cleavage stimulation factor subunit 1 (PF3D7_0620500).

major missing piece of this Ca^{2+} signaling toolkit. Studies using exogenous IP_3 on malaria parasites support the existence of protein sensitive to IP_3 that is capable to trigger a Ca^{2+} response (Passos and Garcia, 1998; Alves et al., 2011). The constant failed to identify this protein in *Apicomplexa* suggests this group has a distinct and unique structure compared to the IP_3 -binding core domain from other eukaryotes. The search for an IP_3 R in *Apicomplexa* requires a different strategy that does not rely exclusively on bioinformatics tools as BLAST (Basic Local Alignment Search Tool) based on previously known IP_3 R.

The use of IP_3 affinity chromatography column has been successfully reported to concentrate proteins that interact with high affinity to IP_3 analogues (Hirata et al., 1990) and retained key components from IP_3 — Ca^{2+} signaling from proteins extract from tissues (Kishigami et al., 2001). In this work, we used a biotin-inositol 1,4,5-triphosphate attached to a high-performance streptavidin-sepharose substrate to initially enriching proteins with IP_3 affinity from unsynchronized isolate *P. falciparum* blood culture. One of the significant limitations of using chromatography affinity column based on a short life IP_3 molecule is the number of naturally present proteins at sample homogenate that degrade this second messenger. *P. falciparum* contains proteins that can dephosphorylate or phosphorylate IP_3 like inositol polyphosphate 5-phosphatase and inositol 1,4,5-trisphosphate 3-kinase (Gardner et al., 2002). In this protocol, we tried to overcome this limitation by keeping all the binding and elution steps under low temperature while adding LiCl in every buffer. LiCl has been previously used to inhibit the dephosphorylation of IP_3 (Elabbadi et al., 1994; Irvine et al., 1985; Thomas et al., 1984). Another risk of using an IP_3 affinity column assumes that protein(s) that might interact with IP_3 in *Plasmodium* do not bind/interact with strong affinity with the sepharose-streptavidin substrate alone. We excluded all 494 proteins that bind with the sepharose-free IP_3 column as a potential IP_3 R candidate (Fig. 1).

From the 201 proteins identified exclusively from the IP_3 -sepharose column, 175 did not contain any TMDs suggesting the protocol used to extract the proteins from parasite lysate benefited mostly soluble proteins that do not strongly interact with lipid bilayers. This protocol can be optimized for future trials by using a protein extraction that targets membrane proteins (MPs). The IP_3 R in mammals is an MP protein

containing 6 TMDs (Joseph, 1996). The presence of TMDs is an essential aspect of any MPs to physically interact with biological membranes (Cosson et al., 2013). It is fair to predict that any protein with the potential function of IP_3 R should have TMDs to interact with membranes. Table 1 list all candidates with TMDs exclusively from IP_3 -column.

Ca^{2+} is a second messenger that regulates a variety of vital functions in apicomplexan parasites (Nagamune and Sibley, 2006; Docampo et al., 2014; Budu and Garcia, 2012). Accordingly, the use of 2-aminoethoxydiphenyl borinate (2-APB), a pharmacological drug that inhibits IP_3 R, abolished spontaneous Ca^{2+} mobilization and compromise intracellular development of blood stage *P. falciparum* (Enomoto et al., 2012). Pecenin and collaborator (Pecenin et al., 2018) failed to obtain any viable parasite expressing IP_3 -sponge. These data suggest that the IP_3 — Ca^{2+} signaling pathway has a vital role during intraerythrocytic development of *P. falciparum* and support our hypotheses that a potential candidate for IP_3 R in *Plasmodium* not only has to present a TMDs, but also has to be essential. A prediction of gene essentiality in *P. falciparum*, based on the work of Zhang and collaborators (Zhang et al., 2018) is available for consultation at the PlasmoDB website.

The pharmacological evidence that supports the IP_3 — Ca^{2+} signaling pathway in the *Apicomplexa* group is not exclusive to malaria parasites but also present in *Toxoplasma gondii* (Lourido and Moreno, 2015; Chini et al., 2005; Lovett et al., 2002) and *Babesia bovis* (Florin-Christensen et al., 2000). The strategy to pinpoint the potential candidate for IP_3 R in apicomplexan should not rely on gene only exclusive to *Plasmodium* species. Adding this extra meta-analysis step, the list of potential candidates presented exclusively on the IP_3 -sepharose column is finally reduced to four proteins: a MDR1; HSP40; an ATCase, and antigen UB05. Among those four, only MDR1 and antigen UB05 currently have an undefined function.

The small number of candidates makes the use of more computational demanding bioinformatic analyses more feasible. A molecular docking allows us to target the structural protein complexes from our candidate list against potential ligand as IP_3 or other potent IP_3 -analogues drugs like adenophostin A (Mak et al., 2001).

Molecular docking on IP_3 on *P. falciparum* MDR1 protein revealed two potential binding sites on TMD: pocket site 1 (binding energy -15.7

kcal/mol) and pocket site 2 (−11.4 kcal/mol), see Fig. 2. This data suggests that MDR1 pocket 1 has a higher affinity to IP₃ compared to IP₃-binding core of mammal IP₃R ($\Delta G = -10.3$ kcal/mol on 23 °C) (Ding et al., 2010) and a lower affinity when compare to IP₃-binding with N-terminal region of mammal IP₃R ($\Delta G = -79.5$ kcal/mol) (Chandran et al., 2019). Nevertheless, the binding of ATP on Q-Loop site on the nucleotide-binding domain (NBD) likely causes profound changes in the TMD region (Jones et al., 2009) making it hard to predict the actual affinity of the MDR1 protein with IP₃.

In *P. falciparum*, the MDR1 gene encodes for a 162.2 kg Daltons P-glycoprotein located on the digestive vacuole (DV) (Cowman et al., 1991) with unclear function. Still, the polymorphisms within this protein are associated with increases *in vitro* resistance against multiple antimalarial drugs like quinine (Sidhu et al., 2002; Sidhu et al., 2006; Sanchez et al., 2008; Basco et al., 1995; Reed et al., 2000; Cowman et al., 1994; Duraisingh et al., 2000; Price et al., 2004). The MDR1 displays a role as a transporter protein that brings solutes into DV. It consists of two distinct homologous regions: one cytosolic nucleotide-binding domain (NBD) and a substrate-binding consisting of 11 TMDs (Friedrich et al., 2014; Rohrbach et al., 2006). Interestingly, in malaria parasites, the DV is an acid compartment known to be a dynamic intracellular Ca²⁺ store (Biagini et al., 2003; Garcia et al., 1998; Borges-Pereira et al., 2020; Varotti et al., 2003), making the subcellular location of MDR1 protein suitable for an IP₃R-like candidate. Moreover, the *in vivo* and *in vitro* treatment with IP₃R inhibitor 2-aminoethoxydiphenyl borinate (2-APB) is associated with reversing resistance to antimalarial chloroquine in *P. falciparum* and *P. chabaudi* parasites, presumably by disrupting Ca²⁺ homeostasis (Mossaad et al., 2015). Multiple antimalarial drugs can also disrupt the Ca²⁺ dynamic on the parasite (Lee et al., 2018; Gazarini et al., 2007), nevertheless, there is no direct evidence that suggests the MDR1 acts as a Ca²⁺ gate.

The lack of information to build a quality 3D model for in-silico analyses on UB05, HSP40 and ATCase candidates does not exclude them as a potential role in sensing IP₃. The next natural step is to obtain functional evidence that these four candidates act as a protein sensitive to IP₃. One suggestion is expressing them on a triple IP₃R knock-out cell lines like DT40 chicken B cell (Winding and Berchtold, 2001) and test its sensitivity to mobilize Ca²⁺ with IP₃.

Considering that agents that disrupt IP₃R channels such as 2-APB block malaria *in vitro* growth (Beraldo et al., 2007; Enomoto et al., 2012; Pecenin et al., 2018), identify this receptor in *Plasmodium* will not only add crucial missing information on malaria Ca²⁺ signaling, but it will also present a potential new target for pharmacological treatment. This work aims to stimulate the use of IP₃-affinity column with bioinformatic strategies as a potential tool to identify proteins that might act as IP₃R in *Apicomplexa*. The MDR1 seems to be a promising candidate waiting to be validated. Nevertheless, this is just an initial but an important first step from a long rewarding task of finding the *Apicomplexa* channel sensitive to IP₃.

Funding

Celia R. S. Garcia is funded by FAPESP (2017/08684-7; 2018/07177-7). Helder Nakaya is funded by FAPESP/2018/14933-2.

Author contribution

All authors have contributed to discuss experimental design, discussing the data and manuscript writing. EA; EG and HN performed experiments.

Declaration of Competing Interest

The authors declare no conflict of financial or commercial interests.

Data availability

Data will be made available on request.

Acknowledgements

We thank Prof. Dr Akio Kishigami for the helpful suggestion on the IP₃-Affinity Column. Ross Tomaino from Taplin Mass Spectrometry Facility for helpful support on mass spectrometry data. Colsan (Associação Beneficente de Coleta de Sangue) for providing the human blood and plasma used on parasite culture.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.crmicr.2022.100179.

References

- Alves, E., Bartlett, P.J., Garcia, C.R., Thomas, A.P., 2011. Melatonin and IP₃-induced Ca²⁺ + release from intracellular stores in the malaria parasite *Plasmodium falciparum* within infected red blood cells. *J. Biol. Chem.* 286 (7), 5905–5912.
- Basco, L.K., Le Bras, J., Rhoades, Z., Wilson, C.M., 1995. Analysis of pfmdr1 and drug susceptibility in fresh isolates of *Plasmodium falciparum* from sub-Saharan Africa. *Mol. Biochem. Parasitol.* 74 (2), 157–166.
- Beraldo, F.H., Mikoshiba, K., Garcia, C.R., 2007. Human malarial parasite, *Plasmodium falciparum*, displays capacitative calcium entry: 2-aminoethyl diphenylborinate blocks the signal transduction pathway of melatonin action on the *P. falciparum* cell cycle. *J. Pineal Res.* 43 (4), 360–364.
- Berridge, M.J., 2009. Inositol trisphosphate and calcium signalling mechanisms. *Biochim. Biophys. Acta* 1793 (6), 933–940.
- Berridge, M.J., Irvine, R.F., 1984. Inositol trisphosphate, a novel second messenger in cellular signal transduction. *Nature* 312 (5992), 315–321.
- Bezprozvanny, I., Ehrlich, B.E., 1993. ATP modulates the function of inositol 1,4,5-trisphosphate-gated channels at two sites. *Neuron* 10 (6), 1175–1184.
- Biagini, G.A., Bray, P.G., Spiller, D.G., White, M.R., Ward, S.A., 2003. The digestive food vacuole of the malaria parasite is a dynamic intracellular Ca²⁺ store. *J. Biol. Chem.* 278 (30), 27910–27915.
- Borges-Pereira, L., Thomas, S.J., Dos Anjos, E.S.A.L., Bartlett, P.J., Thomas, A.P., Garcia, C.R.S., 2020. The genetic Ca(2+) sensor GCaMP3 reveals multiple Ca(2+) stores differentially coupled to Ca(2+) entry in the human malaria parasite *Plasmodium falciparum*. *J. Biol. Chem.* 295 (44), 14998–15012.
- Budu, A., Garcia, C.R., 2012. Generation of second messengers in *Plasmodium*. *Microbes Infect.* 14 (10), 787–795.
- Chandran, A., Chee, X., Prole, D.L., Rahman, T., 2019. Exploration of inositol 1,4,5-trisphosphate (IP₃) regulated dynamics of N-terminal domain of IP₃ receptor reveals early phase molecular events during receptor activation. *Sci. Rep.* 9 (1), 2454.
- Chini, E.N., Nagamune, K., Wetzel, D.M., Sibley, L.D., 2005. Evidence that the cADPR signalling pathway controls calcium-mediated microneme secretion in *Toxoplasma gondii*. *Biochem. J.* 389 (2), 269–277. Pt.
- Cosson, P., Perrin, J., Bonifacino, J.S., 2013. Anchors aweigh: protein localization and transport mediated by transmembrane domains. *Trends Cell Biol.* 23 (10), 511–517.
- Cowman, A.F., Galatis, D., Thompson, J.K., 1994. Selection for mefloquine resistance in *Plasmodium falciparum* is linked to amplification of the pfmdr1 gene and cross-resistance to halofantrine and quinine. *Proc. Natl. Acad. Sci. U. S. A.* 91 (3), 1143–1147.
- Cowman, A.F., Karcz, S., Galatis, D., Culvenor, J.G., 1991. A P-glycoprotein homologue of *Plasmodium falciparum* is localized on the digestive vacuole. *J. Cell Biol.* 113 (5), 1033–1042.
- Ding, Z., Rossi, A.M., Riley, A.M., Rahman, T., Potter, B.V., Taylor, C.W., 2010. Binding of inositol 1,4,5-trisphosphate (IP₃) and adenophostin A to the N-terminal region of the IP₃ receptor: thermodynamic analysis using fluorescence polarization with a novel IP₃ receptor ligand. *Mol. Pharmacol.* 77 (6), 995–1004.
- Docampo, R., Moreno, S.N., Plattner, H., 2014. Intracellular calcium channels in protozoa. *Eur. J. Pharmacol.* 739, 4–18.
- Duraisingh, M.T., Jones, P., Sambou, I., von Seidlein, L., Pinder, M., Warhurst, D.C., 2000. The tyrosine-86 allele of the pfmdr1 gene of *Plasmodium falciparum* is associated with increased sensitivity to the anti-malarials mefloquine and artemisinin. *Mol. Biochem. Parasitol.* 108 (1), 13–23.
- Elabbadi, N., Ancelin, M.L., Vial, H.J., 1994. Characterization of phosphatidylinositol synthase and evidence of a polyphosphoinositide cycle in *Plasmodium*-infected erythrocytes. *Mol. Biochem. Parasitol.* 63 (2), 179–192.
- Enomoto, M., Kawazu, S., Kawai, S., Furuyama, W., Ikegami, T., Watanabe, J., et al., 2012. Blockage of spontaneous Ca²⁺ oscillation causes cell death in intraerythrocytic *Plasmodium falciparum*. *PLoS ONE* 7 (7), e39499.
- Ferris, C.D., Haganir, R.L., Snyder, S.H., 1990. Calcium flux mediated by purified inositol 1,4,5-trisphosphate receptor in reconstituted lipid vesicles is allosterically regulated by adenine nucleotides. *Proc. Natl. Acad. Sci. U. S. A.* 87 (6), 2147–2151.

- Florin-Christensen, J., Suarez, C.E., Florin-Christensen, M., Hines, S.A., McElwain, T.F., Palmer, G.H., 2000. Phosphatidylcholine formation is the predominant lipid biosynthetic event in the hemoparasite *Babesia bovis*. *Mol. Biochem. Parasitol.* 106 (1), 147–156.
- Friedrich, O., Reiling, S.J., Wunderlich, J., Rohrbach, P., 2014. Assessment of *Plasmodium falciparum* PfMDR1 transport rates using Fluo-4. *J. Cell Mol. Med.* 18 (9), 1851–1862.
- Garcia, C.R., Ann, S.E., Tavares, E.S., Dluzewski, A.R., Mason, W.T., Paiva, F.B., 1998. Acidic calcium pools in intraerythrocytic malaria parasites. *Eur. J. Cell Biol.* 76 (2), 133–138.
- Garcia, C.R.S., Alves, E., Pereira, P.H.S., Bartlett, P.J., Thomas, A.P., Mikoshiba, K., et al., 2017. InsP3 signaling in apicomplexan parasites. *Curr. Top. Med. Chem.* 17 (19), 2158–2165.
- Gardner, M.J., Hall, N., Fung, E., White, O., Berriman, M., Hyman, R.W., et al., 2002. Genome sequence of the human malaria parasite *Plasmodium falciparum*. *Nature* 419 (6906), 498–511.
- Gazarini, M.L., Sigolo, C.A., Markus, R.P., Thomas, A.P., Garcia, C.R., 2007. Antimalarial drugs disrupt ion homeostasis in malarial parasites. *Mem. Inst. Oswaldo Cruz* 102 (3), 329–334.
- Gazarini, M.L., Thomas, A.P., Pozzan, T., Garcia, C.R., 2003. Calcium signaling in a low calcium environment: how the intracellular malaria parasite solves the problem. *J. Cell Biol.* 161 (1), 103–110.
- Grosdidier, A., Zoete, V., Michielin, O., 2011. SwissDock, a protein-small molecule docking web service based on EADock DSS. *Nucleic Acids Res.* 39, W270–W277. Web Server issue.
- Hillier, C., Pardo, M., Yu, L., Bushell, E., Sanderson, T., Metcalf, T., et al., 2019. Landscape of the *Plasmodium* interactome reveals both conserved and species-specific functionality. *Cell Rep.* 28 (6), 1635–1647 e5.
- Hirata, M., Yanaga, F., Koga, T., Ogasawara, T., Watanabe, Y., Ozaki, S., 1990. Stereospecific recognition of inositol 1,4,5-trisphosphate analogs by the phosphatase, kinase, and binding proteins. *J. Biol. Chem.* 265 (15), 8404–8407.
- Irvine, R.F., Anggard, E.E., Letcher, A.J., Downes, C.P., 1985. Metabolism of inositol 1,4,5-trisphosphate and inositol 1,3,4-trisphosphate in rat parotid glands. *Biochem. J.* 229 (2), 505–511.
- Jones, P.M., O'Mara, M.L., George, A.M., 2009. ABC transporters: a riddle wrapped in a mystery inside an enigma. *Trends Biochem. Sci.* 34 (10), 520–531.
- Joseph, S.K., 1996. The inositol triphosphate receptor family. *Cell Signal* 8 (1), 1–7.
- Kishigami, A., Ogasawara, T., Watanabe, Y., Hirata, M., Maeda, T., Hayashi, F., et al., 2001. Inositol-1,4,5-trisphosphate-binding proteins controlling the phototransduction cascade of invertebrate visual cells. *J. Exp. Biol.* 204 (3), 487–493. Pt.
- Koenderink, J.B., Kavishe, R.A., Rijpma, S.R., Russel, F.G., 2010. The ABCs of multidrug resistance in malaria. *Trends Parasitol.* 26 (9), 440–446.
- Krogh, A., Larsson, B., von Heijne, G., Sonnhammer, E.L., 2001. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* 305 (3), 567–580.
- Ladenburger, E.M., Sehring, I.M., Korn, I., Plattner, H., 2009. Novel types of Ca²⁺ release channels participate in the secretory cycle of *Paramecium* cells. *Mol. Cell Biol.* 29 (13), 3605–3622.
- Lee, A.H., Dhingra, S.K., Lewis, I.A., Singh, M.K., Siriwardana, A., Dalal, S., et al., 2018. Evidence for regulation of hemoglobin metabolism and intracellular ionic flux by the *Plasmodium falciparum* chloroquine resistance transporter. *Sci. Rep.* 8 (1), 13578.
- Lourido, S., Moreno, S.N., 2015. The calcium signaling toolkit of the Apicomplexan parasites *Toxoplasma gondii* and *Plasmodium* spp. *Cell Calcium* 57 (3), 186–193.
- Lovett, J.L., Marchesini, N., Moreno, S.N., Sibley, L.D., 2002. *Toxoplasma gondii* microneme secretion involves intracellular Ca²⁺ release from inositol 1,4,5-trisphosphate (IP(3))/ryanodine-sensitive stores. *J. Biol. Chem.* 277 (29), 25870–25876.
- Mak, D.O., McBride, S., Foskett, J.K., 2001. ATP-dependent adenophostin activation of inositol 1,4,5-trisphosphate receptor channel gating: kinetic implications for the durations of calcium puffs in cells. *J. Gen. Physiol.* 117 (4), 299–314.
- Martin, S.K., Jett, M., Schneider, I., 1994. Correlation of phosphoinositide hydrolysis with exflagellation in the malaria microgametocyte. *J. Parasitol.* 80 (3), 371–378.
- Michell, R.H., 2011. Inositol and its derivatives: their evolution and functions. *Adv. Enzyme Regul.* 51 (1), 84–90.
- Mikoshiba, K., 2007. IP₃ receptor/Ca²⁺ channel: from discovery to new signaling concepts. *J. Neurochem.* 102 (5), 1426–1446.
- Mossaad, E., Furuyama, W., Enomoto, M., Kawai, S., Mikoshiba, K., Kawazu, S., 2015. Simultaneous administration of 2-aminoethyl diphenylborinate and chloroquine reverses chloroquine resistance in malaria parasites. *Antimicrob. Agents Chemother.* 59 (5), 2890–2892.
- Nagamune, K., Sibley, L.D., 2006. Comparative genomic and phylogenetic analyses of calcium ATPases and calcium-regulated proteins in the apicomplexa. *Mol. Biol. Evol.* 23 (8), 1613–1627.
- Nosol, K., Romane, K., Irobalieva, R.N., Alam, A., Kowal, J., Fujita, N., et al., 2020. Cryo-EM structures reveal distinct mechanisms of inhibition of the human multidrug transporter ABCB1. *Proc. Natl. Acad. Sci. U. S. A.* 117 (42), 26245–26253.
- Passos, A.P., Garcia, C.R., 1998. Inositol 1,4,5-trisphosphate induced Ca²⁺ release from chloroquine-sensitive and -insensitive intracellular stores in the intraerythrocytic stage of the malaria parasite *P. chabaudi*. *Biochem. Biophys. Res. Commun.* 245 (1), 155–160.
- Pecenin, M.F., Borges-Pereira, L., Levano-Garcia, J., Budu, A., Alves, E., Mikoshiba, K., et al., 2018. Blocking IP₃ signal transduction pathways inhibits melatonin-induced Ca²⁺ signals and impairs *P. falciparum* development and proliferation in erythrocytes. *Cell Calcium* 72, 81–90.
- Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., et al., 2004. UCSF Chimera—a visualization system for exploratory research and analysis. *J. Comput. Chem.* 25 (13), 1605–1612.
- Price, R.N., Uhlemann, A.C., Brockman, A., McGready, R., Ashley, E., Phaipun, L., et al., 2004. Mefloquine resistance in *Plasmodium falciparum* and increased pfmdr1 gene copy number. *Lancet* 364 (9432), 438–447.
- Prole, D.L., Taylor, C.W., 2011. Identification of intracellular and plasma membrane calcium channel homologues in pathogenic parasites. *PLoS ONE* 6 (10), e26218.
- Raabe, A.C., Wengelnik, K., Billker, O., Vial, H.J., 2011. Multiple roles for *Plasmodium berghei* phosphoinositide-specific phospholipase C in regulating gametocyte activation and differentiation. *Cell Microbiol.* 13 (7), 955–966.
- Reed, M.B., Saliba, K.J., Caruana, S.R., Kirk, K., Cowman, A.F., 2000. Pgh1 modulates sensitivity and resistance to multiple antimalarials in *Plasmodium falciparum*. *Nature* 403 (6772), 906–909.
- Rohrbach, P., Sanchez, C.P., Hayton, K., Friedrich, O., Patel, J., Sidhu, A.B., et al., 2006. Genetic linkage of pfmdr1 with food vacuolar solute import in *Plasmodium falciparum*. *EMBO J.* 25 (13), 3000–3011.
- Sanchez, C.P., Stein, W.D., Lanzer, M., 2008. Dissecting the components of quinine accumulation in *Plasmodium falciparum*. *Mol. Microbiol.* 67 (5), 1081–1093.
- Sartorello, R., Amaya, M.J., Nathanson, M.H., Garcia, C.R., 2009. The *Plasmodium* receptor for activated C kinase protein inhibits Ca²⁺ signaling in mammalian cells. *Biochem. Biophys. Res. Commun.* 389 (4), 586–592.
- Schwede, T., Kopp, J., Guex, N., Peitsch, M.C., 2003. SWISS-MODEL: an automated protein homology-modeling server. *Nucleic Acids Res.* 31 (13), 3381–3385.
- Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., et al., 2003. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* 13 (11), 2498–2504.
- Sidhu, A.B., Uhlemann, A.C., Valderramos, S.G., Valderramos, J.C., Krishna, S., Fidock, D.A., 2006. Decreasing pfmdr1 copy number in *Plasmodium falciparum* malaria heightens susceptibility to mefloquine, lumefantrine, halofantrine, quinine, and artemisinin. *J. Infect. Dis.* 194 (4), 528–535.
- Sidhu, A.B., Verdier-Pinard, D., Fidock, D.A., 2002. Chloroquine resistance in *Plasmodium falciparum* malaria parasites conferred by pfcr mutations. *Science* 298 (5591), 210–213.
- Simmer, J.P., Kelly, R.E., Rinker Jr., A.G., Zimmermann, B.H., Scully, J.L., Kim, H., et al., 1990. Mammalian dihydroorotase: nucleotide sequence, peptide sequences, and evolution of the dihydroorotase domain of the multifunctional protein CAD. *Proc. Natl. Acad. Sci. U. S. A.* 87 (1), 174–178.
- Streb, H., Irvine, R.F., Berridge, M.J., Schulz, I., 1983. Release of Ca²⁺ from a nonmitochondrial intracellular store in pancreatic acinar cells by inositol-1,4,5-trisphosphate. *Nature* 306 (5938), 67–69.
- Thomas, A.P., Alexander, J., Williamson, J.R., 1984. Relationship between inositol polyphosphate production and the increase of cytosolic free Ca²⁺ induced by vasopressin in isolated hepatocytes. *J. Biol. Chem.* 259 (9), 5574–5584.
- Trager, W., Jensen, J.B., 1976. Human malaria parasites in continuous culture. *Science* 193 (4254), 673–675.
- Usui-Aoki, K., Matsumoto, K., Koganezawa, M., Kohatsu, S., Isono, K., Matsubayashi, H., et al., 2005. Targeted expression of Ip3 sponge and Ip3 dsRNA impairs sugar taste sensation in *Drosophila*. *J. Neurogenet.* 19 (3–4), 123–141.
- Varotti, F.P., Beraldo, F.H., Gazarini, M.L., Garcia, C.R., 2003. *Plasmodium falciparum* malaria parasites display a THG-sensitive Ca²⁺ pool. *Cell Calcium* 33 (2), 137–144.
- Walsh, P., Bursac, D., Law, Y.C., Cyr, D., Lithgow, T., 2004. The J-protein family: modulating protein assembly, disassembly and translocation. *EMBO Rep.* 5 (6), 567–571.
- Winding, P., Berchtold, M.W., 2001. The chicken B cell line DT40: a novel tool for gene disruption experiments. *J. Immunol. Methods* 29 (1–2), 1–16.
- Zhang, M., Wang, C., Otto, T.D., Oberstaller, J., Liao, X., Adapa, S.R., et al., 2018. Uncovering the essential genes of the human malaria parasite *Plasmodium falciparum* by saturation mutagenesis. *Science* 360 (6388).