

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Pullan, RL; Bethony, JM; M Geiger, S; Correa-Oliveira, R; Brooker, S; Quinnell, RJ; (2009) Human helminth co-infection: no evidence of common genetic control of hookworm and *Schistosoma mansoni* infection intensity in a Brazilian community. *International journal for parasitology*, 40 (3). pp. 299-306. ISSN 0020-7519 DOI: <https://doi.org/10.1016/j.ijpara.2009.08.002>

Downloaded from: <http://researchonline.lshtm.ac.uk/4951/>

DOI: <https://doi.org/10.1016/j.ijpara.2009.08.002>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

Published in final edited form as:

Int J Parasitol. 2010 March 1; 40(3): 299–306. doi:10.1016/j.ijpara.2009.08.002.

Human helminth co-infection: no evidence of common genetic control of hookworm and *Schistosoma mansoni* infection intensity in a Brazilian community

Rachel L. Pullan^{a,*}, Jeffrey M. Bethony^{b,c}, Stefan M. Geiger^{b,c}, Rodrigo Correa-Oliveira^b, Simon Brooker^{a,d}, and Rupert J. Quinnell^e

^aDepartment of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, United Kingdom

^bRené Rachou Research Centre FIOCRUZ, Belo Horizonte, Brazil

^cThe George Washington University, Washington D.C., United States of America

^dMalaria Public Health and Epidemiology Group, KEMRI-Wellcome Trust Research Programme, Nairobi, Kenya

^eInstitute of Integrative and Comparative Biology, University of Leeds, Leeds, United Kingdom

Abstract

Strong statistical associations between soil transmitted helminths and schistosomes are frequently observed in co-endemic human populations, although the underlying explanations remain poorly understood. This study investigates the contribution of host genetics and domestic environment to hookworm and *Schistosoma mansoni* infection intensity and evaluates the role of genetic and non-genetic factors in covariation of infection intensity. Detailed genealogical information allowed assignment of 1,303 individuals living in the Brazilian community of Americaninhas, Minas Gerais state, to 25 pedigrees (containing between two and 1,159 members) residing in 303 households. The prevalence of co-infection with both hookworms and schistosomes was high (38.5%), with significant correlation between *Necator americanus* and *S. mansoni* faecal egg counts ($r = 0.242$; $P < 0.0001$). Bivariate variance component analysis demonstrated a modest but significant species-specific heritability for intensity of *N. americanus* ($h^2 = 0.196$) and *S. mansoni* infection ($h^2 = 0.230$). However, after accounting for demographic, socio-economic and household risk factors, no evidence for common genetic control of intensity of hookworm and schistosome infection was observed (genetic correlation $\rho_G = 0.15$; $P = 0.52$). There was some evidence for residual clustering within households (household correlation $\rho_C = 0.45$; $P = 0.06$), but the majority (63%) of the covariance between *N. americanus* and *S. mansoni* infection intensity remained specific to the individual and could not be explained by shared genes, shared environment or other shared demographic, socio-economic or environmental risk factors. Our results emphasize the importance of exposure to hookworm and schistosome infection in driving the association between levels of infection with these species in hosts resident in areas of high transmission and suggest that much of this common exposure occurs outside the home.

*Corresponding author. R.L. Pullan, Department of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, United Kingdom Tel: +44 (0)20 7299 4667 ; Fax: +44 (0)20 7436 5389 rachel.pullan@lshtm.ac.uk.

Keywords

Variance component analysis; Heritability; *Schistosoma mansoni*; *Necator americanus*; Co-infection

1. Introduction

Numerous epidemiological studies indicate that individuals infected with multiple helminths, particularly soil-transmitted helminths (STH; especially hookworms and *Ascaris lumbricoides*) and schistosomes, harbour heavier worm burdens for each species than those infected with a single species (Booth et al., 1998; Needham et al., 1998; Brooker et al., 2000; Tchuem Tchuente et al., 2003; Fleming et al., 2006). Presence (and intensity) of co-infection may result from environmental and/or behavioural factors of the human host; for example, factors specific to the peri-domiciliary environment (Holland et al., 1988; Bethony et al., 2001; Raso et al., 2005; Hotez, 2007) or to family-specific behaviours (Bethony et al., 2001; Clennon et al., 2004). Higher levels of infection in co-infected individuals may alternatively be explained by host immunological or physiological factors (Fulford et al., 1998), phenotypes which in turn may be influenced by host genetics (Bethony et al., 1999), nutrition (Bundy and Golden, 1987) or the parasites themselves (Pearce and MacDonald, 2002; Loukas et al., 2005). Although evidence from quantitative genetic studies and genome scans of humans living in helminth endemic areas suggests a genetic basis for intensity of helminth infection (reviewed by Quinnell, 2003; Bethony and Quinnell, 2008), few studies have investigated whether phenotypic correlation in intensity of helminth infections can be attributed to common host genetic influences in humans. A recent quantitative genetic analysis of individuals infected with multiple helminths in rural Jiangxi Province, China, suggested that, whilst there was some evidence that multiple helminth infections aggregated within families, the risk of multiple infection for an individual was largely environmentally influenced (Ellis et al., 2007).

In a previous study, we explored the role of exposure-related factors in defining the epidemiological and spatial patterns of co-infection with the hookworm *Necator americanus* and *Schistosoma mansoni* in a rural Brazilian community (Pullan et al., 2008). While this study revealed that a limited number of household and environmental factors explained much of the spatial variability in the presence of co-infection, the majority (66%) of between-household variation in the occurrence of co-infection could not be explained by these factors, pointing to the need to evaluate the involvement of additional behavioural and genetic elements. On this basis we applied a multiple outcome approach to separate the effects of host genetic factors, common domestic environmental effects and residual individual variation to evaluate the evidence for shared genetic and non-genetic control of covariation in *N. americanus* and *S. mansoni* infection intensity. The extension of univariate quantitative genetic analysis to investigate multivariate outcomes has been described in detail (Hopper and Matthews, 1982; Lange and Boehnke, 1983; Carey, 1988), but this is the first time, to our knowledge, that it has been applied in the context of parasitic diseases in humans.

2. Materials and methods

2.1. Study area, population and recruitment

The study was conducted in Americaninhas, a region in the municipality of Nova Oriente, in northeastern Minas Gerais state, Brazil in 2004. Details of the study area, recruitment and enrolment procedures, as well as cross-sectional parasitological and survey data have been provided elsewhere (Brooker et al., 2006; Fleming et al., 2006; Brooker et al., 2007b), with

only a summary provided here. Briefly, the study was designed as a total population survey, with the research team visiting all households in a 10 km² area to obtain informed consent using a written and verbal consent form approved by the ethical committee of the Centro de Pesquisas René Rachou-FIOCRUZ, the Brazilian National Committee for Ethics in Research (CONEP), George Washington University Medical Center (USA), and the London School of Hygiene and Tropical Medicine (UK). Each house was assigned a unique household identification number (HHID), and each resident a unique personal identifier (PID).

2.2. Mapping, household questionnaire and parasitological survey

All households in the study area were geo-referenced and information on household socio-economic and physical characteristics was collected using a pre-tested, standardized household questionnaire. Remotely sensed environmental data were extracted for May 2001 from the Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER) satellite sensor at 30 m spatial resolution. ASTER provides information on Normalized Difference Vegetation Index (NDVI), a proxy of vegetation density and soil moisture, and digital elevation (Tatem et al., 2004). During the parasitological survey, stool samples were collected over the course of 2 days. Those confirmed as egg-positive for any helminth species using formalin-ether sedimentation were subsequently examined by Kato–Katz faecal thick smear to quantify the intensity of the infection as eggs per gram of faeces (epg). Two slides were taken from each day's faecal sample for a total of up to four slides from each individual. Individuals positive by formalin-ether sedimentation but negative by Kato-Katz were assigned an egg count of 3 epg, half the Kato-Katz detection limit. Hookworm was confirmed as exclusively *N. americanus* by both morphological examination and PCR (Pullan et al., 2008).

2.3. Pedigree collection and identification

To establish kinship relationships within and between households, questionnaires were administered to either the head male or female of the household to elicit genealogical and demographic information on each household member, including their age, sex and names of parents. Relationships between all household members were recorded, and records were also made of any other first- or second-degree relatives living in other households in the study area to establish genetic ties between households. Individuals were then defined as belonging to the same extended pedigree if they were related to anyone else in the pedigree or were married to anyone in the pedigree. Pedigrees were assembled using PEDSYS (Dyke, 1999), and the number of relative pairs calculated using the 'kinship' function. Doubtful pedigree relationships were confirmed by re-interviewing relevant heads of households. The 1,303 sampled individuals who provided pedigree and household information and at least one phenotype could be assembled into one large multi-generational pedigree comprising 1,159 people, and 24 smaller pedigrees of two to 11 phenotyped individuals each; 36 individuals had no phenotyped relatives in the study area. There were 303 households, with one to 12 (mean 4.3) phenotyped individuals per household.

2.4. Data management and preliminary statistical analysis

The outcomes used in this analysis were intensity of infection with *N. americanus* and *S. mansoni*. The methods used assume the phenotype to be normally distributed; therefore, because of the skewed distribution of infection intensities (faecal egg counts), individual infection intensities were assessed after log-transformation as $\ln(\text{epg}+1)$. Analysis was not undertaken for *A. lumbricoides* because egg counts were truncated at 500 eggs per slide (12,000 epg).

Preliminary investigation of covariates was performed using Stata 9.1 (STATA Corporation, Houston, TX, USA). All available demographic (age, sex), household (toilet facilities, household crowding, flooring material, relative socio-economic status) and environmental (sector, NDVI, altitude and population and household density) covariates were included in a full fixed effect regression model, and non-significant ($P > 0.1$) variables removed sequentially to derive a minimally adequate model. Standard errors were adjusted for dependence between individuals within households using a clustered sandwich estimator. Visual examination of scatter-plots was used to investigate non-linear relationships. Covariates were analysed separately for each infection. Subsequently, the retained covariates were built into bivariate mixed linear regression models outlined below (genetic variance component analysis (VCA)).

2.5. Genetic VCA

Quantitative genetic analyses were conducted by means of a maximum likelihood-based variance decomposition approach implemented in the computer package SOLAR 4.2.0 (Almasy and Blangero, 1998). In brief, this approach incorporates information on genetic relationships within and between households in order to partition the total variation in infection intensity into its genetic, household and other causes. The a priori expectation for the covariance between any two individuals is determined by the degree of relatedness between them (equal to twice the kinship coefficient (ϕ), which represents a genome-averaged probability that two individuals will share an allele identical by descent (Hopper and Matthews, 1982)). Incorporation of a “kinship coefficient matrix” into a random effect model therefore allows the covariance between relatives to depend upon the degree of relatedness between them, providing an estimate of the total cumulative effect of individual genes (the “additive genetic” effect) on phenotypic variation. The relative contribution of fixed covariates to total variation in infection intensity (total phenotypic variance) was estimated by comparing the trait S.D. in models with and without fixed covariates.

For each infection, four initial models were investigated. The ‘sporadic’ model attributes variation in faecal egg excretion entirely to a random, individual-specific effect (e^2). The ‘household’ model introduces an additional household-level random effect (c^2), allowing estimation of the effect of unmeasured factors associated with the common domestic environment, whilst the ‘polygenic’ model introduces a random effect describing additive genetic variation (h^2 heritability) by incorporating a “relationship matrix”. Finally, in the ‘saturated’ model, all of these effects are modelled and estimated together. The variance parameters were standardised by dividing by the total phenotypic variation (i.e. the proportion of total variance attributable to each parameter was estimated). Chi-squared testing based on likelihood ratios was then used to test the significance of variance parameters.

Bivariate variance components were fitted to estimate the role of additive genetics and shared household in the covariance between two traits (e.g. $\ln(\text{epg}+1)$ of hookworm infection and $\ln(\text{epg}+1)$ of schistosome infection). By this approach, the variance-covariance matrix is partitioned into additive genetic, shared-household and random, individual-specific effects, providing information on the pair-wise covariance between traits. The variance-covariance matrices for these bivariate outcome models provided estimates of three additional parameters: the additive genetic correlation (ρ_G), the shared-household correlation (ρ_C) and unexplained correlation (ρ_E) between infection intensity. Respectively, these correlations are estimates of the common additive effects of genes on the phenotypic variance of each trait, and the effects of domestic environmental factors and individual non-genetic factors common to both traits (Carey, 1988). The significance of each of the additional estimated parameters was evaluated by likelihood-ratio tests comparing the general model, in which the covariance parameter is estimated, with a restricted model in

which the covariance parameter is constrained to zero (i.e. no correlation between infections for that effect). Shared genetic control is indicated by an additive genetic correlation (ρ_G) that is significantly different from zero.

An unbiased estimate of the total phenotypic correlation (ρ_P) between the two species can be calculated based on these pair-wise correlations using the following equation:

$$\rho_P = \left(\sqrt{h_1^2} \sqrt{h_2^2} \rho_G \right) + \left(\sqrt{c_1^2} \sqrt{c_2^2} \rho_C \right) + \left(\sqrt{e_1^2} \sqrt{e_2^2} \rho_E \right)$$

where the sub-script 1 and 2 represent variance component estimates for each trait, respectively.

3. Results

3.1. Parasitological and demographic data

In total, 1,303 individuals provided pedigree and household information and at least one parasitological phenotype (87% of the total population): intensity of hookworm infection was available for 1,294 individuals, whilst intensity of *S. mansoni* infection was available for 1,302. Overall, the prevalence of *N. americanus* was 70.9% (mean epg = 1,441), and *S. mansoni* was 47.2% (mean epg = 131); 38.5% of individuals were co-infected with both species. A significant positive phenotypic correlation was observed between log transformed *N. americanus* and *S. mansoni* infection intensities ($r = 0.242$, $n = 1,293$, $P < 0.0001$), as has been described previously (Fleming et al., 2006). Intensity of hookworm infection rose during childhood, reaching a peak between 5 and 10 years of age where it remained constant throughout adulthood; mean intensity was greater in males than females. In contrast *S. mansoni* infection intensity peaked between 10 and 25 years of age and declined throughout adulthood.

Findings from unadjusted linear regression models reflect our previous multinomial analysis of *N. americanus* – *S. mansoni* co-infection: *N. americanus* infection intensity was positively associated with characteristics associated with lower socio-economic status (low socio-economic index, household crowding) and with higher NDVI, whilst both infections showed significant associations with lack of toilet facilities. Mean hookworm infection intensity was lower in the village area (sector 1) compared with the rural sectors (sectors 3-7); whilst for *S. mansoni* mean infection intensity was highest in the village area.

3.2. Variance components analysis of individual helminth infection intensity

To define the contribution of hereditary and environmental influences on infection intensity a quantitative genetic variance component approach was applied. Table 1 describes the number of relative pairs included in these analyses. In total, when considering up to seven degrees of relatedness, 40,476 relative pairs were available. Four models were initially developed: sporadic, household, polygenic and saturated. Maximum-likelihood estimates of the variance components, and the corresponding model log-likelihoods, are provided in Table 2. Results for *N. americanus* and *S. mansoni* are similar. Examination of log-likelihood values reveals that infection intensity was best described by a saturated model; in the absence of covariates, heritability (h^2) was 0.30 for *N. americanus* and 0.29 for *S. mansoni*, with a further 0.15 and 0.17 of variation in egg counts attributable to the domestic environment.

At an individual level, *N. americanus* intensity was modelled by a sex-dependent linear increase with age from 0-5 years, with no change in older hosts; intensity was higher in adult

males than females. Other covariates included in the *N. americanus* model were the absence of toilet facilities, household overcrowding (> 1 people/room), relative socio-economic index (linear), NDVI and sector; the *S. mansoni* model included age (modelled as age and age²), absence of toilet facilities, and sector. Introduction of these demographic, socio-economic and environmental covariates led to a slight reduction in the proportion of remaining variation explained by the additive genetic (to 0.23-0.24) and shared household variance components (to 0.12 for *N. americanus* and 0.11 for *S. mansoni*). It did not however lead to different conclusions for the hypothesis tests; for both infections both additive genetic effects and household effects were highly significant ($P < 0.001$ for χ^2 tests comparing the saturated model with simpler (household or polygenic) models).

3.3. Bivariate variance components analysis of *N. americanus* and *S. mansoni* intensity

To estimate the contribution of additive genetic effects and domestic environment to the phenotypic correlation between hookworm and *S. mansoni* intensities we used a bivariate variance component analysis. Estimates of the covariance components are presented in Table 3. Total unbiased correlation between infection intensities (ρ_p) was 0.27, which reduced to 0.24 after adjusting for covariates. The relatively small influence of covariates (which explained only 11% of the correlation between these two traits) is unsurprising given that the only common explanatory factors identified were age, toilet facilities and sector. Correlation between individuals' hookworm and *S. mansoni* worm burdens was unrelated to the degree of relationship between them, as indicated by the lack of evidence for covariance between the additive genetic effects for each trait ($\rho_G = 0.147$; $P = 0.52$). Evidence for positive correlation due to shared domestic environment was weak ($\rho_C = 0.451$; $P = 0.06$). However, there was strong evidence for residual (non-genetic, non-household) correlation between intensity of infection at the individual level ($\rho_E = 0.233$; $P << 0.001$); after adjusting for covariates an estimated 5% of residual phenotypic variance could be attributed to additional unmeasured factors common to both species ($\rho_E^2 = 0.054$).

The estimates of all fixed effects and variance parameters are shown in Table 4, and resulting estimates of shared and species-specific sources of variation are shown in Fig. 1. Heritability was comparable to that estimated from univariate models; for *N. americanus* $h^2 = 0.196$, for *S. mansoni* $h^2 = 0.230$. A strong age dependency was observed for intensity of infection for both helminths; for the intensity of hookworm infection, this effect was greater in males than females. Together, age and sex explained 11.6% of total variation of hookworm burden and 13.8% of *S. mansoni* burden. Socio-economic and geographical environmental covariates associated with the household explained substantially more of the total variation for hookworm infection intensity than for *S. mansoni* infection intensity (a further 13.1% and 4.2%, respectively), with a further 9.9-11.4% attributable to other factors associated with the domestic environment. When expressed as a proportion of total, rather than residual, variation the contribution of host genetics to the variation in faecal egg excretion was 14.8% for hookworm and 18.9% for schistosomiasis. A remaining 46.5-50.8% of the heterogeneity in faecal egg counts was not explained by any of the factors included in this model.

4. Discussion

In this analysis, we build upon recent literature exploring infection intensity for human helminth infection. By modelling the intensity of *N. americanus* and *S. mansoni* infection as a bivariate outcome, we were able to assess host genetic and non-genetic influences unique to each infection and common to both. As frequently observed in co-endemic communities, there was a strong positive phenotypic correlation between the intensity of infection for these two species ($\rho_p = 0.24$ when adjusting for covariates). However, when modelled in the presence of shared domestic environment, no evidence was found for common genetic

control of infection intensity, although there was weak evidence of a household contribution to the correlation between egg counts for these species within individual hosts ($\rho_C = 0.45$; $P = 0.06$). Importantly, the majority (63%) of the covariance between *N. americanus* and *S. mansoni* infection intensity remained specific to the individual and could not be explained by host genetics, the domestic environment or known demographic, socio-economic and environmental risk factors.

Our conclusions are based on the results of a bivariate genetic variance components analysis. This method has been used extensively to model relationships between biological phenotypes (Edwards et al., 1999; Hokanson et al., 2003; Njajou et al., 2006; Lichtenstein et al., 2009), although we believe this is the first time it has been applied in the context of human parasitic helminth infection. We demonstrate a modest but significant heritability for intensity of *N. americanus* ($h^2 = 0.196$) and *S. mansoni* infection ($h^2 = 0.230$) in hosts from an endemic area. Heritability estimates are population-specific, depending on the magnitude of both genetic and environmental variability; here, heritability is derived from a single large pedigree, which may result in lower genetic and/or environmental variance. However, our estimates are consistent with previous studies establishing a role for host genetics in determining infection burden for the major human STH (Williams-Blangero et al., 1997, 1999, 2002a; Breitling et al., 2008) and schistosome infections (Bethony et al., 2002; King et al., 2004; Seto et al., 2005; Ellis et al., 2006). Importantly, our results suggest that there is no, or very little, shared genetic control of *N. americanus* and *S. mansoni* infection intensity in this population, with a low and non-significant genetic correlation of 0.147.

Strong positive genetic correlations have been previously demonstrated between *Trichuris suis* and *Ascaris suum* burdens in experimentally infected pigs, and between *Haemonchus contortus* and *Trichostrongylus colubriformis* in sheep, suggesting that resistance to intestinal nematodes of different species may involve regulation by overlapping sets of genes (Gruner et al., 2004; Nejsun et al., 2009). The lack of a positive genetic correlation in our study may reflect greater biological differences between hookworms and schistosomes (including substantially different transmission cycles, locations within the host and interactions with the host immune system). However, it should be noted that assessing evidence for biological pleiotropy from statistical correlations can be difficult (Carey, 1988). For example, dominance and epistatic effects that differ across traits, linkage disequilibrium, gene-environment interactions and simple pedigree errors can all serve to decrease (or increase) the estimate of additive genetic correlation (Carey, 1988). Identification of the genes involved in determining intensity of infection for each species will be needed to confirm whether there are any common genetic control. Whilst some loci controlling *S. mansoni* (Marquet et al., 1996) *A. lumbricoides* (Williams-Blangero et al., 2002b) and *Trichuris trichiura* (Williams-Blangero et al., 2008) infection have been mapped, to our knowledge there have been no such studies carried out for either of the hookworm species, an area warranting further investigation.

After accounting for socio-economic, environmental and household covariates, further unmeasured factors associated with the domestic environment accounted for up to 25% of the correlation in intensity of infection between species in this community. This finding is in keeping with a previous observation that occurrences of co-infections with *Schistosoma japonicum* – *T. trichiura* or with *A. lumbricoides* – *T. trichiura* were aggregated within households, but not within families, in a rural Chinese population (Ellis et al., 2007), and suggests that, despite differences in the mode of transmission between schistosomes and STHs, common exposures shared by members of the same household may play an important role in transmission of both species. These observations emphasise that improvements in sanitation and health awareness within the home, as well as targeted treatment of those

living in poor conditions, will impact significantly on the transmission of multiple helminth species.

Importantly however, most of the phenotypic correlation between these two species could be explained by neither the domestic environment nor host genetics, suggesting that common exposures outside the home may be driving the relationship. Although (as mentioned above) the ecology and transmission of *S. mansoni* and hookworm differ considerably, occupation, hygiene behaviours, health knowledge and water-contact patterns may all potentially influence exposure to both infections (Watts et al., 1998; Bethony et al., 2004; Matthys et al., 2007). For example, a recent analysis of risk factors for *S. mansoni* and hookworm in urban farming communities in western Côte d'Ivoire revealed that both infections were related to certain agricultural practices, including contact with irrigation wells that may provide hiding places for open defecation (Matthys et al., 2007). Similarly, water contact was a significant risk factor for both *S. japonicum*, *T. trichiura* and *A. lumbricoïdes* in the Poyang Lake region in China (Ellis et al., 2007). Interestingly, there was very little role for individual-level factors in determining predisposition to hookworm infection in this Brazilian community, suggesting that environmental determinants of co-infection may be variable through time (Quinnell et al., in press). Alternatively, the unexplained correlation between *N. americanus* and *S. mansoni* infection intensity may reflect a direct biological or immunological interaction between species. *Schistosoma mansoni* infection influences immune responses to other helminth infections in mice (Curry et al., 1995; Yoshida et al., 1999), but whether helminth-induced immuno-suppression can influence the intensity of concomitant infection in humans requires further study (for review see Geiger, 2008).

As for single outcome genetic VCA, shared household is only a proxy measure for the influence of domestic environmental factors on phenotype. Given that environmental conditions may be common to multiple households within a community, it is important that potential sources of heterogeneity in environmental exposure to infection be measured and included as covariates. To our knowledge this is the first study of the genetic epidemiology of human helminth infection to include detailed information on socio-economic indicators, physical environment and spatial location (including remotely sensed environmental data) allowing greater quantification of sources of heterogeneities in exposure in this community. Despite these included covariates, nearly half of the total variation in infection intensity remained unexplained. Much of this may be due to measurement error, which is known to be large for faecal egg counts, even when (as in the current study) several slides were counted on separate days (Yu et al., 1998). Moreover, faecal egg counts represent an indirect measure of underlying worm burden; it is likely that density-dependent fecundity and possibly other host factors (including host genetics) may influence this association (Anderson and Schad, 1985; Agnew et al., 1996). Given the location in which the adult worms lay their eggs (the gut lumen versus mesenteric blood vessels) comparability between these two proxies of infection intensity may also be limited. As a consequence, studies such as ours that rely on faecal egg count for the quantitative assessment of the intensity of infection risk underestimate the genetic and non-genetic components (and correlations) for these infections. Unexplained variation in helminth infection intensity may also result from genetic heterogeneity in the parasite population. Molecular studies have revealed allelic and nucleotide diversity in the genomes of human helminth parasite populations (Hawdon et al., 2001; Curtis et al., 2002; Stohler et al., 2005), and further study of the relative importance of parasite genetic diversity in driving patterns of heterogeneity within human populations is warranted.

We believe this study represents the first analysis of the intensity of co-infection with *S. mansoni* and *N. americanus* to quantify the relative importance of host genetic and household factors in the commonly observed correlation in infection levels for these two

helminth species. Our results emphasise the importance of environmental factors in driving this association and suggest that whilst around a quarter of correlation can be explained by factors associated with the domestic environment, much of this common exposure occurs outside the home. Conflicting opinions on the nature of interactions between concomitant infections, particularly in relation to interactions between helminths and malaria, have highlighted the need for robust approaches to studying co-infection (Nacher, 2004; Booth, 2006; Mwangi et al., 2006; Brooker et al 2007a). Numerous studies support the view that, for the majority of infectious diseases, host susceptibility is likely to be polygenic, with the immunomodulatory effects of certain genetic polymorphisms influencing several diseases (Frodsham and Hill, 2004). However, it still remains unclear whether the associations frequently observed between parasitic infections are dictated by an individual's genotype, by direct synergistic or antagonistic interactions between species, or whether they simply reflect common patterns of exposure. By providing insight into the patterns of genetic and non-genetic relationships between traits, bivariate genetic VCA provides us with a valuable tool to help unravel the complex interplay between these infectious diseases.

Acknowledgments

We are very grateful to inhabitants of Americaninhas who kindly participated in the study, to the many members of the Centro de Pesquisas René Rachou staff for their technical assistance, and to Lutz Breitling (University of Leeds) for assembling the pedigrees. We are also most appreciative of those who were responsible for carrying out the fieldwork, which made this analysis possible. Fieldwork was financially supported by the Human Hookworm Vaccine Initiative (HHVI) of the Sabin Institute, which receives support from the Bill and Melinda Gates Foundation. RP is supported by a Medical Research Council DTA-funded studentship and SB is supported by a Career Development Fellowship (081673) from the Wellcome Trust. JB is supported by grants from the Sabin Institute, the Bill and Melinda Gates Foundation and NIAID/NIH.

References

- Agnew A, Fulford A, Mwanje MT, Gachuhi K, Gutschmann V, Kriger FW, Sturrock RF, Vennervald BJ, Ouma JH, Butterworth AE, Deelder AM. Age-dependent reduction of schistosome fecundity in *Schistosoma haematobium* but not *Schistosoma mansoni* infections in humans. *Am. J. Trop. Med. Hyg.* 1996; 55:338–343. [PubMed: 8842126]
- Almasy L, Blangero J. Multipoint quantitative-trait linkage analysis in general pedigrees. *Am. J. Hum. Genet.* 1998; 62:1198–1211. [PubMed: 9545414]
- Anderson RM, Schad GA. Hookworm burdens and faecal egg counts: an analysis of the biological basis of variation. *Trans. R. Soc. Trop. Med. Hyg.* 1985; 79:812–825. [PubMed: 3832493]
- Bethony J, Silveira AM, Alves-Oliveira LF, Thakur A, Gazzinelli G, Correa-Oliveira R, LoVerde PT. Familial resemblance in humoral immune response to defined and crude *Schistosoma mansoni* antigens in an endemic area in Brazil. *J. Infect. Dis.* 1999; 180:1665–1673. [PubMed: 10515830]
- Bethony J, Williams JT, Kloos H, Blangero J, Alves-Fraga L, Buck G, Michalek A, Williams-Blangero S, Loverde PT, Correa-Oliveira R, Gazzinelli A. Exposure to *Schistosoma mansoni* infection in a rural area in Brazil. II: Household risk factors. *Trop. Med. Int. Health.* 2001; 6:136–145. [PubMed: 11251910]
- Bethony J, Williams JT, Blangero J, Kloos H, Gazzinelli A, Soares-Filho B, Coelho L, Alves-Fraga L, Williams-Blangero S, Loverde PT, Correa-Oliveira R. Additive host genetic factors influence fecal egg excretion rates during *Schistosoma mansoni* infection in a rural area in Brazil. *Am. J. Trop. Med. Hyg.* 2002; 67:336–343. [PubMed: 12452486]
- Bethony J, Williams JT, Brooker S, Gazzinelli A, Gazzinelli MF, LoVerde PT, Correa-Oliveira R, Kloos H. Exposure to *Schistosoma mansoni* infection in a rural area in Brazil. Part III: household aggregation of water-contact behaviour. *Trop. Med. Int. Health.* 2004; 9:381–389. [PubMed: 14996368]
- Bethony JM, Quinnell RJ. Genetic epidemiology of human schistosomiasis in Brazil. *Acta Trop.* 2008; 108:166–174. [PubMed: 18207118]

- Booth M, Bundy DA, Albonico M, Chwaya HM, Alawi KS, Savioli L. Associations among multiple geohelminth species infections in schoolchildren from Pemba Island. *Parasitology*. 1998; 116:85–93. [PubMed: 9481778]
- Booth M. The role of residential location in apparent helminth and malaria associations. *Trends Parasitol*. 2006; 22:359–362. [PubMed: 16797235]
- Breitling LP, Wilson AJ, Raiko A, Lagog M, Siba M, Shaw MA, Quinnell RJ. Heritability of human hookworm infection in Papua New Guinea. *Parasitology*. 2008; 135:1407–1415. [PubMed: 18937884]
- Brooker S, Miguel EA, Moulin S, Luoba AI, Bundy DA, Kremer M. Epidemiology of single and multiple species of helminth infections among school children in Busia District, Kenya. *East African Medicine Journal*. 2000; 77:157–161.
- Brooker S, Alexander N, Geiger S, Moyeed RA, Stander J, Fleming F, Hotez PJ, Correa-Oliveira R, Bethony J. Contrasting patterns in the small-scale heterogeneity of human helminth infections in urban and rural environments in Brazil. *Int. J. Parasitol*. 2006; 36:1143–1151. [PubMed: 16814294]
- Brooker S, Akhwale WS, Pullan R, Estambale B, Clarke S, Snow RW, Hotez PJ. Epidemiology of *Plasmodium*-Helminth coinfection in Africa: potential impact on anaemia and prospects for combining control. *Am. J. Trop. Med. Hyg*. 2007a; 77:88–98. [PubMed: 18165479]
- Brooker S, Jardim-Botelho A, Quinnell RJ, Geiger SM, Caldas IR, Fleming F, Hotez PJ, Correa-Oliveira R, Rodrigues LC, Bethony JM. Age-related changes in hookworm infection, anaemia and iron deficiency in an area of high *Necator americanus* hookworm transmission in southeastern Brazil. *Trans. R. Soc. Trop. Med. Hyg*. 2007b; 101:146–154. [PubMed: 17027054]
- Bundy DAP, Golden MH. The impact of host nutrition on gastrointestinal helminth populations. *Parasitology*. 1987; 95:623–635. [PubMed: 3320892]
- Carey G. Inference about genetic correlations. *Behav. Genet*. 1988; 18:329–338. [PubMed: 3219111]
- Clennon JA, King CH, Muchiri EM, Karuiki HC, Ouma JH, Mungai P, Kitron U. Spatial patterns of urinary schistosomiasis infection in a highly endemic area of coastal Kenya. *Am. J. Trop. Med. Hyg*. 2004; 70:443–448. [PubMed: 15100462]
- Curry AJ, Else KJ, Jones F, Bancroft A, Grecis RK, Dunne DW. Evidence that cytokine-mediated immune interactions induced by *Schistosoma mansoni* alter disease outcome in mice concurrently infected with *Trichuris muris*. *J. Exp. Med*. 1995; 181:769–774. [PubMed: 7836929]
- Curtis J, Sorensen RE, Minchella DJ. Schistosome genetic diversity : the implications of population structure as detected with microsatellite markers. *Parasitology*. 2002; 125:S51–S59. [PubMed: 12622328]
- Dyke, B. PEDSYS: A Pedigree Data Management System. Southwest Foundation for Biomedical Research; San Antonio, TX: 1999.
- Edwards KL, Mahaney MC, Motulsky AG, Austin MA. Pleiotropic genetic effects on LDL size, plasma triglyceride and HDL cholesterol in families. *Arterioscler. Thromb. Vasc. Biol*. 1999; 19:2456–2464. [PubMed: 10521376]
- Ellis MK, Li Y, Rong Z, Chen H, McManus DP. Familial aggregation of human infection with *Schistosoma japonicum* in the Poyang Lake region, China. *Int. J. Parasitol*. 2006; 36:71–77. [PubMed: 16321389]
- Ellis MK, Raso G, Li Y-S, Rong Z, Chen H-G, McManus DP. Familial aggregation of human susceptibility to co- and multiple helminth infections in a population from the Poyang Lake region, China. *Int J Parasitol*. 2007; 37:1153–61. [PubMed: 17407778]
- Fleming FM, Brooker S, Geiger SM, Caldas IR, Correa-Oliveira R, Hotez PJ, Bethony JM. Synergistic associations between hookworm and other helminth species in a rural community in Brazil. *Trop. Med. Int. Health*. 2006; 11:56–64. [PubMed: 16398756]
- Frodsham AJ, Hill AVS. Genetics of infectious diseases. *Hum. Mol. Genet*. 2004; 13:R187–R194. [PubMed: 15358724]
- Fulford AJ, Webster M, Ouma JH, Kimani G, Dunne DW. Puberty and age-related changes in susceptibility to schistosome infection. *Parasitol. Today*. 1998; 14:23–26. [PubMed: 17040685]

- Geiger S. Immuno-epidemiology of *Schistosoma mansoni* infections in endemic populations co-infected with soil-transmitted helminths: present knowledge, challenges, and the need for further studies. *Acta Trop.* 2008; 108:118–123. [PubMed: 18582842]
- Gruner L, Bouix J, Brunel JC. High genetic correlation between resistance to *Haemonchus contortus* and to *Trichostrongylus colubriformis* in INRA 401 sheep. *Vet. Parasitol.* 2004; 119:51–58. [PubMed: 15036576]
- Hawdon JM, Li T, Zhan B, Blouin MS. Genetic structure of populations of the human hookworm, *Necator americanus*, in China. *Mol. Ecol.* 2001; 10:1433–1437. [PubMed: 11412366]
- Hokanson JE, Langefeld CD, Mitchell BD, Lange LA, Goff DCJ, Haffner SM, Saad MF, Rotter JI. Pleiotropy and heterogeneity in the expression of atherogenic lipoproteins: the IRAS Family Study. *Hum. Hered.* 2003; 55:46–50. [PubMed: 12890925]
- Holland CV, Taren DL, Crompton DW, Nesheim MC, Sanjur D, Barbeau I, Tucker K, Tiffany J, Rivera G. Intestinal helminthiasis in relation to the socioeconomic environment of Panamanian children. *Soc. Sci. Med.* 1988; 26:209–213. [PubMed: 3347848]
- Hopper JL, Matthews JD. Extensions to multivariate normal models for pedigree analysis. *Ann. Hum. Genet.* 1982; 46:373–383. [PubMed: 6961886]
- Hotez PJ. Hookworm and poverty. *Ann. New York Acad. Sci.* 2007; 1136:38–44. [PubMed: 17954674]
- King CH, Blanton RE, Muchiri EM, Ouma JH, Kariuki HC, Mungai P, Magak P, Kadzo H, Ireri E, Koech DK. Low heritable component of risk for infection intensity and infection-associated disease in urinary schistosomiasis among Wadigo village populations in Coast Province, Kenya. *Am. J. Trop. Med. Hyg.* 2004; 70:57–62. [PubMed: 14971699]
- Lange K, Boehnke M. Extensions to pedigree analysis, IV: covariance components models for multivariate traits. *Am. J. Med. Genet.* 1983; 14:513–524. [PubMed: 6859102]
- Lichtenstein P, Yip BH, Bjork C, Pawitan Y, Cannon TD, Sullivan PF, Hultman CM. Common genetic determinants of schizophrenia and bipolar disorder in Swedish families: a population-based study. *Lancet.* 2009; 373:234–239. [PubMed: 19150704]
- Loukas A, Constant SL, Bethony JM. Immunobiology of hookworm infection. *FEMS Immunol. Med. Microbiol.* 2005; 43:115–124. [PubMed: 15681140]
- Marquet S, Abel L, Hillaire D, Dessein H, Kalil J, Feingold J, Weissenbach J, Dessein AJ. Genetic localization of a locus controlling the intensity of infection by *Schistosoma mansoni* on chromosome 5q31-q33. *Nat. Genet.* 1996; 14:181–184. [PubMed: 8841190]
- Matthys B, Tschannen AB, Tian-Bi NT, Comoé H, Diabaté S, Traoré M, Vounatsou P, Raso G, Gosoniu L, Tanner M, Cissé G, N'Goran EK, Utzinger J. Risk factors for *Schistosoma mansoni* and hookworm in urban farming communities in western Côte d'Ivoire. *Trop. Med. Int. Health.* 2007; 12:709–723. [PubMed: 17550468]
- Mwangi TW, Bethony JM, Brooker S. Malaria and helminth interactions in humans: an epidemiological viewpoint. *Ann. Trop. Med. Parasitol.* 2006; 100:551–570. [PubMed: 16989681]
- Nacher M. Interactions between worm infections and malaria. *Clin. Rev. Allergy Immunol.* 2004; 26:85–92. [PubMed: 15146105]
- Needham C, Kim HT, Hoa NV, Cong LD, Michael E, Drake L, Hall A, Bundy DA. Epidemiology of soil-transmitted nematode infections in Ha Nam Province, Vietnam. *Trop. Med. Int. Health.* 1998; 3:904–912. [PubMed: 9855404]
- Nejsum P, Roepstorff A, Jorgensen CB, Fredholm M, Goring HHH, Anderson TJC, Thamsborg SM. High heritability for *Ascaris* and *Trichuris* infection levels in pigs. *Heredity.* 2009; 102:357–364. [PubMed: 19142203]
- Njajou OT, Alizadeh BZ, Aulchenko Y, Zillikens MC, Pols HA, Oostra BA, Swinkels DW, van Duijn CM. Heritability of serum iron, ferritin and transferrin saturation in a genetically isolated population, the Erasmus Rucphen Family (ERF) Study. *Hum. Hered.* 2006; 61:222–228. [PubMed: 16877869]
- Pearce EJ, MacDonald AS. The immunobiology of schistosomiasis. *Nat. Rev. Immunol.* 2002; 2:499–511. [PubMed: 12094224]

- Pullan R, Bethony J, Geiger S, Cundill B, Correa-Oliveira R, Quinnell RJ, Brooker S. Human helminth co-infection: an analysis of spatial heterogeneity and household and environmental risk factors in a Brazilian community. *PLoS Negl. Trop. Dis.* 2008; 2:e352. [PubMed: 19104658]
- Quinnell RJ. Genetics of susceptibility to human helminth infection. *Int. J. Parasitol.* 2003; 33:1219–1231. [PubMed: 13678637]
- Quinnell RJ, Pullan RL, Breitling L.Ph. Geiger S, Cundill B, Correa-Oliveira R, Brooker S, Bethony JM. Genetic and household determinants of predisposition to human hookworm infection in a Brazilian community. *J. Infect. Dis.* in press.
- Raso G, Utzinger J, Silue KD, Ouattara A, Yapi A, Toty A, Matthys B, Vounatsou P, Tanner M, N’Goran E,K. Disparities in parasitic infections, perceived ill health and access to health care among poorer and less poor schoolchildren of rural Côte d’Ivoire. *Trop. Med. Int. Health.* 2005; 10:42–57. [PubMed: 15655013]
- Seto EY, Zhong B, Kouch J, Hubbard A, Spear RC. Genetic and household risk factors for *Schistosoma japonicum* infection in the presence of larger scale environmental differences in the mountainous transmission areas of China. *Am. J. Trop. Med. Hyg.* 2005; 73:1145–1150. [PubMed: 16354828]
- Stohler RA, Curtis J, Minchella DJ. A comparison of microsatellite polymorphism and heterozygosity among field and laboratory populations of *Schistosoma mansoni*. *Int. J. Parasitol.* 2004; 34:595–601. [PubMed: 15064124]
- Tatem AJ, Goetz SJ, Hay SI. Terra and Aqua: new data for epidemiology and public health. *Int. J. App. Earth Obs. and Geo.* 2004; 6:33–46.
- Tchuem Tchuente LA, Behnke JM, Gilbert FS, Southgate VR, Vercruyse J. Polyparasitism with *Schistosoma haematobium* and soil-transmitted helminth infections among school children in Loum, Cameroon. *Trop. Med. Int. Health.* 2003; 8:975–986. [PubMed: 14629763]
- Watts S, Khallaayoune K, Bensefia R, Laamrani H, Gryseels B. The study of human behavior and schistosomiasis transmission in an irrigated area in Morocco. *Soc. Sci. Med.* 1998; 46:755–765. [PubMed: 9522434]
- Williams-Blangero S, Blangero J, Bradley M. Quantitative genetic analysis of susceptibility to hookworm infection in a population from rural Zimbabwe. *Hum. Biol.* 1997; 69:201–208. [PubMed: 9057344]
- Williams-Blangero S, Subedi J, Upadhayay RP, Manral DB, Rai DR, Jha B, Robinson ES, Blangero J. Genetic analysis of susceptibility to infection with *Ascaris lumbricoides*. *Am. J. Trop. Med. Hyg.* 1999; 60:921–926. [PubMed: 10403321]
- Williams-Blangero S, McGarvey ST, Subedi J, Wiest PM, Upadhayay RP, Rai DR, Jha B, Olds GR, Guanling W, Blangero J. Genetic component to susceptibility to *Trichuris trichiura*: evidence from two Asian populations. *Genet. Epidemiol.* 2002a; 22:254–264. [PubMed: 11921085]
- Williams-Blangero S, VandeBerg JL, Subedi J, Aivaliotis MJ, Rai DR, Upadhayay RP, Jha B, Blangero J. Genes on chromosomes 1 and 13 have significant effects on *Ascaris* infection. *Proc. Natl. Acad. Sci. USA.* 2002b; 99:5533–5538. [PubMed: 11960011]
- Williams-Blangero S, Vandeberg JL, Subedi J, Jha B, Dyer TD, Blangero J. Two quantitative trait loci influence whipworm (*Trichuris trichiura*) infection in a Nepalese population. *J. Infect. Dis.* 2008; 197:1198–1203. [PubMed: 18462166]
- Yoshida A, Maruyama H, Yabu Y, Amano T, Kobayakawa T, Ohta N. Immune response against protozoal and nematodal infection in mice with underlying *Schistosoma mansoni* infection. *Parasitol. Int.* 1999; 48:73–79. [PubMed: 11269328]
- Yu JM, De Vlas SJ, Yuan HC, Gryseels B. Variations in fecal *Schistosoma japonicum* egg counts. *Am. J. Trop. Med. Hyg.* 1998; 59:370–375. [PubMed: 9749627]




Fig. 1. Results of bivariate quantitative genetic analysis of infection intensity for *Necator americanus* and *Schistosoma mansoni*, adjusting for covariates. Total phenotypic variance in (A) *N. americanus* and (B) *S. mansoni* infection intensity is partitioned into that due to covariates, species-specific additive genetic effects, domestic environmental effects (species-specific and those common to both species) and unexplained, individual-specific effects (species-specific and those common to both species).

Table 1Distribution of relative pairs by degree of relationship ($n = 1303$)

Relationship	Degree of relatedness	Co-efficient of relatedness	Number of pairs
Husband - Wife	0	0	554
Parent - offspring	1	0.5	1,373
Sibling	1	0.5	1,281
Other first degree (inc. complex relationships)	1	0.5-0.563	211
Half sibling	2	0.25	243
Grandparent	2	0.25	885
Avuncular	2	0.25	2,292
Other second degree (inc. complex relationships)	2	0.25-0.375	638
Cousin	3	0.125	3,117
Half avuncular	3	0.125	472
Grand avuncular	3	0.125	1,354
Great grand parent	3	0.125	90
Other third degree (inc. complex relationships)	3	0.125-0.219	2,158
Higher-degree relative	4 - 7	0.0039-0.117	26,358
Total	-		40,476

Variance component analysis for variation in infection intensity with *Necator americanus* and *Schistosoma mansoni* showing standardised variance parameter estimates (h^2 additive genetic variation; c^2 household-level variation; e^2 individual-level unexplained variation) and estimated log likelihood values.

Table 2

Hypothesis	N. americanus			S. mansoni			
	h^2	c^2	e^2	h^2	c^2	e^2	
No Covariates:							
Sporadic			1.00	-2174.3		1.00	-1838.0
Shared household		0.34 ± 0.03	0.66 ± 0.03	-2076.5	0.35 ± 0.03	0.65 ± 0.03	-1711.3
Additive genetic	0.48 ± 0.05		0.52 ± 0.05	-2067.7	0.47 ± 0.04	0.53 ± 0.04	-1704.1
Saturated	0.30 ± 0.07	0.15 ± 0.05	0.55 ± 0.04	-2060.8	0.17 ± 0.04	0.54 ± 0.04	-1693.5
With Covariates ^a							
Sporadic			1.00	-1955.3		1.00	-1655.3
Shared household		0.22 ± 0.03	0.78 ± 0.03	-1909.0	0.22 ± 0.03	0.78 ± 0.03	-1610.5
Additive genetic	0.38 ± 0.05		0.62 ± 0.05	-1906.2	0.37 ± 0.05	0.63 ± 0.05	-1605.5
Saturated	0.23 ± 0.06	0.12 ± 0.04	0.65 ± 0.05	-1900.2	0.11 ± 0.04	0.65 ± 0.05	-1600.8

^a *Necator americanus* infection adjusted for age, sex, toilet facilities, overcrowding, sector and Normalised Difference Vegetation Index (NDVI); *S. mansoni* infection adjusted for age (age and age²), toilet facilities and sector.

Table 3

Correlation components for covariance between *Necator americanus* and *Schistosoma mansoni* faecal egg excretion.

Total phenotypic correlation (ρ_P)	0.24
Additive genetic:	
Additive genetic correlation(ρ_G)(SE)	0.147 (0.216)
% of total correlation (ρ_P)	12.8%
<i>P</i> ^b	0.52
Shared-household:	
Household correlation(ρ_C)(SE)	0.451 (0.201)
% of total correlation (ρ_P)	24.5%
<i>P</i> ^b	0.06
Non-genetic individual:	
Unexplained correlation(ρ_E)(SE)	0.233 (0.048)
% of total correlation (ρ_P)	62.6%
<i>P</i> ^b	<<0.001

^a *Necator americanus* infection adjusted for age, sex, toilet facilities, overcrowding, sector and Normalised Difference Vegetation Index (NDVI); *S. mansoni* infection adjusted for age (age and age²), toilet facilities and sector.

^b *P* value for likelihood ratio test comparing general model with restricted model in which correlation for this effect is zero.

Table 4

Covariate and variance estimates from bivariate quantitative genetic analysis of *Necator americanus* and *Schistosoma mansoni* infection intensity expressed as $\ln(\text{epg}+1)$.

	<i>N. americanus</i> $\ln[\text{epg}+1]$		<i>S. mansoni</i> $\ln[\text{epg}+1]$	
Covariate coefficients (SE):				
Demography				
Age ^a	1.04	(0.09)	0.075	(0.009)
Age ²			-0.001	(0.0001)
Sex (female vs male)	-0.49	(0.14)		
Sex × age interaction	-0.27	(0.13)		
Household characteristics				
Toilet (present vs absent)	-0.66	(0.32)	-0.44	(0.21)
Crowding ^b (yes vs no)	0.50	(0.22)		
SES index ^c	-0.23	(0.09)		
Location characteristics				
Remotely sensed data:				
NDVI ^c	2.41	(0.75)		
Sector (vs. sector 1):				
Sector 3	1.50	(0.37)	0.71	(0.26)
Sector 4	1.26	(0.46)	-1.82	(0.34)
Sector 5	-0.90	(0.56)	-2.41	(0.42)
Sector 6	1.49	(0.63)	-1.82	(0.48)
Sector 7	0.23	(0.41)	-1.63	(0.30)
Variance components as proportions of residual variation (SE) after controlling for covariates:				
Additive genetic	0.196	(0.065)	0.230	(0.065)
Shared domestic environment	0.151	(0.043)	0.116	(0.042)
Individual-specific	0.653	(0.047)	0.654	(0.047)

epg, eggs per gram of faeces; SE, standard error.

^aFor *N. americanus*, age was modelled as a linear increase up to 5 years old; for *S. mansoni* age was modelled as a quadratic.

^bA crowded household is defined as having < 1 room/person

^cSocio-economic (SES) index and Normalised Difference Vegetation Index (NDVI) are modelled a continuous variables.