

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Cox, SE; Nweneka, CV; Doherty, CP; Fulford, AJ; Moore, SE; Prentice, AM; (2013) Randomised controlled trial of weekly chloroquine to re-establish normal erythron iron flux and haemoglobin recovery in postmalarial anaemia. *BMJ open*, 3 (7). ISSN 2044-6055 DOI: <https://doi.org/10.1136/bmjopen-2013-002666>

Downloaded from: <http://researchonline.lshtm.ac.uk/1035427/>

DOI: <https://doi.org/10.1136/bmjopen-2013-002666>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: Creative Commons Attribution Non-commercial
<http://creativecommons.org/licenses/by-nc/3.0/>

<https://researchonline.lshtm.ac.uk>

Randomised controlled trial of weekly chloroquine to re-establish normal erythron iron flux and haemoglobin recovery in postmalarial anaemia

Sharon E Cox,^{1,2} Chidi V Nweneka,^{1,3} Conor P Doherty,⁴ Anthony J Fulford,^{1,3} Sophie E Moore,^{1,3} Andrew M Prentice^{1,3}

To cite: Cox SE, Nweneka CV, Doherty CP, *et al*. Randomised controlled trial of weekly chloroquine to re-establish normal erythron iron flux and haemoglobin recovery in postmalarial anaemia. *BMJ Open* 2013;**3**:e002666. doi:10.1136/bmjopen-2013-002666

► Prepublication history and additional material for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2013-002666>).

CVN and SEC contributed equally.

Received 31 January 2013
Revised 11 April 2013
Accepted 30 May 2013

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 3.0 Licence; see <http://bmjopen.bmj.com>

For numbered affiliations see end of article.

Correspondence to

Professor Andrew M Prentice; Andrew.Prentice@lshtm.ac.uk

ABSTRACT

Objective: To determine if low-dose weekly chloroquine (CQ) therapy improves recovery from malaria-associated anaemia.

Design: Proof of concept randomised clinical trial.

Setting: West Kiang District, Lower River Region, The Gambia.

Participants: Children resident in participating communities, aged 12–72 months, with uncomplicated malaria identified using active case detection over two consecutive malaria transmission seasons.

Interventions: In 2007, eligible children were randomised to chloroquine-sulfadoxine/pyrimethamine (CQ-SP) or co-artemether (ACT) antimalarial therapy, and after parasite clearance on day 3 were subsequently re-randomised (double-blind) to weekly low-dose CQ (5 mg/kg) or placebo. In 2008, all eligible children were treated with ACT and subsequently randomised to CQ or placebo.

Outcome measures: The primary outcome was a change in haemoglobin from baseline (day 3 of antimalarial treatment) to day 90 in the CQ and placebo treatment arms. Secondary outcomes were changes in urinary neopterin as a marker of macrophage activation, markers of erythropoietic response and prevalence of submicroscopic parasitaemia. Change in haemoglobin in the placebo arm by initial antimalarial treatment was also assessed.

Results: In 2007, 101 children with uncomplicated malaria were randomised to antimalarial treatment with CQ-SP or ACT and 65 were subsequently randomised to weekly CQ or placebo. In 2008, all children received ACT antimalarial treatment and 31 were subsequently randomised to receive weekly CQ or placebo. Follow-up to day 90 was 96%. There was no effect of weekly CQ vs placebo on change in haemoglobin at day 90 (CQ+10.04 g/L (95% CI 6.66 to 13.42) vs placebo +7.61 g/L (95% CI 2.88 to 12.35)). There was no effect on the secondary outcomes assessed, or effect of initial antimalarial therapy on haemoglobin recovery. Higher day 90 haemoglobin correlated independently with older age, not being stunted, higher haemoglobin at day 0 and adequate iron status at day 3.

ARTICLE SUMMARY

Article focus

- To determine if 3 months of low-dose weekly chloroquine (CQ) (5 mg/kg) therapy after effective treatment of uncomplicated malaria can improve haemoglobin recovery after malaria-associated anaemia in rural African children.
- To test the hypothesis that chloroquine may enhance haematological recovery after a malaria episode through its anti-inflammatory and immuno-modulatory actions, antimacrophageal iron loading effect and antimalarial effect (despite widespread resistance).

Key messages

- Weekly low-dose CQ after effective antimalarial treatment was not effective in improving recovery from mild-to-moderate postmalarial anaemia.
- Weekly low-dose CQ at 5 mg/kg had no clear effects on erythropoietic responses, as determined by reticulocyte percentage, or immune-mediated inflammation as assessed by urinary neopterin, a marker of macrophage activation.

Conclusions: Weekly low-dose CQ after effective antimalarial treatment is not effective in improving recovery from postmalarial anaemia.

Trial registration: The clinical trial registration number is NCT00473837 (ClinicalTrials.gov).

INTRODUCTION

The precise contribution of malaria to anaemia is difficult to calculate mainly because many anaemia-causing conditions coexist in children living in malaria-endemic countries.¹ Murphy and Breman² estimate the annual incidence of severe malarial anaemia (SMA) to be 1.42–5.66 million,

ARTICLE SUMMARY

Strengths and limitations of this study

- The strengths of this study are its randomised design and double blinding of the allocation to weekly CQ or placebo and that few subjects were lost to follow-up. The continuous follow-up and active case detection for all children ensured that malarial infections were detected early and within similar time frames, plus resurgent or new clinical infections were excluded from the analysis of haemoglobin change. A strict malaria case definition was adhered to and the measurement of submicroscopic parasitaemia over time allowed the effects of incomplete treatment to be modelled.
- A limitation of the study was the need to recruit participants over two malaria transmission seasons and the significantly reduced malaria transmission rate in 2008 and thus reduced recruitment, which also appeared to affect the severity of malaria in terms of parasite densities at first detection and haemoglobin concentrations. However, CIs of the effect size in the final multivariable analysis do not include the clinically significant effect size included in the sample size calculation, and thus our observation is unlikely to be due to a type II error as a result of lack of power to detect a true effect.
- It is possible that the relatively low dose of CQ we used was insufficient to mediate the effects that we were testing for and we were unable to conduct pilot studies of CQ drug dosage and duration to determine the potential best treatment regime to test or of the timings of evaluation of possible mechanisms of effect.
- The change in design in 2008 to all children receiving ACT antimalarial treatment did not affect the analysis of the primary outcome. The change did, however, limit our ability to determine the effect of initial antimalaria therapy on haemoglobin response in the placebo group. An effect of antimalarial treatment CQ on Hb when in the context of adequate parasite clearance, as was observed, might have been expected to occur in the early period of recovery.

resulting in between 194 000 and 974 000 deaths in children younger than 5 years of age. SMA increases the risk of death in children with malaria,³ many of whom die shortly after admission before blood transfusion can be started^{4 5} and after discharge in the community.⁶ In addition, blood transfusion, the standard treatment for severe malarial anaemia exposes the child to the risk of transfusion-related infections like HIV and hepatitis.⁷ Among a nationally representative sample of children aged 1–5 years in the Gambia in 1999, 76% had a haemoglobin (Hb) less than 11 g/dL and 15% had Hb less than 7 g/dL.⁸ It is quite likely that most of the cases of moderate-to-severe anaemia were due in great part to malaria.^{9–11} A better understanding of the pathogenesis of malarial anaemia is required to improve its prevention and management.

Multiple immune and non-immune-mediated mechanisms cause anaemia during and after a malarial episode,¹² including the destruction of non-parasitised erythrocytes, which may account for more than 90% of erythrocyte loss.^{13 14} One important component is

inflammation-induced iron delocalisation, which is characterised by sequestration of iron by the monocyte–macrophage system and hepatocytes, resulting in a state of pseudo-iron deficiency, a state that reverses after the inflammation has subsided.¹² The trans-membrane protein ferroportin is the exclusive route for cellular iron efflux and is responsible for releasing absorbed iron from enterocytes and sequestered iron from macrophages.^{12 15 16} Hepcidin, the master regulator of iron metabolism which responds both to iron status and inflammation,¹⁵ blocks ferroportin. Clinically, malaria-induced inflammation has been shown to be associated with a hepcidin-mediated reduction in intestinal absorption of iron.¹⁷ Inflammatory cytokines, particularly interleukin (IL)-6 and IL-1, stimulate hepcidin release,^{18 19} which results both in the reduced iron absorption of dietary iron and impaired macrophageal release of iron, the latter effect being quantitatively much more significant as macrophageal iron recycling and transport to the erythron are 25-fold to 30-fold greater than the influx of dietary iron. In this way, hepcidin is the mediator of the anaemia of chronic disease.¹⁵ Tumour necrosis factor α could also play a role in ferroportin-mediated iron transfer independent of hepcidin.²⁰ Rapid resolution of malaria-induced inflammation could therefore limit the degree and duration of malarial anaemia by reducing the duration of iron delocalisation.

Chloroquine is a cheap and safe antimalarial drug with proven anti-inflammatory properties; it has been shown to reduce iron delocalisation *in vitro*.^{21 22} We hypothesised that chloroquine may enhance haematological recovery after a malarial episode through its anti-inflammatory and immuno-modulatory actions, antimalarial iron loading effect and antimalarial effect (despite widespread resistance). We designed a proof of concept study to investigate the potential use of chloroquine in the management of mild postmalarial anaemia.

MATERIALS AND METHODS**Study design and population**

This study was a population-based randomised controlled trial (RCT) in children aged 12–72 months that was conducted over two malarial seasons in 2007 and 2008 in West Kiang District, Lower River Region, The Gambia. The study was initially designed and conducted as a 2×2 randomised placebo-controlled trial in 2007 but had to be changed, due to changes in the national malaria treatment guidelines, to a two-arm RCT during the second malaria transmission season in 2008. In the 2×2 phase of the study, children with malaria were initially openly randomised to receive a therapeutic dose regime of chloroquine plus sulfadoxine/pyrimethamine (CQ-SP) or co-artemether (ACT) (day 0). After treatment, on day 3, children were subsequently randomised (double blind) to receive weekly chloroquine or placebo until day 90 ([figure 1](#)). In the two-arm phase, all




Figure 1 CONSORT flow diagram describing participants' movements in 2007 and 2008.

enrolled children with malaria were treated with ACT and then randomised and followed up as before (figure 1). A combination of active and passive case detection was used to identify children in participating communities with malaria using a standard case definition of clinical malaria (see below).

The climate of this district is typical of sub-Saharan Africa with a long dry season lasting from November to June and a relatively short rainy season lasting from July to October. Malaria transmission is seasonal, hyperendemic and follows the rainy season, occurring between July and December. The prevalence of children experiencing one or more malarial episodes among a cohort of 1002 children aged between 1 and 6 years under active malaria surveillance during a population-based study in 2003 was 34% (Doherty *et al*, unpublished data). During the study period, standard malaria control activities in the district consisted of insecticide-treated bednet distribution and case finding and treatment. In 2008, the standard first-line antimalarial treatment regimen for uncomplicated malaria in the Gambia was changed from CQ-SP to co-artemether.

Inclusion and exclusion criteria

Eligibility for malaria surveillance and subsequent possible enrolment into the RCT included all children in the participating communities, aged between 12 and

72 months, at the start of the surveillance period and who were not severely wasted, defined as weight-for-height Z scores less than $-3SD$.

Inclusion criteria for enrolment included: (1) uncomplicated malaria at day 0, defined as a history of fever 48 h prior to presentation or a measured temperature of $37.5^{\circ}C$ or more with peripheral parasitaemia (asexual forms of *Plasmodium falciparum* $\geq 500/\mu L$ by microscopy) on day 0 of the study and (2) for randomisation to the postmalarial intervention, children had to have an Hb between 69 and 110 g/L on day 3 with no peripheral parasitaemia.

Children were excluded from the RCT if they were: (1) unable to take oral medications; (2) had features of severe malaria¹; (3) had a known haemoglobinopathy; (4) were enrolled in another project; (5) had already received antimalarial drugs from other sources outside the project team or (6) were prescribed other drugs with potential antimalarial or antianaemic effects such as co-trimoxazole and haematinics.

Following randomisation to weekly CQ or placebo, children were withdrawn from the study if: (1) they developed a second malarial episode or had life-threatening symptoms, more severe disease or serious adverse reactions; (2) left the study area for more than 14 days; (3) received antimalarials from other sources or (4) their Hb fell below 70 g/L.

Chloroquine in postmalaria anaemia

Field and clinical procedures

To effect the surveillance protocol, the study villages were divided into geographically adjacent clusters supervised by a resident study nurse who visited each village morning and evening and provided 24 h clinical coverage to all enrolled children in their cluster. The active surveillance consisted of twice weekly temperature monitoring of enrolled children by village assistants, while the passive surveillance consisted of mothers referring sick children to the visiting study nurse. Children with fever—reported or measured—were screened for malaria using a rapid diagnostic test (MRDT; Cortez diagnostics, Inc) after clinical evaluation. Children testing negative to MRDT were treated based on the likely diagnosis from the symptoms and signs. Children testing positive to MRDT were started on antimalarials immediately (either CQ-SP or ACT in 2007, and ACT in 2008 for 3 days) and directly observed by the study nurse and managed as appropriate. Finger prick blood samples (days 7, 15, 30 and 70) were collected in an EDTA microtainer while venous blood (2 mL—days 0 and 90; and 5 mL—days 3 and 45) samples were collected in an EDTA vacutainer and sent to the study laboratory on the same day. Comorbidities were treated as appropriate, avoiding co-trimoxazole and haematinics. For the subsequent visits after initial malaria diagnosis at day 0, children were seen by the study nurse and blood samples and clinical data collected as per protocol. Only children who fulfilled the inclusion criteria for advancement to the weekly chloroquine/weekly placebo phase of the study were sampled after day 3. These children were also followed up on days 15, 30, 45, 70 (except in 2008) and 90.

Randomisation and blinding

The randomisation to malaria treatment group in 2007 (CQ-SP or ACT) was open (non-blinded to trial staff, investigators or participants) and was carried out by sequential assignment as children were identified. The block randomisation to the postmalaria treatment of weekly CQ or placebo in both 2007 and 2008 was double blinded and was carried out in blocks of eight. The randomisation codes were generated by a staff member independent of the study team and held by the external trial monitor. Treatment codes were labelled A to H and placed in sequentially numbered, opaque sealed envelopes held by the study nurses. Allocation to the treatment was by matching the code in the envelope to a bottle of the intervention labelled with the same code and then labelled with the subject ID.

Interventions

Children openly randomised to malaria treatment at day 0 with CQ-SP received chloroquine syrup (Chlormal, Medreich Sterilab Ltd, Avalahalli, Bangalore, India) consisting of 50 mg chloroquine base per 5 mL as chloroquine phosphate at a dose of 10 mg/kg for 3 days (days 0–2). SP (Micro Laboratories, India) was

co-administered as a 525 mg tablet (500 mg sulfadoxine and 25 mg pyrimethamine) on day 0 only at a dose of half tablet/10 kg. Children who were randomised to ACT treatment in 2007 and all children in 2008 received co-artemether (Coartem) tablets (Beijing Novartis Pharma Ltd, Beijing China for Novartis Pharma AG, Basle, Switzerland) as blister packets of six tablets per card with each tablet containing 20 mg artemether and 120 mg lumefantrine. The treatment dose comprised a stat dose on day 0 and then 8 h later, followed by twice daily doses for the next 2 days with doses depending on body weight (5 for <15 kg, 1 tablet/dose; 15 for <25 kg, 2 tablets/dose; 25 for <35 kg, 3 tablets per dose).

During the second blinded phase of the trial, children received either weekly CQ syrup (as above) but at a dose of 5 mg/kg or matching placebo syrup (prepared by the Royal Victoria Teaching Hospital pharmaceutical laboratory in collaboration with Atlantic Pharmaceuticals Limited, Latrikunda Sabiji, The Gambia) The syrups were in similar amber-coloured bottles with matching caps and labels. Doses were administered in the child's home under direct observation by the study nurses who waited for at least 10 min after administering the drugs to ensure that they were not vomited.

Primary and secondary outcomes

The primary outcome measure was Hb change from day 3 post-treatment of the malarial episode to day 90 in the weekly CQ and placebo arms.

The secondary outcomes were to compare between the weekly CQ and placebo arms: (1a) changes in erythropoietic response between days 3 and 30; and (1b) between days 3 and 90; (2) changes in urinary neopterin from days 3 to 30 as a marker of macrophage activation; (3) prevalence of submicroscopic malaria parasitaemia using PCR at each time point and (4) Hb change from days 3 to 90 in the two placebo arms to investigate the effects of initial malaria treatment therapy.

Sample size

During the initial study design, we used a working assumption of an SD for change in Hb of 15 g/L.²³ With power 80% and significance level set at 5%, we estimated the sample sizes required to detect varying effect sizes for the primary outcome. A sample size of 65 per arm was estimated to be able to detect an effect size of 7.5 g/L. This sample size was thought to be achievable and the effect size of likely public health significance.

Laboratory procedures

Table 1 describes the different laboratory analyses conducted. Zinc protoporphyrin (ZnPP) as a marker of iron status was measured on fresh whole blood using a fluorometer (AVIV Biomedicals, Lakewood, New Jersey, USA). Full blood count (FBC), haemoglobin and automated reticulocyte count were performed using a Cell Dyne 3700 (Abbott Laboratories, Abbott Park, Illinois, USA). Iron deficiency was defined using three

Table 1 Laboratory investigations conducted

Investigations	Specimen	Days collected*	Purpose
Microscopy for malaria parasite	Blood	0, 3, 7, 15, 30, 45, 70 and 90	To assess parasitaemia
Erythrocyte zinc protoporphyrin	Blood	0 and 3	To assess iron status
Full blood count	Blood	0, 3, 7, 15, 30, 45, 70 and 90	To assess Hb change over time
Automated reticulocyte percentage	Blood	3, 7, 15, 30, 45, 70 and 90	To monitor bone marrow response
Quantitative PCR for <i>Plasmodium falciparum</i> parasites	Blood	0, 3, 7, 15, 30, 45, 70 and 90	To assess submicroscopic parasitaemia
Urinary neopterin concentration	Urine	3, 15 and 30	To assess macrophage activation as measure of immune-mediated inflammation

*No blood was collected on days 7 and 70 in 2008. Hb, haemoglobin.

parameters: mean cell volume (MCV), mean cell haemoglobin (MCH) and ZnPP. The cut-off values for the different parameters employed were: $\geq 61 \mu\text{mol/mol}$ of Hb for ZnPP,²⁴ $\leq 73 \text{ fL}$ for MCV²⁵ and $\leq 25 \text{ pg}$ for MCH.²⁵ Combining these parameters has been shown to increase their sensitivity and specificity to identify iron deficiency.²⁶ Malarial parasites were identified in a thick peripheral blood smear stained with Giemsa overnight and read by a light microscope with $\times 100$ magnification in oil immersion. Parasite DNA extraction and subsequent quantitative PCR amplification were carried out using established protocols.²⁷ In brief, DNA extraction was carried out using the QIamp DNA Mini Kit (Qiagen Ltd, UK) according to the manufacturer's instructions with minor modifications. Extracted DNA samples were stored at -20°C . Quantification of parasite density was performed with a TaqMan assay employing a fluorogenic probe to detect specific parasite products and using an Opticon 2 Real Time PCR machine and Opticon V.3.1 software (Bio-Rad Laboratories, Inc). Primers and probes were synthesised by Applied Biosystems, UK. Urinary neopterin levels were measured using an ELISA technique (BRAHMS Neopterin EIA, B.R.A.H.M.S., Germany). The stored urine was allowed to thaw at room temperature and then diluted 1:100, and the rest of the procedure was according to the manufacturer's instructions.

Statistical analysis

Delta haemoglobin summary statistics are presented as means and SDs for normally distributed continuous variables, and counts and percentages for discrete variables. Neopterin and parasite count were not normally distributed and were log transformed (\log_e for neopterin and \log_{10} for parasite count). The geometric means and 95% CIs are presented for log-transformed variables. Two-sided Student *t* tests were used to compare results between two treatment arms.

Visual inspection of Hb change between baseline at day 3 and later time points by treatment arm suggested that, by day 30, Hb change in both treatment arms had reached a plateau. To fully determine the effect of treatment group on Hb change, we used random effects

models using the repeat measures at days 30 and 90. Initial malaria treatment group (CQ-SP vs ACT), Hb at recruitment (day 0 Hb), baseline Hb (day 3), age in months, year of study, and the child's village were included in the models 'a priori' and hence adjusted for.

Linear regression was used to investigate further potential predictors of Hb change in the study population. First, simple linear regression was used to assess the association of each predictor variable with Hb. Thereafter, all variables with a *p* value of 0.1 or less were included in a multiple linear regression to assess confounding. The final regression model consisted of only the variables that maintained a significant association with Hb in the multiple regression models. All analyses were carried out using STATA V.8 (StataCorp, College Station, Texas, USA).

Ethical clearance

The Scientific Coordinating Committee of the Medical Research Council (UK), The Gambia, and the MRC/Gambia Government Joint Ethical Committee approved the study (SCC 1076). Written informed consent was obtained from parents/guardians of the children prior to enrolment. The clinical trial registration number is NCT00473837 (ClinicalTrials.gov).

RESULTS

Characteristics of the study population under malaria surveillance

Figure 1 describes the flow of participants through the study and the reasons for exclusions from the study. In 2007, 1445 of 1817 (80%) potentially eligible children identified through the Keneba MRC DSS database from 14 villages were included in the malaria surveillance. In 2008, 1220 of the 1413 (86%) identified children from five of the original villages were included in the malaria surveillance (smaller villages were excluded as they had contributed few cases in 2007). Table 2 describes the characteristics of the children placed under surveillance for malaria. The proportion of children with a positive malaria rapid test during surveillance (only first event per child counted) fell from 7.3% (105/1445) in 2007

Chloroquine in postmalarial anaemia

Table 2 Baseline characteristics of all participants under surveillance

	2007 N=1445	2008 N=1220	Combined N=2665
Mean age in months (SD)	42.2 (17.2)	40.7 (17.7)	41.5 (17.5)
Mean weight-for-height Z scores (SD)	-1.06 (0.84)	-1.02 (0.85)	-1.04 (0.84)
Mean weight-for-age Z scores (SD)	-1.48 (0.97)	-1.50 (0.94)	-1.49 (0.95)
Mean height-for-age Z scores (SD)	-1.06 (1.21)	-1.17 (1.13)	-1.11 (1.17)
Mean body mass index (kg/m ²) (SD)	14.6 (1.3)	14.7 (1.4)	14.6 (1.4)
Prevalence of a malaria rapid test positive result during surveillance*	105 (7.3)	49 (4.0)	154 (5.8)

*Only the first positive result is counted.

to 4% (49/1220) in 2008. In the first year of the study, 101 (96%) of the children with a positive malaria rapid test were randomised to CQ-SP or ACT treatment, and 65 (64%) subsequently randomised at day 3 to weekly chloroquine or placebo, with all children completing follow-up until day 30 and 33/34 completing follow-up until day 90 in the weekly CQ and 28/31 in the weekly placebo group. In the second year of the study, 31 of the 49 (63%) cases with a first positive malaria rapid test were subsequently randomised to weekly CQ or placebo after treatment with ACT, all of whom completed follow-up until day 90 (16 CQ and 15 placebo). Thus, 96 children were included in the analysis of the primary and secondary outcomes, except for the analysis of effect of initial malaria treatment within the placebo arms which included 31 children.

Table 3 describes the characteristics of participants randomised to weekly chloroquine and weekly placebo for both years combined. Demographic and anthropometric characteristics were similar between the groups as was baseline Hb at day 3. The prevalence of iron deficiency measured at baseline was 43.1% (28/65) in 2007 and 58.1% (18/31) in 2008 with a similar prevalence between the CQ/placebo groups within each year (data not shown) and in both years combined. However, parasite density was lower in the placebo group (table 3). No adverse events were reported in either treatment group.

Effect of weekly CQ on haemoglobin change during follow-up

To assess the effect of weekly chloroquine on Hb recovery after clearance of peripheral parasitaemia, we compared change in Hb from days 3 to 90 and from days 3 to 30 in the two arms. No significant differences in Hb change between the two groups were observed, either for the data for both study years or for each study year separately (both years combined: change in Hb at day 90, CQ +10.04 g/L (95% CI 6.66 to 13.42) vs placebo +7.61 g/L (95% CI 2.88 to 12.35); figure 2). Thus, the effect size of the CQ treatment was 2.43 g/L (95% CI -3.38 to 8.24) compared to the 7.5 g/L we included in the sample size calculation. However, at day 15, children receiving weekly CQ had an increase of 3.5 g/L (95% CI 0.5 to 6.4) compared to 0.8 g/L (95% CI -2.3 to 3.9) in children in the placebo arm, but this difference did not reach statistical significance (p=0.2). The lack of effect of weekly CQ remained when adjusting for factors determined 'a priori' (Hb at recruitment (day 0), Hb at baseline (day 3), age (in months), year of study, initial treatment received (CQ-SP vs ACT) and village group) using a random effects repeated measures model including Hb measured at days 30 and 90 (coefficient=1.43 (95% CI -1.70 to 4.57)). Furthermore, adding parasite density at recruitment, presence of submicroscopic parasitaemia at any time point during follow-up, and

Table 3 Baseline characteristics of children randomised to weekly chloroquine and weekly placebo in 2007 and 2008, pooled

Parameter	Chloroquine group N=50	Placebo group N=46
Male sex (%)	28 (56.0)	24 (52.2)
Age in months, mean (SD)	41.9 (18.1)	38.7 (17.8)
Height-for-age Z scores, mean (SD)	-1.19 (0.91)	-0.94 (1.20)
Weight-for-age Z scores, mean (SD)	-1.44 (0.95)	-1.35 (0.87)
Weight-for-height Z scores, mean (SD)	-0.94 (0.90)	-0.96 (0.79)
BMI (kg/m ²), mean (SD)	14.7 (1.4)	14.7 (1.3)
Parasite density at recruitment (day 0), geometric mean, (95% CI)	47 783 (31 617 to 71 120)	32 496 (21 910 to -47 507)
Hb (g/L) at recruitment (day 0), mean (SD)	103.9 (14.2)	101.5 (14.8)
Hb (g/L) at baseline (day 3), mean (SD)	93.4 (9.6)	93.7 (10.2)
Iron deficient* at baseline (day 3) (%)	23 (46)	23 (50)

*Iron deficiency was defined as zinc protoporphyrin ≥ 61 $\mu\text{mol/mol}$ of Hb, mean cell volume ≤ 73 fL and mean cell haemoglobin ≤ 25 pg.




Figure 2 Mean change in haemoglobin concentration at days 30 and 90 from baseline at day 3 by treatment group for each year of study and for both years combined. Error bars represent the 95% CIs. Change in haemoglobin at day 3 is compared to haemoglobin concentration at presentation with malaria and initiation of antimalarial treatment. Thereafter, the comparison is with baseline levels at day 3 after completion of antimalarial treatment. Error bars represent the 95% CIs.

presence of iron deficiency had no effect on the estimates of effect for CQ vs placebo treatment during follow-up (data not shown).

Predictors of haemoglobin at day 90

In the absence of an effect of weekly CQ on Hb change after successful malaria treatment, we assessed what other factors might determine final Hb. Using simple linear regression, significant positive associations were observed for parasite density at day 0, better nutritional status, age, Hb at day 0 at malaria diagnosis and day 3 of successful treatment. Presence of iron deficiency at baseline was associated with significantly lower final Hb. There was no apparent effect of initial malaria treatment (CQ-SP vs ACT; [table 4](#)). In a final multiple regression model, Hb at day 0 was the strongest predictor of final

Hb and other independent predictors of final haemoglobin were presence of iron deficiency at day 3, degree of stunting (height-for-age Z score), and age (in months). Children with higher Hb at diagnosis and older children had greater final Hb. Children with iron deficiency at day 3 had significantly lower final Hb with a mean difference of 12.1 g/L ([table 5](#)).

Effect of initial antimalaria treatment (CQ-SP vs ACT) on haemoglobin change

A secondary outcome of this study was Hb change from days 3 to 30 and day 90 in the placebo arm to determine whether initial antimalaria treatment (CQ-SP or ACT) treatment affected macrophage iron loading in acute clinical malaria and its subsequent release. As indicated in the analysis of Hb at day 90, there was no significant effect of initial antimalarial treatment. Please see online supplementary figure S1 depicting Hb change during follow-up from days 0 to 90 in the placebo arm by initial antimalarial treatment received and the accompanying text.

Submicroscopic parasitaemia, inflammation, bone marrow response, haemoglobin change and treatment groups

Submicroscopic parasitaemia was commonest in the first 7 days of acute phase follow-up (days 3–15). By day 30, most were negative (77%). The prevalence of submicroscopic parasitaemia at days 30 and 90 combined was 15.1% and there was no evidence of an effect of the post-malaria treatment group or of initial antimalarial treatment ([figure 3](#) and see online supplementary figure S2). Most of the submicroscopic parasitaemia at these time points was of a very low level (1–2 parasites/ μ L). However, higher parasite density at diagnosis was associated with an increased likelihood of subsequent submicroscopic parasitaemia at days 30 or 90 with a one-log increase in parasite count increasing the odds of having submicroscopic parasitaemia by 1.6 ($p=0.005$). Although the presence/absence of submicroscopic parasitaemia

Table 4 Predictors of final haemoglobin at day 90 using univariable linear regression

Parameter	Regression coefficient (95% CI)	p Value
Parasite count at recruitment (day 0)	3.5 (1.1 to 5.8)	0.004
Sex	-0.4 (-3.3 to 2.5)	0.8
Iron deficiency at baseline (day 3)	-12.1 (-14.7 to -9.5)	<0.001
Height-for-age Z scores (HAZ) <-2	3.2 (1.8 to 4.6)	<0.001
Weight-for-age Z scores (WAZ) <-2	2.6 (1.0 to 4.2)	0.001
Weight-for-height (WHZ) Z scores <-2	0.1 (-1.6 to 1.8)	0.9
Age in months	0.4 (0.3 to 0.4)	<0.001
Initial antimalaria treatment arm	1.2 (-2.1 to 4.4)	0.5
CQ vs placebo randomisation group	-0.5 (-3.4 to 2.5)	0.8
Hb (g/L) at recruitment (day 0)	0.6 (0.5 to 0.6)	<0.001
Hb (g/L) at baseline (day 3)	0.7 (0.6 to 0.8)	<0.001
Village	1.9 (0.9 to 3.0)	<0.001
Presence of sub-microscopic parasitaemia at any time point during follow-up	0.6 (-7.1 to 8.2)	0.9
Year of study	-1.3 (-4.4 to 1.8)	0.4

Chloroquine in postmalarial anaemia

Table 5 Independent predictors of final haemoglobin (day 90) using multivariable linear regression

Parameter	Regression coefficient (95% CI)	p Value
Day 0 Hb	0.44 (0.36 to 0.52)	<0.001
Presence of iron deficiency	-3.0 (-5.57 to -0.43)	0.022
Height-for-age Z score <-2	1.48 (0.41 to 2.56)	0.007
Age in months	0.21 (0.14 to 0.28)	<0.001

Model: adjusted $r^2=0.46$, $F=76.6$, $p>F<0.001$.
Hb, haemoglobin.

during follow-up (days 30 and 90) was not associated with final Hb (tables 4 and 5), in children with high qPCR (quantitative PCR parasite count) parasite count (>500/ μ L, N=3) at day 30, Hb at day 30 was significantly lower compared to children with 1–2 parasites/ μ L or none at day 30 (82.3 vs 102.8 g/L; $p=0.01$). Similarly, the mean Hb change was also reduced in those with persistent high parasitaemia at day 30 (-5.0 vs 8.5 g/L; mean Hb difference (95% CI) 13.5 g/L (-0.3 to 27.3); $p=0.06$). Bone marrow response as assessed by the percentage of reticulocytes peaked at day 15 (4.5% in both treatment groups combined), compared to similar levels of around 2% at all other measured time points (days 3, 7, 15 and 90). There was no evidence of an effect of either the postmalaria treatment group (figure 4) or of the initial antimalarial therapy (data not shown) on bone marrow response. Urinary neopterin, a product of activated macrophages and therefore a marker of a pro-inflammatory immune status, was strongly associated with qPCR parasite count at day 30 (regression coefficient: 0.2 (95% CI 0.08 to 0.33), $p=0.003$; adjusted $R^2=0.4091$) and was also associated with Hb at day 30. There was no evidence of an effect of the postmalaria treatment group (figure 5) or of the initial antimalarial treatment on neopterin concentrations (see online supplementary figure S3).




Figure 3 Percentage of participants with submicroscopic malaria (qPCR) over time by postmalarial treatment group.

DISCUSSION

Chloroquine is used in the treatment of several diseases in addition to its antimalarial use.^{28 29} However, we believe this to be the first study to assess its potential role in the management of postmalarial anaemia. We hypothesised that since malarial anaemia is essentially a systemic inflammatory response to infection,¹² chloroquine, by suppressing plasmodium-induced inflammation, could enhance erythropoietic recovery. We tested this hypothesis by comparing two groups of children—one group receiving weekly chloroquine and the other group weekly placebo after complete clearance of peripheral parasitaemia as determined by microscopy. The primary outcome measure was the recovery in Hb at days 30 and 90 compared to the day 3 Hb levels. Our results showed no significant difference in the Hb change between the two groups. The data were interrogated further by a random effects model modelling data at days 30 and 90 simultaneously while adjusting for Hb at day 3 and taking into account random measurement errors and missing data. This confirmed that weekly chloroquine at 5 mg/kg body weight did not improve Hb recovery after uncomplicated clinical malaria. There was a suggestion that weekly CQ may have increased the rate of recovery in Hb change at day 15 accompanied by non-significantly increased reticulocyte percentage.

A limitation of the study was the need to recruit participants over two malaria transmission seasons. In 2008, there was a reduced malaria transmission rate, and thus there were fewer cases available for recruitment and also an apparent effect on the severity of malaria in terms of parasite densities at first detection and effect on




Figure 4 Reticulocyte percentage as an indicator of bone marrow response by treatment group for both years combined.




Figure 5 Urinary neopterin by postmalarial treatment group.

haemoglobin concentrations, with a significantly higher baseline Hb in 2008. However, the change in design that was necessitated in 2008 did not change the primary outcome analysis. Furthermore, CIs of the observed effect size in the final multivariable analysis do not include the effect size included in the sample size calculation, and thus our observation is unlikely to be due to a type II error as a result of lack of power to detect a true effect of this magnitude.

The change in design in 2008 to all children receiving ACT antimalarial treatment did not affect the analysis of the primary outcome. The change did, however, limit our ability to determine the effect of initial antimalarial therapy on haemoglobin response in the placebo group. An effect of antimalarial treatment CQ on Hb when in the context of adequate parasite clearance, as was observed, might have been expected to occur in the early period of recovery.

The recommended anti-inflammatory dose of CQ is 3–4 mg/kg/day. However, owing to a lack of safety data for this condition and considering the potential adverse effects, particularly retinal damage, we used the lower prophylactic dose of 5 mg/kg/week. Using this low dose might have contributed to our inability to detect an effect of CQ. Drenou *et al*³⁰ reported the successful long-term control of sideroblastic anaemia (a condition of reduced incorporation of iron into haem in the presence of sufficient or high iron) in an adult patient using chloroquine but at 100 mg/day.

We had hypothesised that chloroquine will impede macrophageal iron sequestration during the acute clinical episode and improve iron availability for the process of erythropoiesis. Urinary excretion of neopterin is markedly increased in individuals infected with malaria, and corresponds to the concurrent activation of T cells and macrophages.^{31–33} We measured the urinary neopterin at days 3, 15 and 30 and expected to see a more rapid decline in the neopterin level in the weekly chloroquine group than in the weekly placebo group, but no difference was observed between the treatment groups. It is possible that a difference may have occurred at an intermediate time point between days 3 and 15,

when the levels had already almost normalised. Indeed, Brown *et al*³³ showed that in a group of volunteers experimentally infected with *plasmodium* parasites, the timing of maximum neopterin excretion coincided with the treatment and elimination of parasitaemia. In our study, this would coincide with study days 0–3. In future, it might be more informative to carry out more measurements during the first week of malaria diagnosis.

The participants were followed up for 90 days because the reports had suggested that Hb continues to fall even after successful malaria treatment primarily due to continuing inflammation.^{34 35} In this study, we found no evidence of persistent inflammation, except in the few cases with remaining parasitaemia at day 30, which also had decreased absolute Hb and Hb change at day 30 compared with children with no parasitaemia or parasite counts of 1–2/ μ L.

Our observation that children with higher Hb at diagnosis had greater Hb recovery is potentially paradoxical as one would have expected the reverse through statistical regression to the mean and a greater ‘catch up’ in children with a lower starting Hb at day 3. However, as the majority of iron used for Hb recovery is recycled,²⁴ those with higher Hb at day 0 have more iron available for recovery. Our non-provision of exogenous iron during the study may have restricted the ability of the more anaemic children to respond appropriately, as suggested by our observation of reduced Hb recovery in children with apparent iron deficiency at day 0. An earlier study in Gambian children showed improved haematological recovery in children treated for malaria and cosupplemented with iron compared to placebo.³⁶ However, we have also shown in Gambian children reduced iron absorption from supplementation in postmalarial anaemia, while also experiencing greater Hb recovery compared to supplemented children with iron-deficiency anaemia, thus causing us to question the efficacy of iron supplementation given within 2 weeks of antimalarial treatment.²⁴ Without the inclusion of a further randomised treatment arm of postmalarial treatment iron supplementation, it is not possible to make further recommendations concerning the recent argument for the provision of iron supplementation to children in malaria endemic areas with regular malaria surveillance and treatment facilities.³⁷

Our finding that stunted children recovered Hb poorly is consistent with an earlier report of stunting being a determinant of the severity of malaria-associated anaemia.³⁸ In our population, this effect was independent of iron status.

Age has been shown to be strongly associated with risk of severe anaemia in children with acute malaria, the risk decreasing with increasing age.³⁹ In addition, we have shown in this study that older children recover better from anaemia after an acute malaria episode than younger children. Paying special attention to younger children with malarial anaemia could reduce malaria-related mortality in children.

Chloroquine in postmalarial anaemia

In conclusion, weekly chloroquine administered for 3 months to children with mild malarial anaemia at a dose of 5 mg/kg/week resulted in no improvement in erythropoietic recovery compared to placebo. Older children and children with higher initial Hb, less stunting at diagnosis and without iron deficiency had better Hb recovery postmalaria. Similar to other reports in the literature, urinary neopterin was elevated during acute malaria. The role of iron deficiency in erythropoietic recovery postmalaria noted in this study suggests areas for further detailed investigation of the clinical value in administering both antimalarial and iron supplementation to children with malarial anaemia residing in areas of high iron deficiency and malaria burden.

Author affiliations

¹Medical Research Council (MRC) International Nutrition Group, London School of Hygiene & Tropical Medicine, London, UK

²Muhimbili Wellcome Programme, Muhimbili University of Health & Allied Sciences, Dar es Salaam, Tanzania

³MRC Keneba, Keneba, The Gambia

⁴Department of Paediatrics, Royal Hospital for Sick Children, Glasgow, UK

Acknowledgements The authors wish to acknowledge the contribution of all the members of the field team—Kabiru Ceesay, Edrisa Sinjanka, Lamin Ceesay, Lamin Sowe, Kalilu Sanneh, Bakary Sarr, Muhammed Sidibeh, Yahya Sanyang; the data staff, especially Jahid Hassan and Musa Jarjou; the laboratory staff, especially Musa Colley, Karamo Camara, Mustapha Bittaye and Ebrimah Danso; all the drivers of the MRC Keneba, Ms Cheryl Eniyou O, Mrs Fatou Noho and Ms Jankey Jagne; all the village assistants; and the study subjects and their parents.

Contributors CVN, CPD and AMP conceived and designed the study with input from SEC. CVN and SEM were responsible for the study conduct and data acquisition. CVN, SEC and AJF were responsible for the statistical analysis and interpretation. SEC and CVN drafted the paper and circulated it to all authors for critical revision. All authors read and approved the final manuscript.

Funding This study was supported by the UK Medical Research Council (MC-A760-5QX00).

Competing interests None.

Ethics approval The Scientific Coordinating Committee of the Medical Research Council (UK), The Gambia, and the MRC/Gambia Government Joint Ethical Committee approved the study (SCC 1076).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The original data set is available on request to the corresponding author and permission is dependent on the nature and purpose of intended use.

REFERENCES

- World Health Organization. Severe falciparum malaria. World Health Organization, Communicable Diseases Cluster. *Trans R Soc Trop Med Hyg* 2000;94(Suppl 1):S1–90.
- Murphy SC, Breman JG. Gaps in the childhood malaria burden in Africa: cerebral malaria, neurological sequelae, anemia, respiratory distress, hypoglycemia, and complications of pregnancy. *Am J Trop Med Hyg* 2001;64(1–2 Suppl):57–67.
- Brabin BJ, Premji Z, Verhoeff F. An analysis of anemia and child mortality. *J Nutr* 2001;131(2S–2):636S–45S; discussion 46S–8S.
- Bojang KA, Palmer A, Boele van Hensbroek M, et al. Management of severe malarial anaemia in Gambian children. *Trans R Soc Trop Med Hyg* 1997;91:557–61.
- Mulenga M, Malunga F, Bennett S, et al. A randomised, double-blind, placebo-controlled trial of atovaquone-proguanil vs. sulphadoxine-pyrimethamine in the treatment of malarial anaemia in Zambian children. *Trop Med Int Health* 2006;11:1643–52.
- Phiri KS, Calis JC, Faragher B, et al. Long term outcome of severe anaemia in Malawian children. *PLoS ONE* 2008;3:e2903.
- Lackritz EM. Prevention of HIV transmission by blood transfusion in the developing world: achievements and continuing challenges. *AIDS* 1998;12(Suppl A):S81–6.
- NaNA. *Nationwide survey on the prevalence of Vitamin A and Iron deficiency in women and children in The Gambia*. National Nutrition Survey. Banjul: National Nutrition Agency/Medical Research Council, 2001.
- Abdalla SH. Iron and folate status in Gambian children with malaria. *Ann Trop Paediatr* 1990;10:265–72.
- Bojang KA, Van Hensbroek MB, Palmer A, et al. Predictors of mortality in Gambian children with severe malaria anaemia. *Ann Trop Paediatr* 1997;17:355–9.
- Brewster DR, Greenwood BM. Seasonal variation of paediatric diseases in The Gambia, West Africa. *Ann Trop Paediatr* 1993;13:133–46.
- Nweneka CV, Doherty CP, Cox S, et al. Iron delocalisation in the pathogenesis of malarial anaemia. *Trans R Soc Trop Med Hyg* 2010;104:175–84.
- Evans KJ, Hansen DS, van Rooijen N, et al. Severe malarial anemia of low parasite burden in rodent models results from accelerated clearance of uninfected erythrocytes. *Blood* 2006;107:1192–9.
- Price RN, Simpson JA, Nosten F, et al. Factors contributing to anemia after uncomplicated falciparum malaria. *Am J Trop Med Hyg* 2001;65:614–22.
- Ganz T. Molecular pathogenesis of anemia of chronic disease. *Pediatr Blood Cancer* 2006;46:554–7.
- Ganz T. Hepcidin and its role in regulating systemic iron metabolism. *Hematology Am Soc Hematol Educ Program* 2006:29–35, 507.
- Prentice AM, Doherty CP, Abrams SA, et al. Hepcidin is the major predictor of erythrocyte iron incorporation in anemic African children. *Blood* 2012;119:1922–8.
- Nemeth E, Rivera S, Gabayan V, et al. IL-6 mediates hypoferrremia of inflammation by inducing the synthesis of the iron regulatory hormone hepcidin. *J Clin Invest* 2004;113:1271–6.
- Lee P, Peng H, Gelbart T, et al. Regulation of hepcidin transcription by interleukin-1 and interleukin-6. *Proc Natl Acad Sci USA* 2005;102:1906–10.
- Atkinson SH, Rockett KA, Morgan G, et al. Tumor necrosis factor SNP haplotypes are associated with iron deficiency anemia in West African children. *Blood* 2008;112:4276–83.
- Legssyer R, Ward RJ, Crichton RR, et al. Effect of chronic chloroquine administration on iron loading in the liver and reticuloendothelial system and on oxidative responses by the alveolar macrophages. *Biochem Pharmacol* 1999;57:907–11.
- Ghigo D, Aldieri E, Todde R, et al. Chloroquine stimulates nitric oxide synthesis in murine, porcine, and human endothelial cells. *J Clin Invest* 1998;102:595–605.
- Atkinson SH, Rockett K, Sirugo G, et al. Seasonal childhood anaemia in West Africa is associated with the haptoglobin 2–2 genotype. *PLoS Med* 2006;3:e172.
- Doherty CP, Cox SE, Fulford AJ, et al. Iron incorporation and post-malaria anaemia. *PLoS ONE* 2008;3:e2133.
- Worwood M. The laboratory assessment of iron status—an update. *Clin Chim Acta* 1997;259:3–23.
- Hershko C, Bar-Or D, Gaziel Y, et al. Diagnosis of iron deficiency anemia in a rural population of children. Relative usefulness of serum ferritin, red cell protoporphyrin, red cell indices, and transferrin saturation determinations. *Am J Clin Nutr* 1981;34:1600–10.
- Nwakanma DC, Gomez-Escobar N, Walther M, et al. Quantitative detection of Plasmodium falciparum DNA in saliva, blood, and urine. *J Infect Dis* 2009;199:1567–74.
- Cooper RG, Magwere T. Chloroquine: novel uses & manifestations. *Indian J Med Res* 2008;127:305–16.
- Wallace DJ. The use of chloroquine and hydroxychloroquine for non-infectious conditions other than rheumatoid arthritis or lupus: a critical review. *Lupus* 1996;5(Suppl 1):S59–64.
- Drenou B, Guyader D, Turlin B, et al. Treatment of sideroblastic anemia with chloroquine. *N Engl J Med* 1995;332:614.
- Reibnegger G, Boonpucknavig V, Fuchs D, et al. Urinary neopterin is elevated in patients with malaria. *Trans R Soc Trop Med Hyg* 1984;78:545–6.
- Fuchs D, Hausen A, Reibnegger G, et al. Neopterin as a marker for activated cell-mediated immunity: application in HIV infection. *Immunol Today* 1988;9:150–5.

33. Brown AE, Herrington DA, Webster HK, *et al*. Urinary neopterin in volunteers experimentally infected with *Plasmodium falciparum*. *Trans R Soc Trop Med Hyg* 1992;86:134–6.
34. Dondorp AM, Nyanoti M, Kager PA, *et al*. The role of reduced red cell deformability in the pathogenesis of severe falciparum malaria and its restoration by blood transfusion. *Trans R Soc Trop Med Hyg* 2002;96:282–6.
35. Biemba G, Gordeuk VR, Thuma PE, *et al*. Prolonged macrophage activation and persistent anaemia in children with complicated malaria. *Trop Med Int Health* 1998;3:60–5.
36. van Hensbroek MB, Morris-Jones S, Meisner S, *et al*. Iron, but not folic acid, combined with effective antimalarial therapy promotes haematological recovery in African children after acute falciparum malaria. *Trans R Soc Trop Med Hyg* 1995;89:672–6.
37. Ojukwu JU, Okebe JU, Yahav D, *et al*. Oral iron supplementation for preventing or treating anaemia among children in malaria-endemic areas. *Cochrane Database Syst Rev* 2009(3):CD006589.
38. Verhoef H, West CE, Veenemans J, *et al*. Stunting may determine the severity of malaria-associated anemia in African children. *Pediatrics* 2002;110:e48.
39. Reyburn H, Mbatia R, Drakeley C, *et al*. Association of transmission intensity and age with clinical manifestations and case fatality of severe *Plasmodium falciparum* malaria. *JAMA* 2005;293:1461–70.

BMJ Open

Randomised controlled trial of weekly chloroquine to re-establish normal erythron iron flux and haemoglobin recovery in postmalarial anaemia

Sharon E Cox, Chidi V Nweneka, Conor P Doherty, et al.

BMJ Open 2013 3:

doi: 10.1136/bmjopen-2013-002666

Updated information and services can be found at:

<http://bmjopen.bmj.com/content/3/7/e002666.full.html>

These include:

Data Supplement

"Supplementary Data"

<http://bmjopen.bmj.com/content/suppl/2013/07/05/bmjopen-2013-002666.DC1.html>

References

This article cites 36 articles, 17 of which can be accessed free at:

<http://bmjopen.bmj.com/content/3/7/e002666.full.html#ref-list-1>

Open Access

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license. See:

<http://creativecommons.org/licenses/by-nc/3.0/> and

<http://creativecommons.org/licenses/by-nc/3.0/legalcode>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Epidemiology](#) (635 articles)

[Haematology \(incl blood transfusion\)](#) (22 articles)

[Infectious diseases](#) (204 articles)

[Nutrition and metabolism](#) (98 articles)

[Pharmacology and therapeutics](#) (193 articles)

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>