

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Murphy, GA; Asiki, G; Young, EH; Seeley, J; Nsubuga, RN; Sandhu, MS; Kamali, A; (2013) Cardiometabolic risk in a rural Ugandan population. *Diabetes care*, 36 (9). e143. ISSN 0149-5992 DOI: <https://doi.org/10.2337/dc13-0739>

Downloaded from: <http://researchonline.lshtm.ac.uk/1701186/>

DOI: <https://doi.org/10.2337/dc13-0739>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

OBSERVATIONS

Cardiometabolic Risk in a Rural Ugandan Population

Although many studies have examined metabolic syndrome (MetS) and the Framingham Risk Score (FRS), few studies have been carried out in African populations. This limited information on MetS and FRS leaves us with an incomplete understanding of the prevalence and distribution of risk of cardiometabolic disease in sub-Saharan Africa (SSA). It also prevents us from critically evaluating how each of the varying definitions of MetS compares in African populations. A clearer understanding of MetS and FRS in African populations may provide the basis for better identifying the impact of these definitions and tools on disease risk and, furthermore, help to evaluate the usefulness of such tools for research and for informing public health care and prevention policy in SSA.

In this study, we examined the prevalence and distribution of MetS and high (>20%) FRS and compare the World Health Organization (WHO), Adult Treatment Panel (ATP) III, International Diabetes Federation (IDF), and the newly proposed harmonized definitions of MetS in a rural Ugandan population (1). A total of 8,087 participants, aged 13 years and older, were surveyed, of whom 7,423 (55% women) had complete data for analysis. Data were collected using standard procedures, and prediabetes was defined using $HbA_{1c} \geq 5.7\%$ (≥ 39 mmol/mol) (2).

The prevalence of MetS varied by definition used, with the WHO, ATP III, IDF, and harmonized definitions resulting in MetS prevalence of 4.1%, 9.9%, 8.9%, and 13.7%, respectively. The harmonized definition was the most sensitive, capturing all those identified using ATP III and IDF and 85.4% of those identified using the WHO criteria. MetS increased with age (P value < 0.001), with a distinctive peak in the prevalence at ages 50–59 years for men for all definitions.

The age-standardized prevalence of MetS, for all definitions, was higher in women (5.0% [95% CI 4.3–5.6] to 18.6%

[95% CI 17.5–19.7]) than men (1.1% [95% CI 0.7–1.5] to 7.0% [95% CI 6.1–7.9]). The largest difference in MetS prevalence between men and women (1.1% [95% CI 0.7–1.5] vs. 14.5% [95% CI 13.5–15.6]) was found for the IDF definition. This was likely due to the substantial sex difference in central obesity (1.6% [95% CI 1.2–2.0] in men versus 29.7% [95% CI 28.4–31.0] in women), which is required in the IDF definition of MetS. Since there is no validated waist circumference cutoff for Africans, the IDF definition may currently be inappropriate for African populations (3).

The mean value of FRS was 3.30 (SD 6.5), with 3.5% having high FRS. Only 8–28%, depending on the MetS definition, had both MetS and high FRS. By contrast to MetS, high FRS was more common among men (5.8% [95% CI 5.1–6.5]) than women (1.9% [95% CI 1.5–2.3]). High FRS increased with age (P value < 0.001).

We found marked differences in the prevalence and distribution of cardiometabolic disease risk according to FRS and MetS definitions in this rural Ugandan population. These inconsistencies emphasize the need to more reliably assess the impact of these risk classifications in SSA populations. Prospective observational studies will be essential to evaluate and assess the distribution and determinants of cardiometabolic disease risk and to help to inform policy and health care programs in SSA (4).

GEORGINA A.V. MURPHY, MPHIL^{1,2}
 GERSHIM ASIKI, MD, MSC³
 ELIZABETH H. YOUNG, MBBS, PHD^{1,2}
 JANET SEELEY, PHD^{3,4,5}
 REBECCA N. NSUBUGA, PHD³
 MANJINDER S. SANDHU, PHD^{1,2}
 ANATOLI KAMALI, MD, PHD^{3,4}

From the ¹Department of Public Health and Primary Care, University of Cambridge, Cambridge, U.K.; the ²Wellcome Trust Sanger Institute, Hinxton, U.K.; the ³Medical Research Council/Uganda Virus Research Institute, Uganda Research Unit on AIDS, Entebbe, Uganda; the ⁴London School of Hygiene and Tropical Medicine, London, U.K.; and the ⁵School of International Development, University of East Anglia, Norwich, U.K.

Corresponding author: Georgina A.V. Murphy, gm7@sanger.ac.uk.
 DOI: 10.2337/dc13-0739

M.S.S. and A.K. contributed equally to this study.
 © 2013 by the American Diabetes Association.

Readers may use this article as long as the work is

properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—This work was sponsored by Medical Research Council (MRC), U.K. (grants G0801566 and G0901213-92157), awarded to M.S.S. and core funding to MRC/Uganda Virus Research Institute. G.A.V.M. was supported by the Gates Cambridge Scholarship.

No potential conflicts of interest relevant to this article were reported.

G.A.V.M. researched data, wrote the manuscript, and led data collection and management, and reviewed the manuscript. G.A. led data collection and management and reviewed the manuscript. E.H.Y. and J.S. researched data, wrote the manuscript, developed hypotheses, developed the study design, and reviewed the manuscript. R.N.N. led data collection and management and reviewed the manuscript. M.S.S. researched data, wrote the manuscript, developed hypotheses, developed the study design, and reviewed the manuscript. A.K. developed hypotheses, developed the study design, and reviewed the manuscript. M.S.S. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

The authors thank the GPC team and all other MRC staff who contributed to this study.

References

- Alberti KG, Eckel RH, Grundy SM, et al.; International Diabetes Federation Task Force on Epidemiology and Prevention; Hational Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; International Association for the Study of Obesity. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009;120:1640–1645
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2010;33(Suppl. 1):S62–S69
- Crowther NJ, Norris SA. The current waist circumference cut point used for the diagnosis of metabolic syndrome in sub-Saharan African women is not appropriate. *PLoS One* 2012;7:e48883
- Holmes MD, Dalal S, Volmink J, et al. Non-communicable diseases in sub-Saharan Africa: the case for cohort studies. *PLoS Med* 2010;7:e1000244