



Review Article

Pregnancy and helminth infections

H. MPAIRWE,¹ R. TWEYONGYERE² & A. ELLIOTT^{1,3}¹MRC/UVRI Uganda Research Unit on AIDS, Entebbe, Uganda, ²Makerere University College of Veterinary Medicine, Animal Resources and Biosecurity, Kampala, Uganda, ³London School of Hygiene & Tropical Medicine, London, UK

SUMMARY

It has been proposed that helminth infection may be particularly detrimental during pregnancy, through adverse effects on maternal anaemia and on birth outcomes, and that anthelmintic treatment during pregnancy will therefore be particularly beneficial. However, the few treatment trials that have been conducted have given, but little support to this notion and further trials in settings of nutritional stress are needed. It has also been proposed that prenatal exposure to helminth infection has an important effect on the development of the foetal immune response. There is evidence that this may impact, long-term, upon responses to helminth and nonhelminth antigens, and to allergens. Exposure to helminths in utero may also have nonspecific effects that may modify the offspring's susceptibility to diseases mediated by inflammation, including metabolic disorders. The mechanisms of such effects are not known, but they deserve to be explored as current epidemiological findings suggest the possibility of primary prevention for inflammatory conditions such as allergy, through intervention during pregnancy.

Keywords allergy, anaemia, birthweight, helminth, offspring, pregnancy

INTRODUCTION

Helminth infections are highly prevalent. In 2006, it was estimated that over a billion people were infected with one or more species of soil-transmitted helminth (1), while 200 million were infected with schistosomes (2). The prevalence of hookworm tends to increase with age (3) and, although the intensity of schistosomiasis declines with age,

the prevalence remains high among young adults in endemic communities (4). Thus, in endemic settings, there is a high burden of helminth infection among pregnant women, a group already at increased risk from malaria, and susceptible to anaemia and to nutritional stress (5–8). Implications for the mother and foetus include possible physiological effects on haemoglobin levels and on foetal growth and development, and possible immunological effects on both mother and foetus. This review will examine current evidence for such effects, focusing mainly on the latter, and discussing possible mechanisms for some of the immunological effects that have been observed.

Observational studies on the effects of helminth infections in humans are fraught with strong potential for both measured and unmeasured confounding, as helminths occur in the most disadvantaged communities alongside poverty, lack of education and malnutrition. Helminth-infected pregnant women are likely to be younger, less educated and of lower socio-economic status than uninfected women (5) – all factors that may impact upon nutrition and upon health-seeking behaviour (importantly, uptake of iron and folate supplements, preventive treatment for malaria and use of bed nets) for themselves and their infants. They may come from different ethnic backgrounds and live in different environments, compared with uninfected women (5), and these factors may influence customs or behaviour, and exposure of themselves and their infants to helminths and other infections. Co-exposure of the foetus or infant to malaria may be more common for environmental reasons, or because of immunomodulating effects of helminths (9, 10). Randomized clinical trials of anthelmintic treatment should help to address this and to identify reversible effects directly caused by helminths.

ANTHELMINTIC TREATMENT DURING PREGNANCY

Historically, pregnant and breastfeeding women have often been excluded from the treatment for helminths, due to

Correspondence: Harriet Mpairwe, MRC/UVRI Uganda Research Unit on AIDS, P.O. Box 49, Entebbe, Uganda (e-mails: mpairwe.harriet@gmail.com; mpairweus@yahoo.com).

Disclosures: None of the authors has any conflict of interest.

Received: 6 August 2013

Accepted for publication: 22 January 2014

concern about possible adverse effects of the drugs on the developing foetus. However, in 1994, the World Health Organization (WHO) recommended the treatment of pregnant women for hookworm, because of concern about hookworm-induced anaemia (11). Further, in 2002, a WHO informal consultation recommended the use of praziquantel during pregnancy and lactation, and that in schistosomiasis-endemic settings, all girls and women of child-bearing age should be included in praziquantel mass drug administration programmes (12). It was hoped that implementation of these recommendations would result in benefits including reduced maternal anaemia, increased birthweight and reduced perinatal and infant mortality. Neither of these recommendations was based on evidence of benefit or safety from controlled trials, and in the case of praziquantel, a subsequent WHO Scientific Working Group, held in 2006, called for such trials to be conducted (13). However, only a very small number of anthelmintic trials have yet been conducted among pregnant women (14–22) (Table 1), and taken together, the results are, as yet, inconclusive.

PHYSIOLOGICAL EFFECTS OF HELMINTHS IN PREGNANCY

There is substantial evidence from studies in human subjects that certain species of helminth infection during pregnancy are associated with maternal anaemia, as reviewed by Brooker and colleagues in 2008 (23), and observed in several subsequent studies (24–27). It is likely that heavy intensities of hookworm and *Trichuris trichiura* are particularly important (28). However, only two of five trials assessing this outcome have shown a benefit of treatment for soil-transmitted helminths during pregnancy (Table 1) (17, 18). In a trial setting, it is almost imperative to provide adequate iron and folate supplementation, and (where relevant) presumptive treatment for malaria during pregnancy, as standard of care during pregnancy. This may override some of the potential benefits of anthelmintics: Torlesse and colleagues found greater benefit of haematinic supplementation than of albendazole in a hookworm-endemic region of Sierra Leone (18).

Studies in mice suggest that maternal worm infections influence foetal linear growth and the growth of lymphoid tissues and bone (29, 30). There are also observational studies in human subjects suggesting that certain maternal helminth infections adversely affect birthweight and perinatal and infant mortality and that anthelmintics during pregnancy may be beneficial for these outcomes (31–35) [reviewed by Freidman (35) and Imhoff-Kunsch (36)], but trials to date have failed to confirm that these associations are causal by showing a benefit of treatment (Table 1). Anaemia, itself, is an important cause of adverse birth

outcomes (37–39), and if effects of helminths on anaemia are obscured by supplementation with haematinics this may also obscure possible benefits of anthelmintics mediated by reductions in anaemia (34).

If the hypothesis that anthelmintic treatment during pregnancy can have important benefits for maternal and infant birth outcomes is to be rigorously tested, and the policy of routine anthelmintic treatment during pregnancy is to be validated, large trials are still needed in a range of settings. It may be that anthelmintics will be found most useful in settings in which anaemia is prevalent, and where there is nutritional stress and limited access to haematinic supplementation. In such settings, single dose benzimidazoles, [which are easy to administer and have been highly effective against hookworm in most trials to date (15, 16, 18, 19)] may have benefits for birth outcomes. No trials have yet demonstrated benefits of praziquantel treatment during pregnancy for maternal anaemia or for birth outcomes, but a further placebo-controlled trial of praziquantel treatment for *Schistosoma japonicum* during pregnancy is in progress and scheduled for completion this year (40). This will be of particular interest because *S. japonicum* has been shown to be associated with inflammation in maternal, placental and foetal tissues (41), and studies *in vitro* suggest that exposure of trophoblast cells to *S. japonicum* egg antigen may result in impaired placentation (42).

EFFECTS OF PREGNANCY ON SUSCEPTIBILITY TO HELMINTHS AND ON THE RESPONSE TO THEIR TREATMENT

Pregnancy results in major hormonal and nutritional changes, and in immunological changes allowing successful allograft retention (43–45). In mammals other than humans, there is a well-recognized increase in susceptibility to helminths, or at least to helminth egg production, around parturition. Immune modulation resulting from hormonal changes (such as a reduction in circulating cortisol) (46), or nutritional changes, such as protein deficiency (47) may contribute to this effect. In humans, data are conflicting, with studies showing increased (7) or similar (48) helminth egg counts between pregnant and nonpregnant women.

Studies in mice also suggest that pregnancy is associated with increased schistosome-induced pathology, related to reduced parasite-specific interferon (IFN)- γ but increased interleukin (IL)-4 production (49). If similar effects were demonstrated in humans, this would be a further strong argument for emphasizing schistosomiasis treatment in young, and pregnant, women. In the Entebbe Mother and Baby Study (EMaBS), a trial of albendazole and praziquantel

Table 1 Clinical trials of anthelmintic treatment during pregnancy

Country; trial name	Participants	Helminth prevalence	Intervention	Outcomes assessed	Main results	References
Entebbe, Uganda; Entebbe mother and baby study	2507	Hookworm <i>Mansonella perstans</i> <i>Schistosoma mansoni</i> <i>Schistosoma stercoralis</i> <i>Trichuris trichiura</i> <i>Ascaris lumbricoides</i>	Alb 400 mg vs. placebo pzq 40 mg/kg vs. placebo each single dose	Maternal anaemia Maternal HIV load MTCT of HIV Congenital abnormalities Adverse birth outcomes ^a Birthweight Infant mortality Infant response to vaccines Infant malaria, diarrhoea, pneumonia Infantile eczema	No effects Reduced by alb No effects No effects No effects No effects No effects No effects Increased by alb, pzq No effects Benefit	(19–22)
Freetown and Port Loko, Sierra Leone	184	Hookworm ^b <i>Trichuris trichiura</i> <i>Ascaris lumbricoides</i>	Alb 400 mg vs. calcium and vitamin D single dose	Infant cognitive development Change in anaemia btw 1st & 3rd trimester Change in iron status btw 1st & 3rd trimester Miscarriages	No effect Benefit No effect	(18)
Iquitos, Peru	1042	Hookworm <i>Trichuris trichiura</i> <i>Ascaris lumbricoides</i>	Meb 500 mg single dose	Congenital abnormalities Maternal anaemia Birthweight Low birthweight Congenital abnormalities Adverse birth outcomes ^a Helminth cure	No effect No effect No effect Reduced No effect No effect Alb superior for hkwm	(15)
Masindi, Uganda	591 STH-infected 241 comparison ^c	Hookworm <i>Trichuris trichiura</i> <i>Ascaris lumbricoides</i> <i>Schistosoma mansoni</i>	Alb400 mg vs. ivc (dose according to height) vs. ivc plus alb	Miscarriage, prematurity Congenital abnormalities Maternal haemoglobin Birthweight Infant haemoglobin Infant mortality Maternal anaemia (<10.5 g/dL) at term Maternal haemoglobin (Hb) at term Maternal anaemia 4 months post-partum Maternal Hb 4 months post-partum Helminth cure	No effect No effect Benefit	(16)
Rufiji, Tanzania	3080 cluster randomized by antenatal clinic		Albendazole			(17)
Lamberene, Gabon	65	<i>Schistosoma</i> <i>haematobium</i>	Mefloquine 15 mg/kg vs. SP (3 tablets of 500/25 mg) twice during pregnancy	Helminth cure	Mefloquine superior	(14)

^aMiscarriages, still births, neonatal deaths and (in Gyorkos 2006) prematurity; ^bNecator americanus; ^conly infected women were included in the treatment trial; alb, albendazole; pzq, praziquantel; meb, mebendazole; ivc, ivermectin; STH, soil-transmitted helminths; btw, between primary outcomes highlighted.

among pregnant women in Uganda (50), we examined effects of pregnancy on the immune response to schistosomiasis by comparing responses during pregnancy and after delivery in placebo recipients. Interleukin-2, IL-4 and IL-10 responses to schistosome worm and egg antigens were similar during pregnancy and after delivery, whereas IFN- γ , IL-5 and IL-13 responses increased after delivery, suggesting that pregnancy in humans is indeed associated with a modified profile of antischistosome cellular responses (51). Antischistosome antibody levels were also lower during pregnancy (52). Further studies to investigate whether these immunological differences are associated with differences in the progression of pathology would be justified.

Treatment of schistosomiasis with praziquantel in non-pregnant human subjects is associated with release of worm antigen into the blood stream and with a boost in cellular and antibody responses to worm and egg antigen (53). We observed a similar, but smaller boost in responses following treatment during pregnancy, when compared to treatment after delivery (51, 52). This may be important for several reasons. First, the immune response is thought to synergize with the direct effects of praziquantel, helping to kill the parasites (54). A reduced boost might mean that treatment in pregnancy was less effective. However, we found no difference in egg reduction following treatment during pregnancy compared with treatment after delivery in our study (51). Second, the sudden immunological changes following treatment might have implications for the success of the pregnancy. Successful pregnancies are associated with a placental microenvironment that is dominated by a type two cytokine milieu, characterized by increased production of IL-4 and IL-10, and decreased production of IFN γ and IL-2 (55). Animal models of pregnancy have shown that increased systemic and placental IFN γ and TNF α levels are associated with poor pregnancy outcomes (56, 57). Reassuringly, in EMaBS, praziquantel treatment of *Schistosoma mansoni* during pregnancy was not associated with an increased rate of miscarriage or still birth (19). However, the number of women with schistosomiasis in the study was fairly small (458), so its power to detect differences in these rates was limited. Vigilance regarding birth outcomes will be important as the 2002 WHO recommendations (12), to use praziquantel during pregnancy and lactation, are more widely implemented.

IMMUNOLOGICAL EFFECTS OF HELMINTHS ON PREGNANT WOMEN

Helminth infections modulate the host immune response through a variety of mechanisms (58). These allow long-term survival of the parasite within the host, while minimizing

damage to host tissues. They also have implications that may be important during pregnancy.

First, they may impact upon susceptibility to heterologous infections. Effects on susceptibility to malaria might be especially relevant during pregnancy, given its adverse impact on pregnancy outcomes (59). Positive associations between hookworm and malaria have been reported (9, 60, 61), while *S. haematobium* and *Ascaris lumbricoides* have been associated with lower malaria parasite densities (60) and lower malaria risk (61), respectively. While these associations might be confounded by environmental and behavioural factors, measured confounders failed to explain the positive hookworm malaria and *Mansonella* malaria associations observed in our study (9), and immunological mechanisms are also possible.

Second, they may impact upon maternal metabolism – for example glucose metabolism (62). A recent study in mice showed that classical activation of macrophages in adipose tissue (favouring inflammatory immune responses) was associated with glucose intolerance, while alternative activation was associated with enhanced glucose tolerance. Eosinophils in adipose tissue were shown to be a key source of IL-4 for the induction of alternatively activated macrophages, and eosinophils induced by helminth infection were shown to produce a sustained improvement in glucose tolerance (62). Similar effects in humans might be beneficial in relation to gestational diabetes.

IMMUNOLOGICAL EFFECTS OF HELMINTHS DURING PREGNANCY ON THE FOETUS

Arguably, the most important potential impact of worm infections during pregnancy is upon long-term outcomes in the offspring.

Effects on helminth-specific responses

There is considerable evidence from studies in both animal models (63–65) and humans (65–70) that in utero exposure to helminths can influence susceptibility to the same helminth species later in life. For example, in 1991, Lammie and colleagues, working in an area endemic for bancroftian filariasis, showed that maternal, but not paternal, filarial infection was associated with childhood infection and that children of infected mothers had reduced cellular responses to microfilariae compared with children of uninfected mothers (66). Other studies supported this result (67). Subsequently, it was shown that children of infected mothers who were sensitized *in utero* to filarial antigen showed similar immunological and clinical outcomes to unexposed infants, but that children who were exposed

in utero and not sensitized (suggesting the acquisition of tolerance) were slower to develop detectable cellular responses and more likely to become microfilaraemic than either sensitized or unexposed infants (71, 72). These results suggest an immunological explanation for the epidemiological observations. There is evidence for similar effects of intrauterine exposure in humans for onchocerciasis (69), schistosomiasis (70) and soil-transmitted helminth infection (68). While increasing susceptibility, the tolerogenic effects of prenatal exposure may reduce inflammation-induced pathology, resulting in an improved outcome for both helminth and host.

Effects on responses to heterologous vaccines and infections

Observational studies have also suggested that exposure to worm infections during pregnancy can alter the offspring's response to unrelated antigens, including vaccines, infectious diseases and allergens. A study in Kenya found that infants exposed and sensitized to schistosomiasis and filarial worms *in utero* showed a Th2 bias in their immune responses to BCG vaccine, while infants who were exposed but not sensitized showed a Th1 profile of response (73). Such an effect might contribute to explaining the poor efficacy of BCG immunization in tropical developing countries – one of the settings in which protection against tuberculosis is most needed (74, 75). In the EMaBS cohort, in which we compared responses following BCG and tetanus immunization in infants of helminth-infected and helminth-uninfected women, we did not observe effects on the infant Th1 or Th2 response to mycobacterial antigens, but treatment with albendazole reduced Th2 responses to tetanus toxoid among the infants of mothers with hookworm (22). Why was this effect seen for the response to tetanus immunization, but not for the response to BCG? One possibility is that concurrent prenatal exposure to helminths and to the heterologous antigen of interest may be important for the helminth effects. EMaBS women received tetanus immunization during pregnancy, providing a potential opportunity for exposure of the foetus to this antigen (76); indeed the infant cytokine response to tetanus immunization increased with the number of doses that the mother received during pregnancy, suggesting that prenatal priming of the infant anti-tetanus response had occurred (77). We also found that infants of mothers with *M. perstans* infection had elevated IL-10 responses to both BCG-related and tetanus antigens, but these associations were seen only among infants of mothers who received no albendazole, suggesting that prenatal exposure to albendazole, or to its effects on maternal infections, blocked the ability of maternal *M. perstans* to

promote regulatory responses to heterologous antigens in the infants (77).

Besides tuberculosis, HIV infection and malaria are among the most important co-infections in helminth-endemic settings, and there is considerable controversy as to the effects of helminths on susceptibility to both (78). Prenatal exposure to maternal helminths has been postulated to increase the risk of vertical HIV transmission by activation of foetal lymphocytes, increasing their susceptibility to HIV infection. A small observational study in Kenya (among 44 HIV-exposed infants, 13 of whom were HIV infected) supported this hypothesis with an odds ratio, adjusted for maternal malaria [aOR (95% confidence interval)] of 7.3 (2.4–33.7) for any maternal helminth vs. none (79). In fact, the association was prominent among infants of mothers with filariasis, but not for other helminth species. In EMaBS, we studied the association between vertical HIV transmission and maternal helminth infection among 203 HIV-exposed babies (34 HIV-infected), and there was no statistically significant association with helminth infection (aOR for all helminths combined, adjusted for the use of nevirapine for prevention of mother-to-child HIV transmission and for vaginal delivery vs. Caesarian section, 2.23 (0.86–5.79); K. Jacqueline, M.E. Alison, unpublished data). However, anthelmintic treatment during pregnancy showed no benefit for this outcome (22); thus, no firm conclusion can yet be reached as to the effects of maternal helminths on vertical HIV transmission.

In EMaBS, we also investigated associations between maternal helminth and maternal malaria infection and infant susceptibility to malaria. We found that maternal worm infections, including hookworm, *M. perstans* and *S. mansoni* were associated with increased risk of malaria incidence in the offspring (10). These associations persisted after adjusting for potential confounders (maternal age, education, parity, net ownership, socio-economic status, maternal malaria and human immunodeficiency virus infections, and location of residence) and after stratifying for maternal malaria infection status. In fact, there was a suggestion that the effect was stronger among infants of mothers who had malaria during pregnancy, again suggesting that concurrent exposure to maternal malaria and to maternal helminths might be important for helminth-induced effects on the offspring's response to malaria, as discussed above in relation to tetanus immunization.

Effects on allergy-related disease outcomes

The most striking findings of EMaBS was that maternal albendazole during pregnancy was associated with an increased incidence of infantile eczema (20) and of eczema in the first 5 years of life (21). This effect was not confined

to the infants of mothers with susceptible worm infections (hookworm, *Ascaris*, *Trichuris* and *Strongyloides*), so the mechanism of the effect of albendazole was not clear. However, maternal hookworm showed a significant inverse association with eczema in childhood, which increased with intensity (80), and maternal hookworm modified the effect of classical risk factors for childhood eczema, including maternal history of eczema and childhood atopy (H. Mpairwe, J. Ndibazza, E.L. Webb, M. Nampijja, L. Muhangi, B. Apule, S. Lule, H. Akurut, D. Kizito, M. Kakande, F.M. Jones, C.M. Fitzsimmons, M. Muwanga, L.C. Rodrigues, D.W. Dunne, A.M. Elliott, unpublished data). Maternal treatment with praziquantel was associated with an increased incidence of infantile eczema among infants whose mothers were infected with *S. mansoni* during pregnancy (20). Together, these findings strongly suggest that prenatal exposure to maternal helminth infection protects against infantile eczema and against the adverse effects of other exposures on this outcome. Whether this effect persists into later childhood, and influences risk of asthma, is currently under investigation.

MECHANISMS BY WHICH MATERNAL HELMINTH INFECTION MIGHT INFLUENCE OUTCOMES IN THE FOETUS

The mechanisms by which maternal helminth infection can influence outcomes in the foetus are not known. Investigation of these mechanisms might shed important light on the development on the infant immune response and inform the development of possible interventions for

the primary prevention of immunologically mediated conditions, such as allergy (81). Some possible mechanisms are summarized in Figure 1 and discussed below.

Foetal exposure and sensitization to foreign antigens

It is clear that the human foetus can become sensitized to parasite (73, 82–88) and vaccine antigens (76) *in utero*, and at first sight, this is surprising as most macromolecules and immune complexes in the maternal circulation are broken down by macrophages in the trophoblast, providing amino acids for the growth of the placenta and foetus and preventing transfer to the foetal circulation (89). However, filarial (87) and schistosome (90–92) antigens have been detected in cord blood in humans. Similarly, there is evidence that tetanus toxoid (93) and malaria antigen (94) can cross the placenta. Studies in mice also suggest that mycobacterial antigen can cross the placenta (95), and although there are differences in placentation between mice and humans, there is evidence of prenatal sensitization of the human foetus to mycobacterial antigens *in utero* (73). How such antigens cross the placenta is not known. One possibility is that the antigens are transported bound to IgG antibody, protected by the foetal immunoglobulin G receptor (FcRn): this is supported by studies showing that the transfer of tetanus toxoid, or of malaria antigen, is dependent upon the presence or concentration of specific antibody in maternal blood (94), but the precise process of transfer is not fully understood.

With regard to helminth-specific immune responses, epidemiological evidence discussed above suggests that *in utero*




Figure 1 Possible mechanisms by which maternal helminth infections might influence foetal immunological development.

sensitization results in down-regulated responses among the offspring, on encountering the homologous antigen. As reviewed by Dauby and colleagues (96), this may be due to a bias in the foetal and neonatal immune response towards the development T-regulatory and T-helper (Th)17 responses, rather than Th1 responses. The effects of helminth exposure might be analogous to findings for effects of placental malaria: *in vitro* depletion of CD4 + CD25hi regulatory T cells in cord blood demonstrated their contribution to suppression of neonatal responses to malaria antigens in malaria-exposed infants (97).

Transfer of antigen-specific idiotype antibodies across the placenta

In addition to the possibility that antibodies facilitate the transfer of antigen across the placenta, there is evidence that maternal antibody idiotypes specific for helminth antigens can be transferred across the placenta and can themselves induce sensitization of the foetus in both animal models (64) and humans (98). In the case of schistosomiasis, studies in mice suggest that, by promoting IFN- γ responses in the neonate, *in utero* sensitization down-regulates the (Th2-mediated) granuloma formation associated with pathology (64).

Non-helminth-specific effects

These possible mechanisms largely provide explanations for effects of maternal helminth infections on the offspring's response to the same helminth infection. How might maternal helminth infection influence the foetal response to unrelated antigens?

Cross-reactivity between helminth immunogens and allergens

In the case of allergens, one possibility may be cross-reactivity between helminth antigens and allergens. There are many structural similarities between allergens and metazoan parasite immunogens (99), and IgE cross-reactivity has been demonstrated in several cases – for example, in human subjects, IgE cross-reactivity has been described between proteins of the nematodes *Anisakis Simplex* (100) and *A. lumbricoides* (101) and the allergen *Dermatophagoides pteronyssinus*; IgE, IgG and IgG4 cross-reactivity has been demonstrated between filarial antigen and cockroach allergen (102). IgE cross-reactivity has also been demonstrated between peanut allergen and schistosome egg antigen in a setting in Ghana where almost no clinical peanut allergy occurs and this has been shown to be attributable to cross-reactive carbohydrate determinants, rather than

protein antigens (103). We hypothesize that intrauterine exposure to cross-reactive parasite immunogens might result in foetal acquisition of tolerance to allergens, or foetal sensitization to related, parasite-specific IgG idiotypes might modulate the foetal allergen-specific response to a nonpathogenic profile.

Effects on innate immune responses

Alternatively, maternal helminth infection might influence the foetal innate immune system in such a way as to modulate the response induced on co-exposure to a wide range of heterologous antigens including malaria, tetanus and mycobacterial antigen, as well as allergens. Allergologists have explored this possibility to investigate how prenatal exposure to a barn environment might influence the offspring's risk of allergy-related disease. Using a mouse model of experimental allergic airway inflammation, Conrad and colleagues showed that low-grade inflammation in the maternal lung induced by the cowshed-derived bacterium *Acinetobacter lwoffii* was associated with reduced susceptibility to experimentally induced asthma in the offspring and that this was mediated by up-regulation of maternal Toll-like receptor (TLR) expression and circulating pro-inflammatory cytokines, and down-regulation of placental TLR expression (104, 105). Helminths also have systemic effects on circulating inflammatory mediators and TLR expression and might influence placental and foetal antigen-presenting cell profiles in a related manner (106).

CONCLUSION

Helminth infection during pregnancy is common, but the benefits of anthelmintic treatment during pregnancy are not yet clear either for the mother or for the offspring. On the other hand, there is strong epidemiological evidence that prenatal exposure to maternal helminth infection tends to down-modulate the offspring's response following infection, resulting in increased susceptibility to the same helminth species, but reduced pathology. Thus, an unwanted consequence of partially successful intervention against helminths may be increased susceptibility to helminth-induced pathology in some endemic settings as prevalence during pregnancy declines. Of more interest, however, is the increasing evidence that prenatal exposure to helminths may influence the offspring's susceptibility to other conditions, and particularly to allergy-related disease. The mechanisms of this effect deserve urgent investigation as they may indicate possible routes to the primary prevention, or management, of these conditions.

ACKNOWLEDGEMENTS

We thank the Wellcome Trust, which has funded the Entebbe Mother and Baby Study (Grant Numbers 064693, 079110, 95778) with additional funding from the European Community's Seventh Framework Programme (FP7/2007-2013) under EC-GA number 241642. Harriet Mpairwe is currently a Wellcome Trust Uganda Post-doctoral Fellow

in Infection and Immunity, funded by a Wellcome Trust Strategic Award, Grant Number 084344. Robert Tweyongyere is a European Foundations Initiative for Neglected Tropical Diseases Senior Fellow, Grant Number Ref I/84 020, Ref 86 529. We thank Dr Stephen Cose for reviewing the manuscript.

REFERENCES

- Bethony J, Brooker S, Albonico M, *et al*. Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *Lancet* 2006; **367**: 1521–1532.
- Steinmann P, Keiser J, Bos R, Tanner M & Utzinger J. Schistosomiasis and water resources development: systematic review, meta-analysis, and estimates of people at risk. *Lancet Infect Dis* 2006; **6**: 411–425.
- Pullan RL, Kabatereine NB, Quinell RJ & Brooker S. Spatial and genetic epidemiology of hookworm in a rural community in Uganda. *PLoS Negl Trop Dis* 2010; **4**: e713.
- Fitzsimmons CM, Jones FM, Pinot de Moira A, *et al*. Progressive cross-reactivity in IgE responses: an explanation for the slow development of human immunity to schistosomiasis? *Infect Immun* 2012; **80**: 4264–4270.
- Woodburn PW, Muhangi L, Hillier S, *et al*. Risk Factors for helminth, malaria, and HIV infection in pregnancy in Entebbe, Uganda. *PLoS Negl Trop Dis* 2009; **3**: e473.
- van Eijk AM, Lindblade KA, Odhiambo F, *et al*. Geohelminth infections among pregnant women in rural western Kenya: a cross-sectional study. *PLoS Negl Trop Dis* 2009; **3**: e370.
- Adegnik AA, Ramharter M, Agnandji ST, *et al*. Epidemiology of parasitic co-infections during pregnancy in Lambarene, Gabon. *Tropical Med Int Health* 2010; **15**: 1204–1209.
- Belyhun Y, Medhin G, Amberbir A, *et al*. Prevalence and risk factors for soil-transmitted helminth infection in mothers and their infants in Butajira, Ethiopia: a population based study. *BMC Public Health* 2010; **10**: 21.
- Hillier SD, Booth M, Muhangi L, *et al*. *Plasmodium falciparum* and helminth coinfection in a semi urban population of pregnant women in Uganda. *J Infect Dis* 2008; **198**: 920–927.
- Ndibazza J, Webb EL, Lule S, *et al*. Associations between maternal helminth and malaria infections in pregnancy and clinical malaria in the offspring: a birth cohort in Entebbe, Uganda. *J Infect Dis* 2013; **208**: 2007–2016.
- WHO. Report of the WHO informal consultation on hookworm infection and anaemia in girls and women. Geneva, 5–7 December 1994. WHO/CTD/SIP/96.1. 1995.
- WHO. Report of the WHO informal consultation on the use of praziquantel during pregnancy/lactation and albendazole/mebendazole in children under 24 months. Geneva, 8–9 April 2002. WHO/CDS/CEP/PVC/2002.4. 2002.
- WHO. Report of the Scientific Working Group meeting on Schistosomiasis. Geneva, 14–16 November, 2005. TDR/SWG/07. 2006.
- Basra A, Mombo-Ngoma G, Melser MC, *et al*. Efficacy of mefloquine intermittent preventive treatment in pregnancy against *Schistosoma haematobium* infection in Gabon: a nested randomized controlled assessor-blinded clinical trial. *Clin Infect Dis* 2013; **56**: e68–e75.
- Larocque R, Casapia M, Gotuzzo E, *et al*. A double-blind randomized controlled trial of antenatal mebendazole to reduce low birthweight in a hookworm-endemic area of Peru. *Tropical Med Int Health* 2006; **11**: 1485–1495.
- Ndyomugenyi R, Kabatereine N, Olsen A & Magnussen P. Efficacy of ivermectin and albendazole alone and in combination for treatment of soil-transmitted helminths in pregnancy and adverse events: a randomized open label controlled intervention trial in Masindi district, western Uganda. *Am J Trop Med Hyg* 2008; **79**: 856–863.
- Urassa DP, Nystrom L & Carlsted A. Effectiveness of routine antihelminthic treatment on anaemia in pregnancy in Rufiji District, Tanzania: a cluster randomised controlled trial. *East Afr J Public Health* 2011; **8**: 176–184.
- Torlesse H & Hodges M. Albendazole therapy and reduced decline in haemoglobin concentration during pregnancy (Sierra Leone). *Trans R Soc Trop Med Hyg* 2001; **95**: 195–201.
- Ndibazza J, Muhangi L, Akishule D, *et al*. Effects of deworming during pregnancy on maternal and perinatal outcomes in Entebbe, Uganda: a randomized controlled trial. *Clin Infect Dis* 2010; **50**: 531–540.
- Mpairwe H, Webb EL, Muhangi L, *et al*. Anthelmintic treatment during pregnancy is associated with increased risk of infantile eczema: randomised-controlled trial results. *Pediatr Allergy Immunol* 2011; **22**: 305–312.
- Ndibazza J, Mpairwe H, Webb EL, *et al*. Impact of anthelmintic treatment in pregnancy and childhood on immunisations, infections and eczema in childhood: a randomised controlled trial. *PLoS ONE* 2012; **7**: e50325.
- Webb EL, Mawa PA, Ndibazza J, *et al*. Effect of single-dose anthelmintic treatment during pregnancy on an infants response to immunisation and on susceptibility to infectious diseases in infancy: a randomised, double-blind, placebo-controlled trial. *Lancet* 2011; **377**: 52–62.
- Brooker S, Hotez PJ & Bundy DA. Hookworm-related anaemia among pregnant women: a systematic review. *PLoS Negl Trop Dis* 2008; **2**: e291.
- Yatch NJ, Jolly PE, Funkhouser E, *et al*. The effect of malaria and intestinal helminth coinfection on birth outcomes in Kumasi, Ghana. *Am J Trop Med Hyg* 2010; **82**: 28–34.
- Boel M, Carrara VI, Rijken M, *et al*. Complex Interactions between soil-transmitted helminths and malaria in pregnant women on the Thai-Burmese border. *PLoS Negl Trop Dis* 2010; **4**: e887.
- Finkelstein JL, Mehta S, Duggan CP, *et al*. Predictors of anaemia and iron deficiency in HIV-infected pregnant women in Tanzania: a potential role for vitamin D and parasitic infections. *Pub Health Nutr* 2012; **15**: 928–937.
- Makhoul Z, Taren D, Duncan B, *et al*. Risk factors associated with anemia, iron deficiency and iron deficiency anemia in rural Nepali pregnant women. *Southeast Asian J Trop Med Public Health* 2012; **43**: 735–746.
- Gyorkos TW, Gilbert NL, Larocque R, Casapia M & Montresor A. Re-visiting *Trichuris trichiura* intensity thresholds based on anemia during pregnancy. *PLoS Negl Trop Dis* 2012; **6**: e1783.
- Odiere MR, Koski KG, Weiler HA & Scott ME. Concurrent nematode infection and pregnancy induce physiological responses that impair linear growth in the murine foetus. *Parasitology* 2010; **137**: 991–1002.
- Odiere MR, Scott ME, Leroux LP, Dzierszinski FS & Koski KG. Maternal protein deficiency during a gastrointestinal nematode infection alters developmental profile

- of lymphocyte populations and selected cytokines in neonatal mice. *J Nutr* 2013; **143**: 100–107.
- 31 Atukorala TM, de Silva LD, Dechering WH, Dassenaieke TS & Perera RS. Evaluation of effectiveness of iron-folate supplementation and anthelmintic therapy against anemia in pregnancy—a study in the plantation sector of Sri Lanka. *Am J Clin Nutr* 1994; **60**: 286–292.
 - 32 de Silva NR, Sirisena JL, Gunasekera DP, Ismail MM & de Silva HJ. Effect of mebendazole therapy during pregnancy on birth outcome. *Lancet* 1999; **353**: 1145–1149.
 - 33 Christian P, Khatry SK & West KP Jr. Antenatal anthelmintic treatment, birth-weight, and infant survival in rural Nepal. *Lancet* 2004; **364**: 981–983.
 - 34 Passerini L, Casey GJ, Biggs BA, et al. Increased birth weight associated with regular pre-pregnancy deworming and weekly iron-folic acid supplementation for Vietnamese women. *PLoS Negl Trop Dis* 2012; **6**: e1608.
 - 35 Friedman JF, Mital P, Kanzaria HK, Olds GR & Kurtis JD. Schistosomiasis and pregnancy. *Trends Parasitol* 2007; **23**: 159–164.
 - 36 Imhoff-Kunsch B & Briggs V. Antihelminthics in pregnancy and maternal, newborn and child health. *Paediatr Perinat Epidemiol* 2012; **26**(Suppl 1): 223–238.
 - 37 Yatch NJ, Funkhouser E, Ehiri JE, et al. Malaria, intestinal helminths and other risk factors for stillbirth in Ghana. *Infect Dis Obstet Gynecol* 2010; **2010**: 350763.
 - 38 Bodeau-Livinec F, Briand V, Berger J, et al. Maternal anemia in Benin: prevalence, risk factors, and association with low birth weight. *Am J Trop Med Hyg* 2011; **85**: 414–420.
 - 39 Zhang Q, Ananth CV, Li Z & Smulian JC. Maternal anaemia and preterm birth: a prospective cohort study. *Int J Epidemiol* 2009; **38**: 1380–1389.
 - 40 ClinicalTrials.gov. *S. japonicum* and pregnancy.
 - 41 Kurtis JD, Higashi A, Wu HW, et al. Maternal *Schistosomiasis japonica* is associated with maternal, placental, and fetal inflammation. *Infect Immun* 2011; **79**: 1254–1261.
 - 42 McDonald EA, Friedman JF, Sharma S, et al. *Schistosoma japonicum* soluble egg antigens attenuate invasion in a first trimester human placental trophoblast model. *PLoS Negl Trop Dis* 2013; **7**: e2253.
 - 43 Robertson SA. Control of the immunological environment of the uterus. *Rev Reprod* 2000; **5**: 164–174.
 - 44 Guleria I & Sayegh MH. Maternal acceptance of the fetus: true human tolerance. *J Immunol* 2007; **178**: 3345–3351.
 - 45 Mellor AL & Munn DH. Extinguishing maternal immune responses during pregnancy: implications for immunosuppression. *Semin Immunol* 2001; **13**: 213–218.
 - 46 Beasley AM, Kahn LP & Windon RG. The periparturient relaxation of immunity in Merino ewes infected with *Trichostrongylus colubriformis*: endocrine and body compositional responses. *Vet Parasitol* 2010; **168**: 51–59.
 - 47 Sakkas P, Houdijk JG, Jones LA, Knox DP & Kyriazakis I. Dietary protein and energy supplies differentially affect resistance to parasites in lactating mammals. *Br J Nutr* 2011; **106**: 1207–1215.
 - 48 Herter U, Petney T, Pipitgool V, et al. The influence of pregnancy on intestinal parasite infection in Thai women. *Acta Trop* 2007; **101**: 200–206.
 - 49 Farah IO, Langoi D, Nyaundi J & Hau J. Schistosome-induced pathology is exacerbated and Th2 polarization is enhanced during pregnancy. *In Vivo* 2007; **21**: 599–602.
 - 50 Elliott AM, Kizza M, Quigley MA, et al. The impact of helminths on the response to immunization and on the incidence of infection and disease in childhood in Uganda: design of a randomized, double-blind, placebo-controlled, factorial trial of deworming interventions delivered in pregnancy and early childhood [ISRCTN32849447]. *Clinical Trials* (London, England) 2007; **4**: 42–57.
 - 51 Tweyongyere R, Mawa PA, Ngom-Wegi S, et al. Effect of praziquantel treatment during pregnancy on cytokine responses to schistosome antigens: results of a randomized, placebo-controlled trial. *J Infect Dis* 2008; **198**: 1870–1879.
 - 52 Tweyongyere R, Mawa PA, Emojong NO, et al. Effect of praziquantel treatment of *Schistosoma mansoni* during pregnancy on intensity of infection and antibody responses to schistosome antigens: results of a randomised, placebo-controlled trial. *BMC Infect Dis* 2009; **9**: 32.
 - 53 Joseph S, Jones FM, Walter K, et al. Increases in human T helper 2 cytokine responses to *Schistosoma mansoni* worm and worm-tegument antigens are induced by treatment with praziquantel. *J Infect Dis* 2004; **190**: 835–842.
 - 54 Fallon PG, Cooper RO, Probert AJ & Doenhoff MJ. Immune-dependent chemotherapy of schistosomiasis. *Parasitology* 1992; **105**(Suppl): S41–S48.
 - 55 Formby B. Immunologic response in pregnancy. Its role in endocrine disorders of pregnancy and influence on the course of maternal autoimmune diseases. *Endocrinol Metab Clin North Am* 1995; **24**: 187–205.
 - 56 Chaouat G, Menu E, Clark DA, Dy M, Minkowski M & Wegmann TG. Control of fetal survival in CBA x DBA/2 mice by lymphokine therapy. *J Reprod Fertil* 1990; **89**: 447–458.
 - 57 Vassiliadis S, Tsoukatos D & Athanassakis I. Interferon-induced class II expression at the spongiotrophoblastic zone of the murine placenta is linked to fetal rejection and developmental abnormalities. *Acta Physiol Scand* 1994; **151**: 485–495.
 - 58 van Riet E, Hartgers FC & Yazdanbakhsh M. Chronic helminth infections induce immunomodulation: consequences and mechanisms. *Immunobiology* 2007; **212**: 475–490.
 - 59 Shulman CE & Dorman EK. Importance and prevention of malaria in pregnancy. *Trans R Soc Trop Med Hyg* 2003; **97**: 30–35.
 - 60 Thigpen MC, Filler SJ, Kazembe PN, et al. Associations between peripheral *Plasmodium falciparum* malaria parasitemia, human immunodeficiency virus, and concurrent helminth infection among pregnant women in Malawi. *Am J Trop Med Hyg* 2011; **84**: 379–385.
 - 61 Boel M, Carrara VI, Rijken M, et al. Complex interactions between soil-transmitted helminths and malaria in pregnant women on the Thai-Burmese border. *PLoS Negl Trop Dis* 2010; **4**: e887.
 - 62 Wu D, Molofsky AB, Liang HE, et al. Eosinophils sustain adipose alternatively activated macrophages associated with glucose homeostasis. *Science* (New York, NY) 2011; **332**: 243–247.
 - 63 Othman AA, Shoheib ZS, Saied EM & Soliman RH. Congenital exposure to *Schistosoma mansoni* infection: impact on the future immune response and the disease outcome. *Immunobiology* 2010; **215**: 101–112.
 - 64 Montesano MA, Colley DG, Eloi-Santos S, Freeman GL Jr & Secor WE. Neonatal idiopathic exposure alters subsequent cytokine, pathology, and survival patterns in experimental *Schistosoma mansoni* infections. *J Exp Med* 1999; **189**: 637–645.
 - 65 Lewert RM & Mandlowitz S. Schistosomiasis: prenatal induction of tolerance to antigens. *Nature* 1969; **224**: 1029–1030.
 - 66 Lammie PJ, Hitch WL, Walker Allen EM, Hightower W & Eberhard ML. Maternal filarial infection as risk factor for infection in children. *Lancet* 1991; **337**: 1005–1006.
 - 67 Steel C, Guinea A, McCarthy JS & Ottesen EA. Long-term effect of prenatal exposure to maternal microfilariemia on immune responsiveness to filarial parasite antigens. *Lancet* 1994; **343**: 890–893.
 - 68 Mehta RS, Rodriguez A, Chico M, et al. Maternal geohelminth infections are associated with an increased susceptibility to geohelminth infection in children: a case-control study. *PLoS Negl Trop Dis* 2012; **6**: e1753.
 - 69 Elson LH, Days A, Calvopina M, et al. In utero exposure to *Onchocerca volvulus*: relationship to subsequent infection intensity and cellular immune responsiveness. *Infect Immun* 1996; **64**: 5061–5065.
 - 70 Nash TE, Cheever AW, Ottesen EA & Cook JA. Schistosome infections in humans: perspectives and recent findings. NIH conference. *Ann Intern Med* 1982; **97**: 740–754.
 - 71 Malhotra I, Ouma JH, Wamachi A, et al. Influence of maternal filariasis on childhood infection and immunity to *Wuchereria bancrofti* in Kenya. *Infect Immun* 2003; **71**: 5231–5237.
 - 72 Malhotra I, Mungai PL, Wamachi AN, et al. Prenatal T cell immunity to *Wuchereria bancrofti* and its effect on filarial

- immunity and infection susceptibility during childhood. *J Infect Dis* 2006; **193**: 1005–1013.
- 73 Malhotra I, Mungai P, Wamachi A, *et al.* Helminth- and Bacillus Calmette-Guerin-induced immunity in children sensitized in utero to filariasis and schistosomiasis. *J Immunol* 1999; **162**: 6843–6848.
- 74 Fine PE. Variation in protection by BCG: implications of and for heterologous immunity. *Lancet* 1995; **346**: 1339–1345.
- 75 Colditz GA, Berkey CS, Mosteller F, *et al.* The efficacy of bacillus Calmette-Guerin vaccination of newborns and infants in the prevention of tuberculosis: meta-analyses of the published literature. *Pediatrics* 1995; **96** (1 Pt 1): 29–35.
- 76 Gill TJ 3rd, Repetti CF, Metlay LA, *et al.* Transplacental immunization of the human fetus to tetanus by immunization of the mother. *J Clin Invest* 1983; **72**: 987–996.
- 77 Elliott AM, Mawa PA, Webb EL, *et al.* Effects of maternal and infant co-infections, and of maternal immunisation, on the infant response to BCG and tetanus immunisation. *Vaccine* 2010; **29**: 247–255.
- 78 Elliott A & Yazdanbakhsh M. Troubles never come alone. *Curr Opin HIV AIDS* 2012; **7**: 211–213.
- 79 Gallagher M, Malhotra I, Mungai PL, *et al.* The effects of maternal helminth and malaria infections on mother-to-child HIV transmission. *AIDS* 2005; **19**: 1849–1855.
- 80 Mpairwe H, Ndibazza J, Webb EL, *et al.* Exposure to hookworm prenatally and to worm infections in early childhood is inversely associated with eczema in childhood: results from a birth cohort in Uganda. *Allergy* 2013; **68**: (s67) 55.
- 81 Warner JO. Parasites and allergy. *Pediatr Allergy Immunol* 2009; **20**: 405.
- 82 King CL, Malhotra I, Mungai P, *et al.* B cell sensitization to helminthic infection develops in utero in humans. *J Immunol* 1998; **160**: 3578–3584.
- 83 Malhotra I, Ouma J, Wamachi A, *et al.* In utero exposure to helminth and mycobacterial antigens generates cytokine responses similar to that observed in adults. *J Clin Invest* 1997; **99**: 1759–1766.
- 84 King CL, Malhotra I, Wamachi A, *et al.* Acquired immune responses to *Plasmodium falciparum* merozoite surface protein-1 in the human fetus. *J Immunol* 2002; **168**: 356–364.
- 85 Novato-Silva E, Gazzinelli G & Colley DG. Immune responses during human *Schistosomiasis mansoni*. XVIII. Immunologic status of pregnant women and their neonates. *Scand J Immunol* 1992; **35**: 429–437.
- 86 Tweyongyere R, Mawa PA, Kihembo M, *et al.* Effect of praziquantel treatment of *Schistosoma mansoni* during pregnancy on immune responses to schistosome antigens among the offspring: results of a randomised, placebo-controlled trial. *BMC Infect Dis* 2011; **11**: 234.
- 87 Bal MS, Mandal NN, Das MK, Kar SK, Sarangi SS & Beuria MK. Transplacental transfer of filarial antigens from *Wuchereria bancrofti*-infected mothers to their offspring. *Parasitology* 2010; **137**: 669–673.
- 88 Guadalupe I, Mitre E, Benitez S, Chico ME, Nutman TB & Cooper PJ. Evidence for in utero sensitization to *Ascaris lumbricoides* in newborns of mothers with ascariasis. *J Infect Dis* 2009; **199**: 1846–1850.
- 89 Schneider H & Miller RK. Receptor-mediated uptake and transport of macromolecules in the human placenta. *Int J Dev Biol* 2010; **54**: 367–375.
- 90 Romia SA, Handoussa AE, Youseff SA & el Zayat MM. Transplacental transfer of schistosomal antigens and antibodies. *J Egypt Soc Parasitol* 1992; **22**: 575–582.
- 91 Hassan MM, Hassounah OA, Hegab M, Salah K, el-Mahrouky L & Galal N. Transmission of circulating schistosomal antigens from infected mothers to their newborns. *J Egypt Soc Parasitol* 1997; **27**: 773–780.
- 92 Attallah AM, Ghanem GE, Ismail H & El Waseef AM. Placental and oral delivery of *Schistosoma mansoni* antigen from infected mothers to their newborns and children. *Am J Trop Med Hyg* 2003; **68**: 647–651.
- 93 Malek A. Ex vivo human placenta models: transport of immunoglobulin G and its subclasses. *Vaccine* 2003; **21**: 3362–3364.
- 94 May K, Grube M, Malhotra I, *et al.* Antibody-dependent transplacental transfer of malaria blood-stage antigen using a human ex vivo placental perfusion model. *PLoS ONE* 2009; **4**: e7986.
- 95 Rahman MJ, Degano IR, Singh M & Fernandez C. Influence of maternal gestational treatment with mycobacterial antigens on postnatal immunity in an experimental murine model. *PLoS ONE* 2010; **5**: e9699.
- 96 Dauby N, Goetghebuer T, Kollmann TR, Levy J & Marchant A. Uninfected but not unaffected: chronic maternal infections during pregnancy, fetal immunity, and susceptibility to postnatal infections. *Lancet Infect Dis* 2012; **12**: 330–340.
- 97 Bisseye C, van der Sande M, Morgan WD, Holder AA, Pinder M & Ismaili J. *Plasmodium falciparum* infection of the placenta impacts on the T helper type 1 (Th1)/Th2 balance of neonatal T cells through CD4(+) CD25(+) forkhead box P3(+) regulatory T cells and interleukin-10. *Clin Exp Immunol* 2009; **158**: 287–293.
- 98 Eloi-Santos SM, Novato-Silva E, Maselli VM, Gazzinelli G, Colley DG & Correia-Oliveira R. Idiotypic sensitization in utero of children born to mothers with schistosomiasis or Chagas disease. *J Clin Invest* 1989; **84**: 1028–1031.
- 99 Fitzsimmons CM & Dunne DW. Survival of the fittest: allergology or parasitology? *Trends Parasitol* 2009; **25**: 447–451.
- 100 Bernardini R, Mistrello G, Novembre E, *et al.* Cross-reactivity between IgE-binding proteins from *Anisakis simplex* and *Dermatophagoides pteronyssinus*. *Int J Immunopathol Pharmacol* 2005; **18**: 671–675.
- 101 Nakazawa T, Khan AF, Yasueda H, *et al.* Immunization of rabbits with nematode *Ascaris lumbricoides* antigens induces antibodies cross-reactive to house dust mite *Dermatophagoides farinae* antigens. *Biosci Biotechnol Biochem* 2013; **77**: 145–150.
- 102 Santiago HC, LeeVan E, Bennuru S, *et al.* Molecular mimicry between cockroach and helminth glutathione S-transferases promotes cross-reactivity and cross-sensitization. *J Allergy Clin Immunol* 2012; **130**: 248–256 e9.
- 103 Amoah AS, Obeng BB, Larbi IA, *et al.* Peanut-specific IgE antibodies in asymptomatic Ghanaian children possibly caused by carbohydrate determinant cross-reactivity. *J Allergy Clin Immunol* 2013; **132**: 639–647.
- 104 Conrad ML, Ferstl R, Teich R, *et al.* Maternal TLR signaling is required for prenatal asthma protection by the nonpathogenic microbe *Acinetobacter lwoffii* F78. *J Exp Med* 2009; **206**: 2869–77.
- 105 Holt PG & Strickland DH. Soothing signals: transplacental transmission of resistance to asthma and allergy. *J Exp Med* 2009; **206**: 2861–4.
- 106 Hartgers FC, Obeng BB, Kruize YC, *et al.* Lower expression of TLR2 and SOCS-3 is associated with *Schistosoma haematobium* infection and with lower risk for allergic reactivity in children living in a rural area in Ghana. *PLoS Negl Trop Dis* 2008; **2**: e227.