

RESEARCH

Open Access

Treatment guided by rapid diagnostic tests for malaria in Tanzanian children: safety and alternative bacterial diagnoses

George Mtove¹, Ilse CE Hendriksen², Ben Amos³, Hedwiga Mrema³, Victor Mandia^{3†}, Alphaxard Manjurano⁴, Florida Muro⁴, Alma Sykes³, Helena Hildenwall⁵, Christopher JM Whitty⁶ and Hugh Reyburn^{6,7*}

Abstract

Background: WHO guidelines for the treatment of young children with suspected malaria have recently changed from presumptive treatment to anti-malarial treatment guided by a blood slide or malaria rapid diagnostic test (RDT). However, there is limited evidence of the safety of this policy in routine outpatient settings in Africa.

Methods: Children 3-59 months of age with a non-severe febrile illness and no obvious cause were enrolled over a period of one year in a malaria endemic area of Tanzania. Treatment was determined by the results of a clinical examination and RDT result, and blood culture and serum lactate were also collected. RDT-negative children were followed up over 14 days.

Results: Over the course of one year, 965 children were enrolled; 158 (16.4%) were RDT-positive and treated with artemether-lumefantrine and 807 (83.4%) were RDT-negative and treated with non-anti-malarial medicines. Compared with RDT-positives, RDT-negative children were on average younger with a lower axillary temperature and more likely to have a history of cough or difficulty in breathing. Six (0.6%) children became RDT-positive after enrolment, all of whom were PCR-negative for *Plasmodium falciparum* DNA at enrolment. In addition, 12 (1.2%) children were admitted to hospital, one with possible malaria, none of whom died. A bacterial pathogen was identified in 9/965 (0.9%) children, eight of whom were RDT-negative and one was RDT-positive, but slide-negative. Excluding three children with *Salmonella typhi*, all of the children with bacteraemia were ≤ 12 months of age. Compared to double-read research slide results RDTs had a sensitivity of 97.8% (95%CI 96.9-98.7) and specificity of 96.3% (95%CI 96.3-98.4).

Conclusions: Use of RDTs to direct the use of anti-malarial drugs in young children did not result in any missed diagnoses of malaria although new infections soon after a consultation with a negative RDT result may undermine confidence in results. Invasive bacterial disease is uncommon in children with non-severe illness and most cases occurred in infants with a current fever.

Background

The features of malaria overlap with those of other common illnesses and, where diagnosis is based on clinical history and examination alone, this often results in a large degree of overtreatment for malaria [1]. For many years this has been considered as a price worth paying to ensure prompt treatment of malaria in young

children, but over the last decade a number of important trends have driven a movement towards parasitological testing to guide the use of anti-malarial drugs [2]. These include the replacement of older anti-malarial drugs with the more expensive artemisinin combination therapy (ACT), the need to minimize selection pressure for drug-resistant strains of *Plasmodium falciparum*, the need for more reliable surveillance data as malaria has declined in many areas of Africa, and the relative neglect of alternative diagnoses to malaria [3-7].

* Correspondence: hugh.reyburn@lshtm.ac.uk

† Contributed equally

⁶London School of Hygiene and Tropical Medicine, Keppel St, London WC1E7HT, UK

Full list of author information is available at the end of the article

In parallel to this, accurate and affordable rapid diagnostic tests for malaria (RDTs) have become available over the last decade [8] and the 2010 WHO malaria treatment guidelines now propose that wherever possible all patients suspected of malaria should be tested by blood slide or RDT and only those with a positive test result should receive anti-malarial treatment [9]. While this should lead to a number of benefits, there is still relatively little direct evidence that non-malarial treatment in young febrile children in highly or moderately endemic areas is safe, with a concern that a significant number of malaria cases may be missed. The risks of 'test negative malaria' are consistently raised by clinicians cautious about this approach [10]. In addition clinicians need more evidence on the probabilities of different non-malarial diagnoses, especially those that require treatment with antimicrobial agents.

This study was conducted in a malaria endemic area of Tanzania to document the clinical outcomes and blood culture results of children with febrile illness whose treatment had been guided by the results of a commonly used RDT.

Methods

Study area

The study was conducted in the mother and child health (MCH) clinic of Teule Hospital, the District Designated Hospital of Muheza District, Tanzania. This serves a predominantly rural population exposed to perennial transmission of malaria with seasonal peaks coinciding with the short and long rainy seasons. In common with much of East Africa, malaria incidence has declined in recent years; in 1990 parasite prevalence in children under the age of five years was approximately 80% and this had declined to 30% in 2008 [11,12]. In order to facilitate patient follow-up the study area was defined as residence in one of the 85 villages from which there had been > 10 paediatric admissions to Teule Hospital in 2006.

Clinical and laboratory procedures

All children presenting to the MCH clinic were screened for the following inclusion criteria; age 3-59 months and resident within the study area, reported fever in the previous two days or current axillary temperature ≥ 37.5 C, no obvious non-malarial cause of fever and no known chronic illness. Children with evidence of severe illness were admitted to the paediatric ward and excluded from the study.

Following consenting procedures, a medical history and clinical examination were undertaken according to IMCI guidelines. Venous blood was drawn for culture (2-5ml), haemoglobin concentration (Hemocue™, Angelholm, Sweden), serum lactate (Lactate-Pro™,

Arkray Inc, Kyoto, Japan), HRP-2 based RDT for *P. falciparum* (Paracheck™, Orchid Biomedical, Mumbai, India).

Giemsa-stained blood slides were prepared from venous blood and independently double-read, with discordant results (either positive/negative discordance or > 2-fold density difference above 400 parasites/ μ L) resolved by an independent third reading. Microscopists were blinded to RDT results. Parasite densities were calculated from the geometric mean of the two closest counts of asexual parasites/200 white blood cells assuming 8,000 WBC/ μ L. Blood for culture was inoculated into a BactALERT™ Paediatric-fan bottle (bioMérieux, France) and incubated in the BacT/ALERT 3D automated microbial detection system. Cultures that flagged positive were identified by standard methods. Cultures positive for non-yeast fungi, coagulase-negative Staphylococcus, Micrococcus, Corynebacterium, Bacillus sp. or viridans group Streptococci were considered contaminants and classified as 'negative culture' unless a pathogenic organism was also isolated. The laboratory participated in a bacteriology external quality assurance programme coordinated by a reference laboratory in Moshi, Tanzania that participates successfully in external quality assurance programmes of the College of American Pathologists. At all visits, a malaria RDT, blood slide and blood spot for parasite DNA identification by polymerase chain reaction (PCR) were obtained according to established methodology [13].

Children were treated according to IMCI guidelines but anti-malarials or other medication with anti-malarial activity was avoided in RDT-negative children. RDT-negative children were seen and checked on Days 2, 7 and 14 following the initial visit (Day 0) with home visits the following day (and up to 2 more days until seen) if they did not attend. RDT-positive children were treated with artemether-lumefantrine (ALu) as recommended in the Tanzanian Guidelines for Malaria Diagnosis and Treatment (2006). RDT-positive children were not routinely followed up but results of positive blood cultures were promptly reviewed by a study clinician and, if the isolate was judged to be a pathogen, home visits were conducted for three consecutive days or until the child was seen and appropriate treatment and follow up was arranged. Children were also seen at additional times on parental demand and admissions to the hospital ward for children in the study were identified and clinical details obtained from hospital records.

Data management and analysis

Data were double-entered in MS Access and analysed in Stata-11. Simple proportions and confidence intervals were calculated for prevalence of pathogens, and risk of malaria in RDT-negative cases at follow-up. Sensitivity

and specificity of Paracheck against the gold standard of double-read research microscopy was undertaken on baseline samples. Groups were compared using Wilcoxon rank-sum test for continuous variables and chi-square test for proportions. The study was able to detect prevalence of pathogens in 800 RDT-negative patients at 1% with 95%CI 0.5-2% and a 5% prevalence with a 95% CI 3.6-6.6%.

Ethics

Written informed consent was obtained from the parent or caretaker of each child, or witnessed verbal consent if the parent could not read. The study was approved by the Ethical Review Boards of the National Institute for Medical Research in Tanzania (NIMR/HQ/R.8a/Vol.IX/785) and the London School of Hygiene & Tropical Medicine in the UK (LSHTM ethics # 5523).

Results

The study was conducted over one year from April 2009; 2592 children were screened, and 965 were eligible and enrolled (Figure 1). Of these, 158 were RDT-positive and treated with ALu and 807 were RDT-

negative and received non-anti-malarial treatment. Baseline characteristics by RDT result are shown in Table 1; children who were RDT-negative were on average younger with a lower axillary temperature and more likely to have a history of cough or difficulty in breathing (Table 1).

RDTs had a sensitivity of 98.6% (95%CI 97.8-99.3) and specificity of 97.3% (95%CI 96.3-98.4) when compared to double-read research slide results; 14.3% of blood slides were positive with a negative predictive value of RDT results of 99.8% (95%CI 99.4-100).

Of the 807 patients who were RDT-negative, 765 (94.8%) remained in the study for the 14 days of follow-up at which time 461 (60.3%) of whom were reported by the caretaker to be 'completely well'. Of 42 patients who exited before the end of the study, the parents of 6 children administered an anti-malarial drug during the 14-day follow-up period and all of these were RDT-negative at their next research visit and a further six children became RDT-positive after enrolment and were treated with ALu, two of whom were slide-positive with 16,676 and 1,234 asexual parasites/ μ l respectively and all of whom were PCR-negative for *P. falciparum* DNA at




Table 1 Clinical and laboratory characteristics of RDT positive compared to RDT-negative children at enrolment into the study

	RDT-Positive, n = 158	RDT-Negative, n = 807	p*
Age in months, median(mean)	26 (28)	16 (20)	< 0.001
Days ill, median(mean)	3 (3.8)	3 (3.3)	0.002
Haemoglobin g/dl, median(mean)	9.7 (9.6)	10.5 (10.4)	< 0.001
Resp. rate/min, median (mean)	48 (46.3)	48 (47.4)	0.16
WHO criteria of pneumonia, n (%) [†]	48 (30.4%)	417 (51.6%)	< 0.001
Diarrhoea or vomiting, n (%)	40 (25.3%)	155 (19.2%)	0.079
Axillary temperature, °C, median(mean)	38.1 (38.1)	37.4 (37.5)	< 0.001
Blood lactate, mmol/L, median(mean)	2.2 (2.5)	1.8 (2.0)	< 0.001

*Wilcoxon rank-sum test for continuous variables or chi-square test for proportions

[†] Cough or difficulty in breathing plus raised respiratory rate for age (> 40 breaths/min 12-59 months, > 50 breaths per minute 2-11 months of age)

enrolment. Twenty children were lost to follow-up of whom 6 (30%) were untraceable and 14 (70%) had left the study area but were reported (by telephone or friends) to be well at the time they exited the study. In addition, 12 children were admitted to hospital for whom details are provided in Table 2.

Diagnoses made by the study clinician at the first consultation are shown in Table 3. Almost half (383, 47.5%) of RDT-negative and 29 (18.4%) of RDT-positive children met WHO criteria for non-severe pneumonia and were prescribed amoxicillin and an additional 34 (4.2%) of RDT-negative and 19(12.0%) of RDT-positive children also met these criteria but were not treated with amoxicillin. There was no evidence of a trend of the proportion of children with raised respiratory rate or a recent history of 'cough or difficulty in breathing' with month of the study that might indicate increased prescriber confidence to withhold antibiotic treatment ($p = 0.37$ and $p = 0.43$ respectively). While 73% of respiratory rate values were even, there was no apparent preference for values at or above cut-off levels; the commonest values were 36/min (9.7% of values) and 48/min (15.4% of values).

A bacterial pathogen was identified from blood culture in 9/965 (0.9%) children in the study, 8 of whom were RDT-negative and one was RDT-positive but slide-negative (Table 4). In addition, an organism judged to be a contaminant was isolated in 56 (5.8%) of cultures. All of the bacteraemic children were successfully treated as outpatients and the majority (7/9) were treated with amoxicillin at the first consultation before the blood culture result was available. Clinical and laboratory features associated with bacteraemia are shown in Table 5, none of which reached 5% significance although 'current fever' (measured axillary temperature $\geq 37.5^{\circ}\text{C}$) was positively associated with bacteraemia ($p = 0.08$) and with sensitivity and specificity of 77.8% (95%CI 75.2-80.4) and 46.5% (95%CI 43.4-49.6) respectively. In addition, with the exception of children with *S. typhi*, all of the children with bacteraemia were 12 months of age or younger

Discussion

While age, haemoglobin level and presence of respiratory symptoms provide some guide to diagnostic probability, clinical predictors of malaria are still insufficient

Table 2 Diagnosis, treatment and outcome of 12 children admitted to hospital during the 14 days of follow-up.

Age (months)	Diagnosis or clinical syndrome	RDT result	Blood slide result	Treatment	Outcome
24 m	Acute diarrhoea	Negative	Negative	Quinine and gentamycin	Discharged well
7 m	Acute diarrhoea	Negative	Negative	IV rehydration	Discharged well
22 m	Acute diarrhoea	Negative	Negative	IV rehydration	Discharged well
8 m	Suspected pneumonia	Negative	Negative	Cotrimoxazole	Discharged well
12 m*	Clinical diagnosis of malaria	Negative	Not available	Quinine	Discharged well
8 m	Acute diarrhoea	Negative	Not available	IV rehydration	Discharged well
7 m	Febrile convulsion	Not done	Not available	Paracetamol	Discharged well
24 m*	Fever and vomiting	Not done	Not available	Quinine	Discharged well
18 m*	Fever and vomiting	Not done	Not available	Quinine	Discharged well
29 m	Fever, convulsion, prostration	Positive	Not available	Quinine	Discharged well
18 m*	Suspected dysentery	Negative	Negative	IV rehydration	Discharged well
18 m	Pneumonia	Not done	Negative	Amoxicillin	Discharged well

*Admitted to non-study facility

Table 3 Diagnoses made by examining clinician at enrolment by RDT result

	RDT-Negative			RDT-Positive		
	First diagnosis	Secondary diagnoses	Total (%)	First diagnosis	Secondary diagnoses	Total (%)
Upper resp. tract infection	150	5	155 (14.2)	4	6	10 (3.6)
Non-specific fever	182	6	188 (17.2)	0	0	0
Diarrhoea	72	40	112 (10.2)	2	7	9 (3.2)
Pneumonia	375	8	383 (35.0)	20	9	29 (10.6)
Malaria	0	0		127	31	158 (57.0)
Anaemia	0	206	206 (18.8)	0	59	59 (21.3)
Bronchitis/wheeze	0	3	3 (0.3)	0	0	0
Other	28	19	47 (4.4)	5	7	11 (4.1)
TOTAL	807	287	1,094 (100)	158	119	277 (100)

to guide treatment. The advent of RDTs provides the potential to target anti-malarial treatment on a scale that has hitherto been impossible. However, doubts remain in the minds of many clinicians about the safety of not treating parasite-negative patients for malaria, and these have probably contributed to the observation from a number of studies in Africa that up to half of those with a negative rapid test or blood slide result for malaria are treated with anti-malarials anyway. Additionally the question 'if it is not malaria what is it?' is important for framing a rational diagnostic and treatment strategy. Appropriate use of anti-microbial drugs is a key issue given the high mortality of children admitted to hospital with invasive bacterial disease but whose treatment is restricted by high levels of resistance to commonly used agents [14].

This study adds to other evidence that anti-malarial treatment based on parasitological testing misses few

true cases of malaria. Njama-Meyer *et al* used expert microscopy to guide the treatment of 2,359 illness episodes in children under the age of 10 years in a malaria-endemic area of Uganda; fewer than 1% of initial blood slide readings were false-negatives, and only 13 (0.8%) of 1,602 slide-negative illness episodes became positive over the following even days, and none of these was severely ill [15]. D'Acromont *et al* followed up 603 RDT-negative children who had not been treated for malaria and only three children developed a positive result in the following week, all of whom were successfully treated [16].

Although the study results are reassuring, the policy of RDT-directed treatment for suspected malaria represents a substantial change in the delivery of routine care in Africa and some questions still remain. Firstly, it is impossible to distinguish between 'new' or 'missed' diagnoses of malaria during routine follow-up of RDT-

Table 4 Clinical and laboratory data on the 9 children whose blood grew a bacterial pathogen

Age (mon)	Blood slide/RDT	Organism	Summary clinical and laboratory data	*Antibiotic susceptible	Antibiotic resistant
8	Neg/Neg	<i>Acinetobacter Iwoffii</i>	7 days cough, temp 36.8°, pulse 138/min, resp. rate 63/min, lactate 1.4 mmol/L	amox, cip, ceft, gen, nal	chlor, cip
8	Neg/Neg	β -Haemolytic streptococcus	3 days cough, temp 37.2°, pulse 144/min, resp. rate 56/min, lactate 1.1 mmol/L	eryth, ceft.	Pen, amox.
7	Neg/Neg	Gram-negative coccus	3 days cough, temp 37.4°, pulse 167/min, resp. rate 64/min, lactate 1.8 mmol/L	No data	
8	Neg/Neg	<i>Haemphilus influenzae</i>	3 days cough and coryza, temp 37.7°, pulse 150/min, lactate 3.1 mmol/L	amox, ceft, gen	chlor, cip, nal
10	Neg/Neg	Non-Typhi <i>Salmonella</i>	2 days cough, temp 39.5°, pulse 156, resp rate 64, lactate 2.0 mmol/L	amox, chlor, cip, ceft, gen, nal	
43	Neg/Neg	<i>Salmonella Typhi</i>	2 days cough, temp 39.2, pulse 145/min, resp rate 39/min, lactate 1.9 mmol/L	cip, ceft, gen	chlor, cotrim, amox
38	Neg/Neg	<i>Salmonella Typhi</i>	4 days diarrhoea, temp 38.1°, pulse 120/min, resp rate 42/min, lactate 3.3 mmol/L	chl, cip, cef, gen	amox, cotrim.
47	Neg/Neg	<i>Salmonella Typhi</i>	7 days febrile illness, temp 38.8, pulse 140/min, resp rate 36/min, lactate 0.9 mmol/L	chl, cip, cef, nal	cotrim, amox
12	Neg/Pos	<i>Staphylococcus aureus</i>	1 day cough, temp 37.4°, pulse 163/min, resp. rate 52/min, lactate 0.9 mmol/L	chl, eryth, gen	cotrim, amox

*Anti-microbial agents tested for in vitro susceptibility: amoxicillin (amox), chloramphenicol (chlor), cotrimoxazole (cotrim), ciprofloxacin (cip), erythromycin (eryth), oxacillin(pen), ceftriaxone (ceft), nalidixic acid (nal).

Table 5 Clinical and laboratory features of children by presence of a bacterial pathogen isolated from blood culture

	Bacterial pathogen, n = 9	No bacterial pathogen, n = 956	p [†]
Median (mean) age in months	10 (20.0)	17 (21.0)	0.56
Days of illness, median (mean)	3 (3.6)	3 (3.4)	0.68
Cough or difficulty in breathing* (%)	7 (77.8%)	601 (62.8%)	0.35
Diarrhoea or vomiting* (%)	1 (11.1%)	194 (20.3%)	0.49
Median (mean) axillary temp, °C	38.1 (38.2)	37.5 (37.6)	0.08
Respiratory rate, median (mean)/min	52 (51.8)	48 (47.2)	0.15
Median (mean) blood lactate mmol/L	1.9 (2.1)	1.8 (2.0)	0.94
Median (mean) haemoglobin, g/dL	10.2 (10.4)	10.4 (10.3)	0.93
WHO criteria of non-severe pneumonia (%)	5 (55.6%)	460 (48.1%)	0.65

* Symptoms reported at any time in the previous 2 days

negative patients. As observed in the current study, a significant minority of children will be admitted for malaria within days of being seen with a non-severe febrile illness and a negative RDT result. Although the PCR primers used in this study were not the most sensitive, the results suggest that the malaria diagnoses following enrolment into the study were 'new' infections and that it is rational and safe to treat RDT-negative children for a non-malarial illness. However, the perception of clinicians and parents may be that a child with malaria has been denied the best treatment as a result of application of new guidelines with RDTs. Given evidence of a strong preference for a diagnosis of malaria among parents and clinicians [17], such an interpretation is quite likely and may undermine efforts to implement the policy of RDT-directed anti-malarial treatment. To guard against this, clinicians should be reminded that 'no test is perfect' and to advise parents to bring their child back if they remain unwell. Secondly, there is still a possibility that presumptive treatment for malaria in all children with fever might provide a benefit due to intermittent prophylaxis that might protect against malaria and/or anaemia. Recent results of trials of intermittent treatment for malaria in children suggest that this is possible, and a cluster randomized trial of presumptive compared to RDT-directed anti-malarial treatment is in progress [18].

The commonest indication for anti-microbial treatment in our study was the IMCI diagnosis of non-severe pneumonia. While the overlap between malaria and pneumonia has been previously described both in severe and non-severe illness, the proportion of children meeting WHO criteria for non-severe pneumonia was higher than expected from previous studies [15,19-21]. Preferential recording by study clinicians in order to justify prescribing an anti-microbial drug seems unlikely as the number of children meeting the WHO definition was slightly higher than the number actually diagnosed. In addition, there was no evidence of a change in the proportion diagnosed with pneumonia as the study

progressed, nor was there evidence of clustering of respiratory counts near the cut-off values that define raised levels. It's possible that 'cough or difficulty breathing' was over-interpreted or there may be local variations in the perception or reporting of respiratory problems in children [22]. In addition, the finger-prick undertaken approximately 15 minutes before examination may have resulted in transiently increased respiratory rates.

As new guidelines and RDTs are introduced across Africa it is possible that there will be a compensatory over-use of the IMCI diagnosis of non-severe pneumonia but this will be difficult to assess since there is no gold standard definition of the diagnosis of pneumonia. Even of the true pneumonias some will be viral, including the vaccine-preventable RSV pneumonias, where antibiotics will make little positive impact on outcome [23,24]. In addition, studies of the quality of care suggest that IMCI procedures to diagnose pneumonia, in particular examining the chest and recording respiratory rate, are not well undertaken outside research settings [25,26].

Consistent with at least one other study, we found that bacteraemia was uncommon in children with non-severe illness, especially in children who were RDT-positive [27]. However, blood culture lacks sensitivity and early recognition and treatment of blood stream infections has the potential to avert progression to severe disease with high associated mortality [14,28]. While RDTs can improve targeting of anti-malarial drugs there is no comparable test for bacterial disease. The POC measure of blood lactate was increased in children who were RDT-positive compared to RDT-negative, but was unhelpful in distinguishing between children with and without invasive bacterial infection. While the small numbers of children with bacteraemia in our study reduced the likelihood of finding a statistically significant association with clinical and laboratory features, our findings are consistent with other studies that bacteraemia occurs predominantly in young children, up to

three-quarters of whom present with a 'current fever'. However, these indicators lack both sensitivity and specificity and there is still a need for more accurate measures of bacterial infection; devices measuring acute phase proteins such as C-reactive protein or procalcitonin merit further research [29].

This study has a number of inevitable limitations. The study was conducted in hospital outpatients where the patient population may differ in a number of ways from those attending primary health facilities generally. The accuracy of RDT interpretation may have been higher than under routine practice, and certainly higher than routine microscopy in some clinics in the region. Clinicians were aware that they were being observed (in the sense that their findings and prescriptions were being recorded) and may have been more likely to follow IMCI guidelines and to prescribe antibiotics, thus over-estimating antibiotic usage that would occur under similar conditions in routine practice. The clinical care provided in the study followed IMCI guidelines and did not have access to an otoscope or urine examination resulting in possible neglect of otitis media or urinary tract infections. The lack of validating investigations for the IMCI diagnosis of pneumonia has already been discussed.

In conclusion, this study demonstrates that use of RDT's to direct the use of anti-malarial drugs in young children is a safe strategy and did not result in any missed diagnoses of malaria. However, new episodes of malaria do occur within a short period of an initial consultation in malaria endemic areas and these may undermine confidence in parasitological testing for malaria. Invasive bacterial disease is relatively uncommon in children with non-severe illness; while most cases occurred in infants with a current fever there is a need for more specific diagnostic tests to detect these infections at the point of care.

Conflict of interest

The authors declare that they have no competing interests.

Acknowledgements

Thanks to the staff and patients of Teule Hospital for their cooperation during study. The following were involved in clinical and laboratory data collection: Aikande Shoo, Selina Wycliffe, Juma Kimera, Levogat Tarimo, Weston Lemmanya, Colin Sutherland and Rachel Hallet quality-checked the PCR results. Our sympathies to the friends and family of Dr Victor Mandia who died earlier this year. Core funding for the study was provided by the Bill and Melinda Gates Foundation though the ACT Consortium awarded to LSHTM.

Author details

¹National Institute for Medical Research, Amani Centre, Muheza, Tanga, Tanzania. ²Mahidol-Oxford Research Unit, Mahidol University, Bangkok 10500, Thailand. ³Teule Hospital, Muheza, Tanzania. ⁴Kilimanjaro Christian Medical

Centre, Moshi, Tanzania. ⁵Division of Global Health, Karolinska Institute, Nobels V13, 17177 Stockholm, Sweden. ⁶London School of Hygiene and Tropical Medicine, Keppel St, London WC1E7HT, UK. ⁷Joint Malaria Programme, P.O. Box 2228, KCMC, Moshi, Tanzania.

Authors' contributions

GM was responsible for clinical data collection and drafting the MS, IH was responsible for drafting study documents and supervising data collection, BA was responsible for the laboratory work and provided critical review of the MS, HM was responsible for data collection, VM was responsible for clinical care, AM was responsible for the PCR analysis, FM assisted in the analysis and provided critical review of the MS, AS contributed to the design of the study and provided critical review of the MS, HH assisted in the analysis and writing and provided critical review of the MS, CW initiated the idea for the study and contributed to the writing of the MS, HR was involved in the design of the study, analysed the data and co-wrote the MS. All authors read and approved the final manuscript.

Received: 8 June 2011 Accepted: 6 October 2011

Published: 6 October 2011

References

1. Chandramohan D, Jaffar S, Greenwood B: Use of clinical algorithms for diagnosing malaria. *Trop Med Int Health* 2002, **7**:45-52.
2. D'Acremont V, Lengeler C, Mshinda H, Mtasiwa D, Tanner M, Genton B: Time to move from presumptive malaria treatment to laboratory-confirmed diagnosis and treatment in African children with fever. *PLoS Med* 2009, **6**:e252.
3. Wongsrichanalai C, Baird K: Linking malaria rapid diagnostic test (RDT) and new drug combinations. *BMJ* 2004, *bmj.com* July 23.
4. Yeung S, Van Damme W, Socheat D, White NJ, Mills A: Cost of increasing access to artemisinin combination therapy: the Cambodian experience. *Malar J* 2008, **7**:84.
5. Zurovac D, Ndhlovu M, Sipilanyambe N, Chanda P, Hamer DH, Simon JL, Snow RW: Paediatric malaria case-management with artemether-lumefantrine in Zambia: a repeat cross-sectional study. *Malar J* 2007, **6**:31.
6. Lubell Y, Reyburn H, Mbakilwa H, Mwangi R, Chonya K, Whitty CJ, Mills A: The cost-effectiveness of parasitologic diagnosis for malaria-suspected patients in an era of combination therapy. *Am J Trop Med Hyg* 2007, **77**:128-132.
7. Yeung S, Van Damme W, Socheat D, White NJ, Mills A: Access to artemisinin combination therapy for malaria in remote areas of Cambodia. *Malar J* 2008, **7**:96.
8. Murray CK, Gasser RA, Magill AJ, Miller RS: Update on rapid diagnostic testing for malaria. *Clin Microbiol Rev* 2008, **21**:97-110.
9. WHO: Guidelines for malaria diagnosis and treatment. *WHO-Roll Back Malaria* WHO-Geneva; 2010, vol. ISBN: 9789241547925.
10. Chandler CI, Jones C, Boniface G, Juma K, Reyburn H, Whitty CJ: Guidelines and mindlines: why do clinical staff over-diagnose malaria in Tanzania? A qualitative study. *Malar J* 2008, **7**:53.
11. Bernard J, Mtove G, Mandike R, Mtei F, Maxwell C, Reyburn H: Equity and coverage of insecticide-treated bed nets in an area of intense transmission of *Plasmodium falciparum* in Tanzania. *Malar J* 2009, **8**:65.
12. Maxwell CA, Chambo W, Mwaimu M, Magogo F, Carneiro IA, Curtis CF: Variation of malaria transmission and morbidity with altitude in Tanzania and with introduction of alphacypermethrin treated nets. *Malar J* 2003, **2**:28.
13. Snounou G, Viriyakosol S, Zhu XP, Jarra W, Pinheiro L, do Rosario VE, Thaithong S, Brown KN: High sensitivity of detection of human malaria parasites by the use of nested polymerase chain reaction. *Mol Biochem Parasitol* 1993, **61**:315-320.
14. Nadjim B, Amos B, Mtove G, Ostermann J, Chonya S, Wangai H, Kimera J, Msuya W, Mtei F, Dekker D, Malahiyo R, Olomi R, Crump JA, Whitty CJ, Reyburn H: WHO guidelines for antimicrobial treatment in children admitted to hospital in an area of intense *Plasmodium falciparum* transmission: prospective study. *BMJ* 2010, **340**:c1350.
15. Njama-Meya D, Clark TD, Nzarubara B, Staedke S, Kanya MR, Dorsey G: Treatment of malaria restricted to laboratory-confirmed cases: a prospective cohort study in Ugandan children. *Malar J* 2007, **6**:7.

16. d'Acromont V, Malila A, Swai N, Tillya R, Kahama-Marro J, Lengeler C, Genton B: **Withholding antimalarials in febrile children who have a negative result for a rapid diagnostic test.** *Clin Infect Dis* 2010, **51**:506-511.
17. Reyburn H, Mbatia R, Drakeley C, Carneiro I, Mwakasungula E, Mwerinde O, Saganda K, Shao J, Kitua A, Olomi R, Greenwood BM, Whitty CJ: **Overdiagnosis of malaria in patients with severe febrile illness in Tanzania: a prospective study.** *Bmj* 2004, **329**:1212.
18. Konate AT, Yaro JB, Ouedraogo AZ, Diarra A, Gansane A, Soulama I, Kangoye DT, Kabore Y, Ouedraogo E, Ouedraogo A, Tiono AB, Ouedraogo IN, Chandramohan D, Cousens S, Milligan PJ, Sirima SB, Greenwood B, Diallo DA: **Intermittent preventive treatment of malaria provides substantial protection against malaria in children already protected by an insecticide-treated bednet in Burkina Faso: a randomised, double-blind, placebo-controlled trial.** *PLoS Med* 2011, **8**: e1000408.
19. Kallander K, Nsungwa-Sabiiti J, Peterson S: **Symptom overlap for malaria and pneumonia—policy implications for home management strategies.** *Acta Trop* 2004, **90**:211-214.
20. O'Dempsey TJ, McArdle TF, Laurence BE, Lamont AC, Todd JE, Greenwood BM: **Overlap in the clinical features of pneumonia and malaria in African children.** *Trans R Soc Trop Med Hyg* 1993, **87**:662-665.
21. English M, Punt J, Mwangi I, McHugh K, Marsh K: **Clinical overlap between malaria and severe pneumonia in Africa children in hospital.** *Trans R Soc Trop Med Hyg* 1996, **90**:658-662.
22. Hildenwall H, Rutebemberwa E, Nsabagasani X, Pariyo G, Tomson G, Peterson S: **Local illness concepts—implications for management of childhood pneumonia in eastern Uganda.** *Acta Trop* 2007, **101**:217-224.
23. Weber MW, Dackour R, Usen S, Schneider G, Adegbola RA, Cane P, Jaffar S, Milligan P, Greenwood BM, Whittle H, Mulholland EK: **The clinical spectrum of respiratory syncytial virus disease in The Gambia.** *Pediatr Infect Dis J* 1998, **17**:224-230.
24. Weber MW, Mulholland EK, Greenwood BM: **Respiratory syncytial virus infection in tropical and developing countries.** *Trop Med Int Health* 1998, **3**:268-280.
25. English M, Esamai F, Wasunna A, Were F, Ogutu B, Wamae A, Snow RW, Peshu N: **Delivery of paediatric care at the first-referral level in Kenya.** *Lancet* 2004, **364**:1622-1629.
26. Reyburn H, Mwakasungula E, Chonya S, Mtei F, Bygbjerg I, Poulsen A, Olomi R: **Clinical assessment and treatment in paediatric wards in the north-east of the United Republic of Tanzania.** *Bull World Health Organ* 2008, **86**:132-139.
27. Brent AJ, Ahmed I, Ndiritu M, Lewa P, Ngetsu C, Lowe B, Bauni E, English M, Berkley JA, Scott JA: **Incidence of clinically significant bacteraemia in children who present to hospital in Kenya: community-based observational study.** *Lancet* 2006, **367**:482-488.
28. Berkley JA, Lowe BS, Mwangi I, Williams T, Bauni E, Mwarumba S, Ngetsu C, Slack MP, Njenga S, Hart CA, Maitland K, English M, Marsh K, Scott JA: **Bacteremia among children admitted to a rural hospital in Kenya.** *N Engl J Med* 2005, **352**:39-47.
29. Diez-Padr s N, Bassat Q, Machevo S, Quinto L, Morais L, Nhampossa T, O'Callaghan-Gordo C, Torres A, Alonso PL, Roca A: **Procalcitonin and C-reactive protein for invasive bacterial pneumonia diagnosis among children in Mozambique, a malaria-endemic area.** *PLoS One* 2010, **5**: e13226.

doi:10.1186/1475-2875-10-290

Cite this article as: Mtove et al.: Treatment guided by rapid diagnostic tests for malaria in Tanzanian children: safety and alternative bacterial diagnoses. *Malaria Journal* 2011 **10**:290.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

