



Detection of lipoarabinomannan (LAM) in urine is indicative of disseminated TB with renal involvement in patients living with HIV and advanced immunodeficiency: evidence and implications

Stephen D. Lawn^{a,b,*} and Ankur Gupta-Wright^{a,c}

^aDepartment of Clinical Research, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, UK;

^bThe Desmond Tutu HIV Centre, Institute of Infectious Disease and Molecular Medicine, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa; ^cMalawi-Liverpool-Wellcome Trust Clinical Research Program, University of Malawi College of Medicine, Blantyre, Malawi

*Corresponding author: Tel: +44 20 7927 2546; E-mail: stephen.lawn@lshtm.ac.uk

Received 2 November 2015; revised 10 December 2015; accepted 11 December 2015

TB is the leading cause of HIV/AIDS-related deaths globally. New diagnostic tools are urgently needed to avert deaths from undiagnosed HIV-associated TB. Although simple assays that detect lipoarabinomannan (LAM) in urine have been commercially available for years, their specific role and utility were initially misunderstood, such that they have been slower to emerge from the diagnostics pipeline than otherwise might have been expected. In this article, we review and explain how urine-LAM assays should be understood as diagnostics for disseminated TB in HIV-positive patients with advanced immunodeficiency, in whom haematogenous TB dissemination to the kidneys serves as the primary mechanism by which LAM enters the urine. These insights are critical for the appropriate design of studies to evaluate these assays and to understand how they might be most usefully implemented. This understanding also supports the 2015 WHO recommendations on the restricted use of these assays in sick HIV-positive patients with advanced immunodeficiency.

Keywords: Diagnosis, Disseminated TB, HIV-associated TB, LAM, Lipoarabinomannan, Urine

Introduction

In this new era of the United Nations Sustainable Development Goals (SDGs), the international community has committed to ending the HIV/AIDS and TB epidemics by 2030.¹ In 2014, the World Health Assembly adopted the WHO's post-2015 End TB Strategy, with targets to assess progress using data from 2015 as the baseline.² Goals to be achieved by 2030 include an 80% reduction in new TB cases and a 90% reduction in TB deaths. Furthermore, achieving an incidence of <10 cases per 100 000 people per year was defined as being sufficient to 'end' the global TB epidemic.

The remaining task, however, is huge.³ WHO reported that in 2014 alone, 9.6 million people developed TB (12% HIV co-infected) and 1.4 million died from TB disease (43% HIV co-infected). Overall, TB remains the leading cause of HIV-related deaths globally. A meta-analysis of hospital post-mortem studies of HIV-related deaths included nine studies from across sub-Saharan Africa.⁴ These found that between 32 and 67% (pooled summary estimate, 43.2%; 95%CI 38.0–48.3) of HIV-positive adult cadavers had evidence of TB at autopsy, and that in over 90% of cases it was deemed the primary cause of

death.⁴ Disease was disseminated in 87.9% of TB cases, and remained undiagnosed at death in nearly half of cases (45.8%). These data highlight the critical need to improve the ascertainment of disseminated HIV-associated TB. To achieve its ambitious goals, the WHO End TB strategy has outlined the requirement for 'intensified research and innovation',¹ including the development and implementation of new, simple and accurate diagnostic assays for TB.

For many years, the development of point-of-care assays has been foremost on the TB research agenda. One such assay methodology for TB diagnosis to emerge over the past 20 years has been the detection of the mycobacterial cell wall glycolipid antigen lipoarabinomannan (LAM) in urine using immune-capture assays. Much of the pioneering work on this was led by Prof. Stefan Svenson at the Karolinska Institute in Sweden, who demonstrated the proof of concept of urine-LAM detection as a means for TB diagnosis.⁵ This early methodology required complex sample processing, which was beyond the capacity of most laboratories in high TB burden countries, and this was viewed as an insurmountable obstacle to implementation. However, an ELISA that required only minimal urine processing was developed and marketed just over 10 years ago. The results of the first

assessment of its diagnostic accuracy in Tanzania were very encouraging,⁶ with high specificity among asymptomatic volunteers and high sensitivity in patients with laboratory-confirmed pulmonary TB. Moreover, diagnostic accuracy was similar in HIV-positive and HIV-negative patients. Had these findings been replicated in larger studies in different settings, this assay would have constituted a substantial breakthrough.

Optimism regarding this assay quickly evaporated, however, as many subsequent studies reported poor diagnostic accuracy. A meta-analysis in 2011⁷ concluded these assays had ‘suboptimal sensitivity for routine clinical use’ and many dismissed urine-LAM detection as a viable diagnostic methodology. In retrospect, many of the earlier studies targeted non-ideal patient groups (HIV-negative patients or HIV-positive patients with high CD4 counts). However, progress came when the assay was applied to HIV-positive adults with advanced immunodeficiency,⁸ in whom specificity was high and sensitivity, although only moderate at best, was highest among the sickest, most vulnerable patients in whom slow culture-based diagnosis was otherwise often required. The next breakthrough came with the development of a 30-minute lateral-flow assay (strip-test) version of the assay marketed as ‘Determine TB-LAM’ (Alere Inc. Waltham, MA, USA). This assay format required neither laboratory infrastructure nor equipment, and was far lower cost per test than the Xpert MTB/RIF assay (Cepheid, Sunnyvale, CA, USA). The results of the ELISA and lateral flow assay formats were remarkably similar, both showing markedly much greater sensitivity among those patients with the lowest CD4 counts.⁹ Thus, urine LAM detection emerged as a means for rapid diagnosis of TB in people living with HIV and advanced immunodeficiency.

Urine-LAM immune-capture assays have not been embraced by the HIV and TB communities as fully or rapidly as one might have expected. We propose that this is not only due to unfavourable data arising from earlier studies in which the assay was evaluated in inappropriate or non-ideal patient groups and using sub-optimal study methodologies. We believe that, in addition to these factors, the fundamental basis of the assays had not been correctly understood—in particular, the mechanism by which LAM enters urine. At first, this might seem an esoteric, academic issue, but we highlight why this issue is of fundamental importance for the appropriate understanding and application of these assays.

Early literature concerning urine-LAM assays presumed that LAM-antigenuria resulted from simple renal filtration of free circulating LAM in the bloodstream.^{6,10} This notion seemed plausible since it was known that LAM was produced in large quantities by *Mycobacterium tuberculosis* (MTB) at sites of disease from which it could enter the blood-stream.¹¹ Moreover, as a 17 kDa glycolipid (similar in size to myoglobin) it should readily traverse the renal glomerular basement membrane. Once produced at sites of TB, whether pulmonary or extrapulmonary, it was known that LAM was detectable in blood, albeit following extensive sample processing in view of the high degree of immune-complexing with host anti-LAM antibodies.¹² Renal filtration of circulating LAM also plausibly explained why the assay detected both pulmonary and extrapulmonary forms of TB. However, here we present a range of direct and indirect evidence that collectively indicates that LAM antigenuria results from renal involvement with TB and not from simple renal filtration of LAM from the bloodstream into urine (Table 1).

Simple filtration of LAM at the renal glomerulus is implausible

In addition to being immune-complexed within blood by anti-LAM antibodies, LAM in the bloodstream is also incorporated within high density lipoprotein (HDL) particles.¹³ Both immune-complexed or HDL-associated LAM would be unable to pass from the systemic circulation through the intact basement membrane in the renal glomerulus.¹⁴ Whilst glomerular dysfunction might hypothetically allow LAM-containing-complexes to pass into the urine, clinical studies have not found associations between urine-LAM detection and proteinuria (a marker for glomerular damage),^{15,16} and there is no histological evidence of glomerular damage at autopsy in patients who were urine-LAM positive.¹⁷

LAM is released in large quantities by actively replicating or dying mycobacteria, serving as an immune-modulating virulence factor. If LAM were freely filtered into urine, it would be expected that during early TB therapy when massive mycobacterial killing occurs, urine concentrations of LAM would rise in parallel with blood concentrations. However, the fact that urine-LAM levels do not increase during early TB treatment is suggestive of a mechanism other than simple renal filtration.¹⁵

Urine microbiology supports the renal TB hypothesis (Table 1)

Urine culture has previously been shown to provide a high diagnostic yield in an American cohort of patients with AIDS and extra-pulmonary TB in whom 77% tested urine-culture- positive.¹⁸ If the presence of LAM in the urine of HIV-infected patients were the result of renal TB, one would expect that *M. tuberculosis* would be culturable from LAM-positive urine samples. However, data correlating urine LAM status and urine culture status are unfortunately lacking, at present. Nevertheless, other important evidence comes from testing of urine samples with the Xpert MTB/RIF assay. The Xpert MTB/RIF assay detects the presence of whole MTB bacilli through amplification of organism-associated DNA (rather than detection of free DNA).^{19,20} Studies testing the urine of HIV-TB patients using TB-LAM and Xpert MTB/RIF have shown a large overlap between the assays, with ≥50% of LAM-positive urine samples also testing positive by Xpert MTB/RIF.^{21–23} The detection of whole MTB bacilli in urine of these LAM-positive patients strongly corroborates the notion of renal TB as the source of urinary-LAM.

Evidence of renal TB from post-mortem studies

Renal TB in the form of microabscesses in this patient population arises as a result of haematogenous dissemination and is a common autopsy finding in patients who have died from disseminated HIV-associated TB (50–69%).^{24–27} In patients with advanced HIV-infection (typically CD4 cell counts of <100 cells/mm³) a strong association between MTB bacteraemia and urine LAM-positivity has been observed, with 70–90% of bacteraemic patients also having detectable LAM-antigenuria.^{28–31} Renal TB readily provides a mechanism for this association.

The strongest evidence explaining the mechanism of urine LAM-antigenuria is provided by post-mortem studies. Cox et al. demonstrated histological evidence of renal TB in the majority

Table 1. Direct and indirect evidence that urine lipoarabinomannan (LAM) antigenuria is due to renal involvement with haematogenously disseminated TB, and not free filtration of LAM into the urine

Observation	Evidence	References
LAM is unlikely to be able to pass from systemic circulation into the urine via the renal glomerulus	<ul style="list-style-type: none"> LAM in the bloodstream is mostly immune-complexed or bound to high-density lipoprotein These LAM containing molecules are too large to freely filter in the renal glomerulus 	Sada et al. 1992 ¹² Sakamuri et al. 2013 ¹³ Haraldsson et al. 2008 ¹⁴
LAM concentration in the urine does not increase during the first weeks of TB treatment	<ul style="list-style-type: none"> Serum LAM concentration likely increases early after the massive mycobactericidal effect of anti-TB treatment. If LAM was freely filtered in the kidneys, LAM concentration in the urine would also be expected to increase early after treatment 	Wood et al. 2012 ¹⁵ Bekker et al. 1998 ⁴⁵
Most LAM-positive urines test XpertMTB/RIF-positive	<ul style="list-style-type: none"> Xpert MTB/RIF detects <i>M. tuberculosis</i> DNA in whole bacilli (not free DNA), suggesting most LAM-positive urines contain whole <i>M. tuberculosis</i> bacilli 	Wood et al. 2012 ¹⁵ Lawn et al. 2012 ²¹ Blakemore et al. 2010 ¹⁹ Manabe et al. 2014 ²⁸ Nakiyingi et al. 2015 ²⁹ Nakiyingi et al. 2014 ³⁰ Lawn & Kerkhoff 2015 ³¹ Lanjewar et al. 2011 ²⁵ Ansari et al. 2002 ²⁴ Cox et al. 2015 ¹⁷ Cox et al. 2015 ¹⁷
Frequent LAM-positive urine in patients with <i>M. tuberculosis</i> bacteraemia	<ul style="list-style-type: none"> The strong association between <i>M. tuberculosis</i> bacteraemia and LAM-positive urine is very plausibly linked mechanistically by haematogenously disseminated renal TB 	
In patients with disseminated TB at autopsy, renal TB is common	<ul style="list-style-type: none"> Prevalence of renal TB is similar to the sensitivity of LAM-positive urine in this population 	
Post-mortem renal TB in HIV-positive patients is associated with LAM-positive urine	<ul style="list-style-type: none"> A Ugandan Autopsy study revealed frequent renal TB in those whose urine also tested LAM-positive and all LAM-positive patients had haematogenously disseminated TB No patients with renal TB were urine LAM-negative 	

of deceased patients with LAM detected in their urine.¹⁷ All LAM-positive patients had evidence of either renal or disseminated TB. Renal TB is also frequently found in cadavers of adults with disseminated TB at post-mortem, typically being seen in more than 50% of such cases.^{24–27}

That LAM-antigenuria is due to renal involvement with haematogenously disseminated TB rather than filtration of LAM also explains several observations about urine-LAM assays, with implications for future research.

What are the implications?

Thus far, we have presented evidence supporting the notion that LAM-antigenuria results from renal involvement with TB arising from haematogenous TB dissemination and not from simple renal filtration of circulating LAM. This understanding is critical as it provides many important insights into the urine-LAM assays and their particular niche within the TB diagnostics armamentarium. Here we now discuss the implications of this knowledge (Tables 2 and 3).

Renal TB explains why urine-LAM assays only have utility in HIV-infected patients

Diagnostic accuracy studies have found the utility of urine-LAM assays to be almost entirely restricted to HIV-positive adults with advanced immunodeficiency.⁷ Disseminated TB, and

therefore renal TB, are comparatively much less common among immune-competent patients. Furthermore, assay sensitivity is very strongly associated with low CD4 cell counts, being highest among those with CD4 counts <50 cells/ μ L,^{7,32} the very patient group with highest risks of mycobacteraemia and disseminated and renal TB. Thus, understanding the mechanism of LAM-antigenuria directly explains why urine LAM-assays have utility restricted to a defined patient group and why the earlier diagnostic accuracy studies focusing on other patient groups generated largely unfavourable data. The results of studies of urine-LAM assays were also disappointing in children with no useful diagnostic accuracy observed, which is likely to be explained by the paucibacillary nature of paediatric TB, contrasting with the overwhelming multi-bacillary nature of disseminated TB in adults with advanced HIV.^{4,33}

Higher mortality in urine-LAM positive patients

Several studies have now demonstrated that HIV-TB co-infected patients who test urine-LAM-positive have much poorer prognostic characteristics and higher mortality risk compared to cases testing LAM-negative.^{28,34–36} This increased mortality risk persists even after adjustment for important risk factors such as CD4 cell count and anaemia.^{35,37,38} The fact that urine LAM-positivity serves as a marker of disseminated/renal TB, is an entirely plausible reason for this association with prognosis, as HIV-TB patients with mycobacteraemia are known to have high mortality risk.^{39,40} Diagnosing mycobacteraemia by liquid culture of blood samples can take 2 to 6 weeks, and is largely unavailable outside research

Table 2. Observations about urine lipoarabinomannan (LAM) assays explained by LAM-antigenuria resulting from renal involvement with TB

Observation	Evidence	References
The utility of the assay is almost entirely restricted to testing adult patients who are HIV-infected.	<ul style="list-style-type: none"> • Haematogenously disseminated TB, and therefore renal TB, is rare amongst immune-competent patients • If LAM were freely filtered sensitivity amongst pulmonary TB cases should be higher 	Minion et al. 2011 ⁷
Assay sensitivity is very strongly associated with blood CD4 cell count, (highest among CD4 counts <50 cells/ μ L and lowest among CD4 counts >200 cells/ μ L). The assay has no utility among children	<ul style="list-style-type: none"> • Haematogenously disseminated TB, and therefore renal TB, is more common with more advanced immunosuppression in HIV • Paediatric TB is usually paucibacillary with rare dissemination of TB disease, even in the context of HIV • Diagnostic yield of LAM is unrelated to respiratory symptoms (cough or sputum production) 	Lawn et al. 2009 ⁸ Lawn et al. 2012 ⁹ Minion et al. 2011 ⁷ Nicol M et al. 2014 ³³
Urine LAM assays should not be specifically targeted to the investigation of HIV-positive patients with respiratory symptoms of TB		Lawn et al. 2014 ²³
LAM-antigenuria is a strong independent predictor of mortality among patients with HIV-associated TB.	<ul style="list-style-type: none"> • Renal and haematogenously disseminated TB is associated with greater mycobacterial burden, which is likely to adversely impact prognosis 	Lawn et al. 2014 ³⁵ Talbot et al. 2012 ³⁷
Studies of urine-LAM find much higher specificity when the reference standard for TB diagnosis includes testing of extra-pulmonary samples in addition to sputum	<p>Testing extra-pulmonary samples increases the chances that disseminated TB is correctly detected, and fewer urine-LAM positive patients will be misclassified as 'not TB' due to inadequate reference standards.</p>	Drain et al. 2015 ³⁸ Cox et al. 2015 ¹⁷ Lawn et al. 2014 ³⁵ Nakiyingi et al. 2014 ³⁰

Table 3. Implications of lipoarabinomannan (LAM) antigenuria being a consequence of disseminated renal

Issue	Implications	References
Targeting the assay to appropriate patient groups	<ul style="list-style-type: none"> • LAM assays should be prioritised for testing sick HIV-positive adults, such as those requiring acute hospital admission rather than ambulatory out-patients • Interpretation of urine-LAM results requires knowledge of HIV status • There are no data showing utility for testing in children 	Nakiyingi et al. 2014 ³⁰
Design of diagnostic accuracy studies	<ul style="list-style-type: none"> • The reference standard for disseminated TB requires testing more than one specimen type, a single sputum culture is inadequate • Ideally this would include mycobacterial blood cultures and/or urine Xpert MTB/RIF assay testing • Should include patients both with and without respiratory symptoms 	Lawn et al. 2015 ⁴¹
Design of studies to assess incremental diagnostic yield of urine-LAM testing		Lawn et al. 2014 ⁴⁶
Studies of clinical impact of Urine-LAM testing		Gupta et al. 2015 ⁴ Lawn & Kerkhoff 2015 ³¹
Patients testing LAM-positive have poorer prognosis than those testing LAM-negative	<ul style="list-style-type: none"> • Studies should focus on acute HIV-positive medical admissions in whom undiagnosed TB and mycobacteraemia is common. e.g. LAMRCT and STAMP trials^{42,43} • Patients testing urine-LAM-positive are much sicker and have poorer prognostic characteristics compared with those with TB who testing LAM-negative 	Manabe et al. 2014 ²⁸ Lawn et al. 2013 ³⁶

facilities in most high burden settings. However, urine-LAM testing can quickly (a 30 min test) identify 80% of patients with mycobacteraemia,³¹ therefore expediting the diagnosis of MTB bacteraemia among patients with high mortality risk. Further research is required to see if interventions in addition to expediting TB treatment can reduce this very high mortality risk.

Underestimation of urine-LAM specificity due to an inadequate reference standard for disseminated TB in diagnostic accuracy studies

Finally, understanding that urine LAM-positivity represents disseminated and renal TB helps to explain why assessments of Determine

TB-LAM specificity are sub-optimal in many studies in which reference standard testing for TB was restricted to a single respiratory sample.⁴¹ LAM-positive disseminated and renal TB may be very easily missed by such a reference standard, especially among very sick patients unable to produce good quality sputum samples and would, therefore, be inappropriately misclassified as 'TB-free'. Specificity of urine LAM-detection is $\geq 99\%$ when the reference standard includes comprehensive extra-pulmonary sampling that is much more likely to detect disseminated TB and thereby accurately classifies patients as having TB or being TB-free.^{17,23,41}

Conclusion

The advent of a simple ELISA-based format for urine-LAM detection for TB diagnosis was greeted with much optimism in 2005, and yet many were perplexed by the results of subsequent studies, which were designed based on a limited understanding of these assays, including the mechanism of LAM-antigenuria, which has profound implications. In effect, the advent of urine-LAM assays heralded the development of simple assays for rapid diagnosis of disseminated HIV-associated TB with renal involvement, providing not only rapid diagnosis but also key prognostic information. This emerging understanding has now culminated in randomised trials of urine-based diagnosis to determine whether their addition to routine diagnostic screening algorithms averts deaths from HIV-associated TB by improving and accelerating the TB diagnostic yield.^{42,43} Results of such studies are awaited.

This paper highlights the critical importance that new diagnostic assays are thoroughly understood mechanistically and are applied to the most appropriate patient groups. Without this, there is a risk that some of the output of the TB diagnostics developmental pipeline might potentially be squandered. However, the mechanistic framework for understanding the utility of the urine LAM assays that we have described in this article is strongly supportive of the WHO recommendations regarding the use of these assays for diagnosis. These specifically restrict or target the use in HIV-positive patients who have the highest risk of disseminated TB (notably in-patients who are sick and have CD4 cell counts <100 cells/ μL).⁴⁴

Authors' contributions: Both SDL AND AGW contributed equally to the manuscript and conceived, designed, drafted, wrote and revised the manuscript. Both authors read and approved the final manuscript. SDL is the guarantor of this paper.

Funding: SDL and AGW are both funded by a Global Clinical Trials Grant from the MRC/DFID/Wellcome Trust (STAMP Trial; grant no. MR/M007375/1). AGW is also funded by the Royal College of Physicians, London, UK (JMGF fellowship).

Competing interests: None declared.

Ethical approval: None required.

References

- 1 United Nations. Transforming Our World: The 2030 Agenda for Sustainable Development. New York: United Nations; 2015.
- 2 WHO. Global Strategy and Targets for Tuberculosis Prevention, Care and Control after 2015. Geneva; World Health Organization; 2014.
- 3 WHO. Global Tuberculosis Report 2015. Geneva: World Health Organization; 2015.
- 4 Gupta RK, Lucas SB, Fielding KL et al. Prevalence of tuberculosis in post-mortem studies of HIV-infected adults and children in resource-limited settings: a systematic review and meta-analysis. *AIDS* 2015;29:1987–2002.
- 5 Hamasur B, Bruchfeld J, Haile M et al. Rapid diagnosis of tuberculosis by detection of mycobacterial lipoarabinomannan in urine. *J Microbiol Methods* 2001;45:41–52.
- 6 Boehme C, Molokova E, Minja F et al. Detection of mycobacterial lipoarabinomannan with an antigen-capture ELISA in unprocessed urine of Tanzanian patients with suspected tuberculosis. *Trans R Soc Trop Med Hyg* 2005;99:893–900.
- 7 Minion J, Leung E, Talbot E et al. Diagnosing tuberculosis with urine lipoarabinomannan: systematic review and meta-analysis. *Eur Respir J* 2011;38:1398–405.
- 8 Lawn SD, Edwards DJ, Kranzer K et al. Urine lipoarabinomannan assay for tuberculosis screening before antiretroviral therapy diagnostic yield and association with immune reconstitution disease. *AIDS* 2009;23:1875–80.
- 9 Lawn SD, Kerkhoff AD, Vogt M et al. Diagnostic accuracy of a low-cost, urine antigen, point-of-care screening assay for HIV-associated pulmonary tuberculosis before antiretroviral therapy: a descriptive study. *Lancet Infect Dis* 2012;12:201–9.
- 10 Peter J, Green C, Hoelscher M et al. Urine for the diagnosis of tuberculosis: current approaches, clinical applicability, and new developments. *Curr Opin Pulm Med* 2010;16:262–70.
- 11 Strohmeier GR, Fenton MJ. Roles of lipoarabinomannan in the pathogenesis of tuberculosis. *Microbes Infect* 1999;1:709–17.
- 12 Sada E, Aguilar D, Torres M et al. Detection of lipoarabinomannan as a diagnostic test for tuberculosis. *J Clin Microbiol* 1992;30:2415–8.
- 13 Sakamuri RM, Price DN, Lee M et al. Association of lipoarabinomannan with high density lipoprotein in blood: implications for diagnostics. *Tuberculosis* 2013;93:301–7.
- 14 Haraldsson B, Nyström J, Deen WM. Properties of the glomerular barrier and mechanisms of proteinuria. *Physiol Rev* 2008;88:451–87.
- 15 Wood R, Racow K, Bekker L-G et al. Lipoarabinomannan in urine during tuberculosis treatment: association with host and pathogen factors and mycobacteruria. *BMC Infect Dis* 2012;12:47.
- 16 Peter JG, Theron G, van Zyl-Smit R et al. Diagnostic accuracy of a urine lipoarabinomannan strip-test for TB detection in HIV-infected hospitalised patients. *Eur Respir J* 2012;40:1211–20.
- 17 Cox JA, Lukande RL, Kalungi S et al. Is urinary lipoarabinomannan the result of renal Tuberculosis? Assessment of the renal histology in an autopsy cohort of Ugandan HIV-infected adults. *PLoS One* 2015;10:e0123323.
- 18 Shafer RW, Kim DS, Weiss JP et al. Extrapulmonary tuberculosis in patients with human immunodeficiency virus infection. *Medicine (Baltimore)* 1991;70:384–97.
- 19 Blakemore R, Story E, Helb D et al. Evaluation of the analytical performance of the Xpert MTB/RIF assay. *J Clin Microbiol* 2010;48:2495–501.
- 20 Lawn SD, Nicol MP. Xpert® MTB/RIF assay: development, evaluation and implementation of a new rapid molecular diagnostic for tuberculosis and rifampicin resistance. *Future Microbiol* 2011;6:1067–82.
- 21 Lawn SD, Kerkhoff AD, Vogt M et al. High diagnostic yield of tuberculosis from screening urine samples from HIV-infected

- patients with advanced immunodeficiency using the Xpert MTB/RIF assay. *J Acquir Immune Defic Syndr* 2012;60:289–94.
- 22 Peter JG, Theron G, Muchinga TE et al. The diagnostic accuracy of urine-based Xpert MTB/RIF in HIV-infected hospitalized patients who are smear-negative or sputum scarce. *PLoS One* 2012;7:e39966.
 - 23 Lawn SD, Kerkhoff AD, Burton R et al. Systematic investigation for tuberculosis in HIV-infected patients on the first day of admission to a South African hospital: incremental diagnostic yield, accuracy and prognostic value of a urine LAM lateral-flow assay. 45th World Conference on Lung Health of the International Union Against Tuberculosis and Lung Disease (The Union). Barcelona, Spain, 2014, HIV Late Breaker Oral Presentation.
 - 24 Ansari NA, Kombe AH, Kenyon TA et al. Pathology and causes of death in a group of 128 predominantly HIV-positive patients in Botswana, 1997–1998. *Int J Tuberc Lung Dis* 2002;6:55–63.
 - 25 Lanjewar DN. The spectrum of clinical and pathological manifestations of AIDS in a consecutive series of 236 autopsied cases in mumbai, India. *Patholog Res Int* 2011;2011:547618.
 - 26 Eza D, Cerrillo G, Moore DAJ et al. Postmortem findings and opportunistic infections in HIV-positive patients from a public hospital in Peru. *Pathol Res Pract* 2006;202:767–75.
 - 27 Marques LP, Rioja LS, Oliveira CA et al. AIDS-associated renal tuberculosis. *Nephron* 1996;74:701–4.
 - 28 Manabe YC, Nonyane BA, Nakiyingi L et al. Point-of-Care lateral flow assays for Tuberculosis and Cryptococcal antigenuria predict death in HIV infected adults in Uganda. *PLoS One* 2014;9:e101459.
 - 29 Nakiyingi L, Ssengooba W, Nakanjako D et al. Predictors and outcomes of mycobacteremia among HIV-infected smear- negative presumptive tuberculosis patients in Uganda. *BMC Infect Dis* 2015;15:1–8.
 - 30 Nakiyingi L, Moodley VM, Manabe YC et al. Diagnostic accuracy of a rapid urine lipoarabinomannan test for tuberculosis in HIV-infected adults. *J Acquir immune Defic Syndr* 2014;66:270–9.
 - 31 Lawn SD, Kerkhoff AD. Rapid diagnosis of TB in HIV-positive in-patients with *M. tuberculosis* bacteraemia in sub-Saharan Africa. *Int J Tuberc Lung Dis* 2015;19:1557–9.
 - 32 Lawn SD. Point-of-care detection of lipoarabinomannan (LAM) in urine for diagnosis of HIV-associated tuberculosis: a state of the art review. *BMC Infect Dis* 2012;12:103.
 - 33 Nicol MP, Allen V, Workman L et al. Urine lipoarabinomannan testing for diagnosis of pulmonary tuberculosis in children: a prospective study. *Lancet Glob Heal* 2014;2:e278–84.
 - 34 Kerkhoff AD, Wood R, Vogt M et al. Prognostic value of a quantitative analysis of lipoarabinomannan in urine from patients with HIV-associated Tuberculosis. *PLoS One* 2014;9:e103285.
 - 35 Lawn SD, Kerkhoff AD, Burton R et al. Massive diagnostic yield of HIV-associated Tuberculosis using rapid urine diagnostic assays in South Africa. Program and Abstracts of the Conferences on Retroviruses and Opportunistic Infections (CROI). 2014, Abstract 2813.
 - 36 Lawn SD, Kerkhoff AD, Vogt M et al. HIV-associated tuberculosis: relationship between disease severity and the sensitivity of new sputum-based and urine-based diagnostic assays. *BMC Med* 2013;11:231.
 - 37 Talbot E, Munseri P, Teixeira P et al. Test characteristics of urinary lipoarabinomannan and predictors of mortality among hospitalized HIV-infected tuberculosis suspects in Tanzania. *PLoS One* 2012;7:e32876.
 - 38 Drain PK, Gounder L, Grobler A et al. Urine lipoarabinomannan to monitor antituberculosis therapy response and predict mortality in an HIV-endemic region: a prospective cohort study. *BMJ Open* 2015;5:e006833–e006833.
 - 39 Feasey NA, Banada PP, Howson W et al. Evaluation of Xpert MTB/RIF for detection of tuberculosis from blood samples of HIV-infected adults confirms Mycobacterium tuberculosis bacteraemia as an indicator of poor prognosis. *J Clin Microbiol* 2013;51:2311–6.
 - 40 Cummings MJ, Donnell MRO. Inverting the pyramid : increasing awareness of mycobacterial sepsis in sub-Saharan Africa. *Int J Tuberc Lung Dis* 2015;19:1128–34.
 - 41 Lawn SD, Kerkhoff AD, Nicol MP et al. Underestimation of the true specificity of the urine lipoarabinomannan point-of-care diagnostic assay for HIV-associated Tuberculosis. *J Acquir Immune Defic Syndr* 2015;69:e144–6.
 - 42 Peter J. A Trial of the Urine LAM Strip Test for TB Diagnosis Amongst Hospitalized HIV-infected Patients (LAMRCT) NCT01770730. <https://clinicaltrials.gov/ct2/show/NCT01770730> [accessed 5 December 2015].
 - 43 Lawn SD. Rapid urine-based Screening for Tuberculosis to reduce AIDS-related Mortality in hospitalized Patients in Africa (STAMP) trial. <http://www.isrctn.com/ISRCTN71603869> [accessed 10 January 2016].
 - 44 WHO. The Use of Lateral Flow Urine Lipoarabinomannan Assay (LF-LAM) for the Diagnosis and Screening of Active Tuberculosis in People Living with HIV: Policy. Geneva: World Health Organization; 2015.
 - 45 Bekker LG, Maartens G, Steyn L et al. Selective increase in plasma tumor necrosis factor-alpha and concomitant clinical deterioration after initiating therapy in patients with severe tuberculosis. *J Infect Dis* 1998;178:580–4.
 - 46 Lawn SD, Kerkhoff AD, Burton R et al. Underestimation of the incremental diagnostic yield of HIV-associated tuberculosis in studies of the determine TB-LAM Ag urine assay. *AIDS* 2014;28:1846–8.