

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Johnston, LG; Hakim, AJ; Dittrich, S; Burnett, J; Kim, E; White, RG; (2016) A Systematic Review of Published Respondent-Driven Sampling Surveys Collecting Behavioral and Biologic Data. *AIDS and behavior*, 20 (8). pp. 1754-76. ISSN 1090-7165 DOI: <https://doi.org/10.1007/s10461-016-1346-5>

Downloaded from: <http://researchonline.lshtm.ac.uk/2535624/>

DOI: <https://doi.org/10.1007/s10461-016-1346-5>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

A Systematic Review of Published Respondent-Driven Sampling Surveys

Collecting Behavioral and Biologic Data

Authors here based on contribution

Lisa G Johnston^{1,2}, Avi J Hakim³, Samantha Dittrich³, Janet Burnett³, Evelyn Kim³, Richard G White⁴

Author's Affiliations:

1. University of California, San Francisco, Global Health Sciences, San Francisco, California, USA,
2. Tulane University, School of Public Health and Tropical Medicine, New Orleans, Louisiana, USA
3. US Centers for Disease Control and Prevention, Division of Global HIV/AIDS, Atlanta, USA
4. Faculty of Epidemiology & Population Health, London School of Hygiene and Tropical Medicine, London, United Kingdom

Address correspondence to: Lisa G. Johnston, 64F Nieuwezijds Voorburgwal, Amsterdam, Netherlands.

Disclaimer: The findings and conclusions in this paper are those of the authors and do not necessarily represent the official position of the US Centers for Disease Control and Prevention.

Funding: This project has been supported in part by the President's Emergency Plan for AIDS Relief (PEPFAR) through the Centers for Disease Control and Prevention (CDC).

Acknowledgments: We would like to thank Kate Orroth for conducting the literature search for the STROBE-RDS Guidelines and allowing us to use it for this analysis.

Abstract word count: 160

Word count: 3060

ABSTRACT

Reporting key details of respondent-driven sampling (RDS) survey implementation and analysis is essential for assessing the quality of RDS surveys. RDS is both a recruitment and analytic method and, as such, it is important to adequately describe both aspects in publications. We extracted data from peer-reviewed literature published through September, 2013 that reported collected biological specimens using RDS. We identified 151 eligible peer-reviewed articles describing 222 surveys conducted in seven regions throughout the world. Most published surveys reported basic implementation information such as survey city, country, year, population sampled, interview method, and final sample size. However, many surveys did not report essential methodological and analytical information for assessing RDS survey quality, including number of recruitment sites, seeds at start and end, maximum number of waves, and whether data were adjusted for network size. Understanding the quality of data collection and analysis in RDS is useful to effectively plan for public health service delivery and funding priorities.

INTRODUCTION

The first respondent-driven sampling (RDS) surveys to assess HIV prevalence in addition to risk behaviors were conducted in 2004 (1–3). Since then hundreds of surveys have been conducted worldwide to capture data from populations considered at higher risk for HIV exposure, including people who inject drugs (PWID), men who have sex with men (MSM), female sex workers (FSW), and other populations without a readily available sampling frame (4–6). Over the past decade, many organizations, including the U.S. Centers for Disease Control and Prevention, UNAIDS, WHO, Global Fund and others have endorsed the use of RDS to establish baseline and trend measurements of HIV and other infections (Syphilis, Hepatitis, etc.) prevalence, risk behaviors, and program impact through biological and behavioral surveys (6–9).

RDS is an important recruitment and analysis tools for sampling populations with no sampling frame that are linked through social networks, including those considered “hard-to-reach” due to stigma and the practice of illegal behaviors (4–6). RDS, a chain-referral sampling technique, uses statistical adjustments for network sizes and recruitment effort to produce estimates representative of a population’s network. Beginning with a set number of participants, “seeds”, selected purposefully by the research team from the target population, RDS builds a sample based on the recruitment of individuals with pre-existing relationships. Using a limited number of coupons for each participant limits the possibility of overrepresentation of those with more ties to others in the population. Providing ‘incentives’ for those participating in and for recruiting peers into the survey helps ensure ongoing participation and recruitment. Ideally, this process results in long recruitment chains made up of numerous “waves” of

recruits (10,14). As recruitment chains lengthen, the structure of the sample becomes less dependent on the purposefully selected seeds and more increasingly similar to the population being sampled.

In addition to these implementation steps, RDS is premised upon several assumptions, most importantly, random walk models (10). Briefly, these assumptions include 1) reciprocal ties between respondents (i.e., know one another as members of the sampled population); 2) respondents are connected by a single network component; 3) sampling occurs with replacement; 4) respondents provide accurate personal network sizes (i.e., number of relatives, friends, and acquaintances they know from the sampled population); 5) peers are recruited randomly from the recruiter's network; and, 6) each respondent can recruit at least one peer (11). RDS also involves a complex analytical component crucial for generating population estimates and confidence intervals using data collected about each participant's social network size (10–14). Estimates generated through RDS should allow for inferences about the network of the population being sampled.

Findings from surveys purporting to use RDS are vital for developing national and international policies, guiding service delivery, informing budgets and dictating funding priorities. Quality reporting of data collected and analyzed using RDS methods allows users to assess their usefulness in decision making. However, there is ample potential for bias when using this method, many of which are related to implementation and analytical failures (15–20). The allure of RDS as a more robust alternative to convenience snowball sampling methods has resulted in partial incorporation of RDS techniques (i.e., the use of coupons) while ignoring some of the more complex aspects

which ensure the mitigation of chain referral related biases (13). Indeed, numerous published surveys report having used RDS, but present insufficient methodological and analytical information to support this assertion (21).

Building upon the STROBE RDS guidelines (22) to recommend improvements in the reporting of survey data, we extracted peer-reviewed literature that reported using RDS for collecting HIV and other infections biological and behavioral data through September, 2013. Specifically, we evaluate a set of general and RDS-specific survey indicators based on the STROBE RDS guidelines (21,22) to describe the extent, consistency, and changes over time for planning, implementation, and analysis as reported in peer reviewed journals. In addition, we provide reasons why some published surveys were not included in the extraction and examples of surveys that reported using RDS when, in fact, evidence suggests they did not. We hope to build upon other efforts to increase accuracy in conducting RDS and to encourage more thorough and standardized reporting of RDS methods and analysis (4,22).

Location of study, citation	Year of study	Population	Pre-survey assessment	Sites, #	Interview method†	Seeds at start, #	Final seeds, #	Primary incentive of value‡	Secondary incentive of value‡	Target sample size
Africa										
Kenya, Kisumu(23)	2008	FSW	Yes	1	ACASI	15	NR	4.00	1.25	480
Nigeria, Abuja(24)	2010	MSM	NR	NR	ACASI/SA	NR	NR	NR	NR	NR
Nigeria, Cross River(25)	2007	MSM	NR	1	IA	10	10	4.00	NR	293
Nigeria, Cross River(26)	2010	PWID	NR	>1	IA	10	10	4.00	4.00	266
Nigeria, Federal Capital Territory(26)	2010	PWID	NR	>1	IA	10	10	4.00	4.00	266
Nigeria, Ibadan(27)	2006	MSM	NR	NR	IA	38	38	4.00	NR	NR
Nigeria, Ibadan(24)	2010	MSM	NR	NR	SA ACASI/SA	NR	NR	NR	NR	NR
Nigeria, Kaduna(26)	2010	PWID	NR	>1	IA	10	10	4.00	4.00	266
Nigeria, Kano(26)	2010	PWID	NR	>1	IA	10	10	4.00	4.00	266
Nigeria, Kano(25)	2007	MSM	NR	1	IA	10	10	4.00	NR	293
Nigeria, Lagos(25)	2007	MSM	NR	1	IA	10	14	4.00	NR	293
Nigeria, Lagos(26)	2010	PWID	NR	>1	IA	10	10	4.00	4.00	266
Nigeria, Lagos(27)	2006	MSM	NR	NR	IA	38	38	4.00	NR	NR
Nigeria, Lagos(24)	2010	MSM	NR	NR	SA ACASI/SA	NR	NR	NR	NR	NR
Nigeria, Oyo(26)	2010	PWID	NR	>1	IA	10	10	4.00	4.00	266
Mauritius(28)	2009	PWID	Yes	2	IA	6	6	7.00	3.50	500
Mauritius(29)	2010	FSW	NR	2	IA	5	5	17.50	7.00	NR

Somalia, Hargeisa, Somaliland(30)	2008	FSW	Yes	1	IA w/ HAPI	6	NR	4.00	3.00	146	2
South Africa, Durban(31)	2011	MSM	Yes	1	SA	4	15	5.00 & 5.00 voucher	5.00 & 5.00 voucher	200	8
South Africa, Johannesburg(31)	2011	MSM	Yes	1	IA or SA	5	14	5.00 & 5.00 voucher	5.00 & 5.00 voucher	200	2
South Africa, Soweto(32)	2008	MSM	NR	1	IA	15	15	NR	NR	NR	3
South Africa, W. Cape Province(33-35)	2006	Heterosexual men	Yes	1	IA	8	20	8.00 phone voucher	2.70 phone voucher	430	4
South Africa, W. Cape Province(36)	2008	Heterosexual men	Yes	1	IA	19	19	8.50 phone voucher	2.85 phone voucher	430	4
South Africa, W. Cape Province(37)	2007	Young women	Yes	1	SA	5	5	8.00 make-up voucher	2.50	270	2
South Africa, W. Cape Province(38)	2011	Heterosexual women	Yes	1	ACASI	15	15	7.50 grocery voucher	7.50 grocery voucher	756	8
Sudan, Khartoum(39)	2008	FSW	NR	1	IA	NR	NR	10.00	10.00	NR	3
Tanzania, Zanzibar(40,41)	2007	MSM	Yes	1	IA	10	10	3.00	1.50	500	5
Uganda, Kampala(42)	2008-09	MSM	Yes	1	ACASI	8	14	3.00	1.00	600	3
Eastern Mediterranean											
Egypt, Cairo(43)	2006	IDU	Yes	1	IA	28	NR	7.00	5.30	406	4
Iran, Kerman(44)	2010	FSW	NR	1	IA	8	12	4.00	2.00	NR	1
Lebanon, Beirut(45,46)	2007-08	FSW	NR	NR	NR	NR	NR	6.60	2.00	NR	1
Lebanon, Beirut(45)	2007-08	MSM	NR	NR	NR	NR	NR	6.60	2.00	NR	1
Lebanon, Beirut(45,47)	2007-08	PWID	NR	NR	NR	NR	NR	6.60	2.00	NR	8
Libya, Tripoli(48)	2010	PWID	Yes	1	IA	7	7	20.00	9.00	NR	3
Libya, Tripoli(49)	2010	MSM	Yes	1	IA	NR	14	NR	NR	NR	2
Libya, Tripoli(49)	2010	FSW	Yes	1	IA	NR	13	NR	NR	314	6
Morocco, Agadir(50)	2010-11	MSM	NR	NR	IA	NR	10	7.00	3.50	NR	3
Morocco, Marrakesh(50)	2010-11	MSM	NR	NR	SA	NR	8	7.00	3.50	NR	3
Palestine, East Jerusalem(51)	2010	PWID	Yes	1	IA	NR	7	NR	NR	NR	1
Europe											
Albania, Tirana(52)	2005	PWID	Yes	3	IA	15	15	12.00	7.00	NR	2
Albania, Tirana(53)	2008	MSM	NR	1	IA	12	NR	10.00	5.00	NR	1
Croatia, Zagreb(54-56)	2006	MSM	NR	1	SA	8	10	18.00	9.00	400	3
Croatia, Zagreb(54)	2012	MSM	Yes	1	SA	10	15	None	9.60	370	4
England, Bristol(57,58)	2006	PWID	NR	NR	CASI	7	7	15.00	10.00	NR	2
England, Bristol(57)	2009	PWID	NR	NR	NR	6	6	NR	NR	NR	2
Estonia, Kohtla Jarve(59,60)	2005	PWID	NR	NR	SA	NR	NR	NR	NR	NR	1
Estonia, Tallinn(61-63)	2007	PWID	Yes	1	SA	5	5	10.00 food voucher	5.00 food voucher	NR	3
Estonia, Tallinn(63,64)	2005-06	FSW	Yes	1; other	SA	6	43	10.00 shop voucher	11.00 shop voucher	NR	2
Estonia, Tallinn(60,65)	2005	PWID	NR	NR	SA	6	NR	NR	NR	NR	3
Estonia, Tallinn(62)	2009	PWID	NR	NR	SA	6	6	10.00 food voucher	5.00 food voucher	NR	3
Kazakhstan, Almaty(66)	2010	MSM	Yes	NR	IA	4	4	10.00	2.50	400	4
Moldova, Balti(67)	2009-10	FSW	NR	NR	IA	5	5	7.00	5.00	350	3
Moldova, Balti(68)	2007-08	PWID	Yes	NR	IA	NR	NR	Items, cash (value NR)	NR	NR	3
Moldova, Balti(69)	2010	MSM	Yes	NR	IA	5	5	8.30	5.80	250	2
Moldova, Chisinau(68)	2007-08	PWID	Yes	NR	IA	NR	NR	Items, cash (value NR)	NR	NR	3
Moldova, Chisinau(67)	2009-10	FSW	NR	NR	IA	5	5	16.00	12.00	350	2
Moldova, Chisinau(69)	2010	MSM	Yes	NR	IA	8	8	8.30	5.80	250	1
Moldova, Tiraspol(68)	2007-08	PWID	Yes	NR	IA	NR	NR	Items, cash (value NR)	NR	NR	3
Montenegro, Podgorica(70)	2008	PWID	NR	1	SA	5	NR	20.80	7.00	NR	3

Montenegro, Podgorica(71)	2005	PWID	NR	1	ACASI	NR	NR	13.00	6.00	NR	3
Russia, Ivanovo(72)	2010	PWID	NR	NR	IA	11	11	Items, food (value NR)	Items, food (value NR)	NR	3
Russia, Novosibirsk(72)	2010	PWID	NR	NR	IA	10	10	Items, food (value NR)	Item, food (value NR)	NR	2
Russia, St. Petersburg(73–75)	2005-08	PWUD/PWID	NR	NR	CASI	48; 108	156	10.00 (items)	items (value NR)	NR	6
Russia, St. Petersburg(74,75)	2005-06	PWID	NR	NR	CASI	35	NR	10.00 (items)	NR	NR	3
Serbia, Belgrade(76)	2010	Youth	Yes	NR	IA	8	8	13.00	6.00	371 ¶	2
Serbia, Belgrade(71)	2005	PWID	NR	1	ACASI	NR	NR	13.00	6.00	NR	4
Serbia, Kragujevac(76)	2010	Youth	Yes	NR	IA	4	4	13.00	6.00	370 ¶	1
Ukraine, Poltava(77)	2011	PWID	NR	NR	SA	4	4	3.00	2.00	NR	2
Ukraine, Khmel'nitsky(77)	2011	PWID	NR	NR	SA	7	7	3.00	2.00	NR	2
Ukraine, Dnipropetrovsk(77)	2011	PWID	NR	NR	SA	6	6	3.00	2.00	NR	1
Ukraine, Cherkasy(77)	2011	PWID	NR	NR	SA	3	3	3.00	2.00	NR	1
Ukraine, Donetsk(77)	2011	PWID	NR	NR	SA	6	6	3.00	2.00	NR	4
Ukraine, Kharkov(77)	2011	PWID	NR	NR	SA	5	5	3.00	2.00	NR	1
Ukraine, Kherson(77)	2011	PWID	NR	NR	SA	4	4	3.00	2.00	NR	2
Ukraine, Kirovograd(77)	2011	PWID	NR	NR	SA	4	4	3.00	2.00	NR	1
Ukraine, Kyiv(77)	2011	PWID	NR	NR	SA	8	8	3.00	2.00	NR	4
Ukraine, Lugansk(77)	2011	PWID	NR	NR	SA	6	6	3.00	2.00	NR	2
Ukraine, Lutsk(77)	2011	PWID	NR	NR	SA	4	4	3.00	2.00	NR	1
Ukraine, Lviv(77)	2011	PWID	NR	NR	SA	7	7	3.00	2.00	NR	1
Ukraine, Mykolaiv(77)	2011	PWID	NR	NR	SA	6	6	3.00	2.00	NR	2
Ukraine, Odesa(77)	2011	PWID	NR	NR	SA	6	6	3.00	2.00	NR	4
Ukraine, Simferopol(77)	2011	PWID	NR	NR	SA	5	5	3.00	2.00	NR	2
Ukraine, Sumy(77)	2011	PWID	NR	NR	SA	5	5	3.00	2.00	NR	1
Latin America and Caribbean											
Argentina, Buenos Aires(78,79)	2009	MSM	NR	NR	SA web-based	16	16	NR	NR	NR	5
Brazil, Belo Horizonte(80–82)	2009	MSM	Yes	NR	IA	6	NR	10.00	6.67	350	M
Brazil, Belo Horizonte(17,83)	2008-09	FSW	NR	NR	ACASI	5 to 10	NR	Misc. (value NR)	4.00	300	2
Brazil, Brasilia(80–82)	2009	MSM	Yes	NR	IA	6	NR	10.00	6.67	350	M
Brazil, Brasilia(17,83)	2008-09	FSW	NR	NR	ACASI	5 to 10	NR	Misc. (value NR)	4.00	300	3
Brazil, Campinas(84)	2005-06	MSM	NR	NR	ACASI	10	30	NR	NR	NR	6
Brazil, Campo Grande(80–82)	2009	MSM	Yes	NR	IA	6	NR	10.00	6.67	350	M
Brazil, Campo Grande(17,83)	2008-09	FSW	NR	NR	ACASI	5 to 10	NR	Misc. (value NR)	4.00	150	1
Brazil, Curitiba(80–82)	2009	MSM	Yes	NR	IA	6	NR	10.00	6.67	350	M
Brazil, Curitiba(17,83)	2008-09	FSW	NR	NR	ACASI	5 to 10	NR	Misc. (value NR)	4.00	200	2
Brazil, Fortaleza(85)	2008	Transvestite	Yes	NR	IA	6	NR	6.00 food voucher	3.00	NR	3
Brazil, Fortaleza(86)	2005	MSM	Yes	2	IA	10	10	5.00	5.00	400	4
Brazil, Itajaí(80–82)	2009	MSM	Yes	NR	IA	6	NR	10.00	6.67	350	M
Brazil, Itajaí(17,83)	2008-09	FSW	NR	NR	ACASI	5 to 10	NR	Misc. (value NR)	4.00	100	9
Brazil, Manaus(80–82)	2009	MSM	Yes	NR	IA	6	NR	10.00	6.67	350	M
Brazil, Manaus(17,83)	2008-09	FSW	NR	NR	ACASI	5 to 10	NR	Misc. (value NR)	4.00	200	1
Brazil, Recife(80–82)	2009	MSM	Yes	NR	IA	6	NR	10.00	6.67	350	M
Brazil, Recife(17,83)	2008-09	FSW	NR	NR	ACASI	5 to 10	NR	Misc. (value NR)	4.00	200	2
Brazil, Rio de Janeiro(80–82)	2009	MSM	Yes	NR	IA	6	NR	10.00	6.67	350	M
Brazil, Rio de Janeiro(17,83)	2008-09	FSW	NR	NR	ACASI	5 to 10	NR	Misc. (value NR)	4.00	600	6
Brazil, Salvador(80–82)	2009	MSM	Yes	NR	IA	6	NR	10.00	6.67	350	M
Brazil, Salvador(17,83)	2008-09	FSW	NR	NR	ACASI	5 to 10	NR	Misc. (value NR)	4.00	300	2
Brazil, Santos(80–82)	2009	MSM	Yes	NR	IA	6	NR	10.00	6.67	350	M
Brazil, Santos(17,83)	2008-09	FSW	NR	NR	ACASI	5 to 10	NR	Misc. (value NR)	4.00	150	1
Dominican Rep., Barahona(87)	2008	MSM	Yes	NR	IA	8	NR	9.00	3.00	300	2

Dominican Rep., La Altagracia(87)	2008	MSM	Yes	NR	IA	7	NR	9.00	3.00	300	2
Dominican Rep., Santiago(87)	2008	MSM	Yes	NR	IA	6	NR	9.00	3.00	300	3
Dominican Rep., Santo Domingo(87)	2008	MSM	Yes	NR	IA	7	NR	9.00	3.00	500	5
El Salvador, San Miguel(88,89)	2008	MSM	Yes	NR	CASI w/ interviewer	5	5	4.00 as items	2.70 (items)	200	1
El Salvador, San Salvador(89)	2008	FSW	NR	1	CASI w/ interviewer	10	10	Items (value NR)	Items (value NR)	NR	7
El Salvador, San Salvador(88,89)	2008	MSM	Yes	NR	CASI w/ interviewer	11	11	4.00 as items	2.70 (items)	600	5
El Salvador, Sonsonate(89)	2008	FSW	NR	1	CASI w/ interviewer	5	5	Items (value NR)	Items (value NR)	NR	7
Honduras, Comayagua(90)	2006	FSW	Yes	1	ACASI	5	5	Purse (value <2.00)	Items (value 3.50)	200	1
Honduras, La Ceiba(90)	2006	FSW	Yes	1	ACASI	7	7	"	"	200	2
Honduras, San Pedro Sula(90)	2006	FSW	Yes	1	ACASI	7	7	"	"	200	1
Honduras, Tegucigalpa(90)	2006	FSW	Yes	1	ACASI	5	5	"	"	200	2
Peru, Lima(91)	2012	Transwoman	Yes	6	SA	8	11	7.00	NR	420	4
North America											
Mexico, Juarez(92-96)	2005	PWID	NR	1	SA	NR	NR	NR	NR	NR	2
Mexico, Juarez(97)	2005	PWID	NR	1	SA	9	17	20.00	5.00	200	1
Mexico, Tijuana(92-96)	2005	PWID	NR	3	SA	15	15	10.00	5.00	200	2
Mexico, Tijuana(98-102)	2006-07	PWID	NR	NR	SA	32	NR	20.00	NR	NR	1
USA, Appalachia(103)	NR	DU	NR	NR	NR	NR	NR	50.00	10.00	NR	5
USA, Atlanta(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, Baltimore(104)	2002-04	Youth IDU	NR	NR	ACASI, w/ w/out interviewer	NR	NR	NR	NR	NR	7
USA, Baltimore(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	7
USA, Baltimore(105)	2006	PWID	NR	NR	CAPI	20	20	20.00	10.00	NR	6
USA, Boston(106,107)	2008	MSM	NR	2	SA	8	21	50.00	10.00	NR	1
USA, Boston(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	4
USA, Chicago(104)	2002-04	Youth IDU	NR	NR	ACASI, w/ w/out interviewer	NR	NR	NR	NR	NR	5
USA, Chicago(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, Dallas(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, Denver(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, Detroit(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, Ft. Lauderdale(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	3
USA, Houston(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, Houston(108)	2006-07	High risk heterosexuals	Yes	1	CAPI	NR	NR	40.00	10.00	750	9
USA, Las Vegas(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	3
USA, Los Angeles(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	6
USA, Los Angeles(75,109-111)	2005-06	DU/PWID/MSM	NR	1	ACASI	25	25	50.00	20.00	NR	4
USA, Miami(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	6
USA, Nassau(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, New Haven(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, Las Cruces(97)	2005	PWID	NR	1	SA	NR	NR	NR	NR	NR	1
USA, New York City(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, New York City(112)	2009	PWID	Yes	NR	SA	NR	3	NR	NR	NR	4
USA, New York City(113-118)	2006-07	High risk heterosexuals	Yes	NR	SA	8	NR	30.00	11.00	NR	8

USA, New York City(1,119,120)	2004	DU	NR	NR	NR	NR	NR	20.00	10.00	NR	4
USA, Newark(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	4
USA, Norfolk(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	4
USA, Oakland(121)	2011-12	High risk/HIV pos. African American	Yes	4	SA	48	NR	10.00 gift card	Varied	NR	2
USA, Philadelphia(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, San Diego(122)	2009-10	PWID	NR	NR	ACASI	NR	NR	NR	10.00	NR	5
USA, San Diego(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, San Francisco(123)	2007-08	MSM	Yes	1	CAPI	10	10	40.00	10.00	NR	2
USA, San Francisco(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, San Juan(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, Seattle(124)	2009	PWID	NR	1	IA	6	6	40.00	10.00	NR	4
USA, Seattle(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	4
USA, St. Louis(6)	2005-06	PWID	NR	NR	SA	8 to 10	NR	25.00	10.00	NR	5
USA, Wash. DC(125)	2006-07	High risk heterosexuals	Yes	1	SA	NR	NR	35.00	10.00	NR	7
USA, Wash. DC(126)	2009	PWID	NR	1	SA	NR	NR	30.00	10.00	NR	5
USA, Texas, El Paso(97)	2006	PWID	NR	1	SA	NR	NR	NR	NR	NR	1
South East Asia											
Bangladesh, Dhaka(127)	2006	MSM	Yes	1	SA	5	8	2.14	1.43	530	5
India, Bishenpur District, Manipur(128)	2006	PWID	NR	NR	SA	NR	NR	NR	NR	400	4
India, Chennai(129)	2008	MSM	Yes	2	SA	NR	19	6.00	None	NR	7
India, Churachandpur District, Manipur(128)	2006	PWID	NR	NR	SA	NR	NR	NR	NR	400	4
India, Coimbatore(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Dimapur District, Nagaland(130)	2006	FSW	NR	NR	SA	10	10	NR	NR	400	4
India, Dindigul(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Erode(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Goa(131,132)	2005	FSW	Yes	NR	SA	59	59	2.50	1.50	318	3
India, Kanyakumari(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Madurai(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Mumbai & Thane Districts(128)	2006	PWID	NR	NR	SA	NR	NR	NR	NR	400	3
India, Nagapattinam(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Nilgiris(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Perambalur(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Phek District, Nagaland(128)	2006	PWID	NR	NR	SA	NR	NR	NR	NR	400	4
India, Pudukottai(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Ramanathapuram(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Salem(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Sivagangai(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Thanjavur(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Tiruchy(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Tirunelveli(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Tiruvarur(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Tuticorin(129)	2008	MSM	Yes	1	SA	NR	19	6.00	None	NR	7
India, Wokha District, Nagaland(128)	2006	PWID	NR	NR	SA	NR	NR	NR	NR	400	4
Pakistan, Abbottabad(133)	2007	MTSW	Yes	2	IA	NR	NR	NR	NR	NR	1
Pakistan, Abbottabad(133)	2007	FSW	Yes	2	IA	NR	NR	NR	NR	NR	1
Pakistan, Lahore(133)	2007	FSW	Yes	3	SA	3	NR	NR	NR	726	7
Pakistan, Rawalpindi(133)	2007	MTSW	Yes	2	IA	NR	NR	NR	NR	NR	8

Pakistan, Rawalpindi(133)	2007	FSW	Yes	2	IA	NR	NR	NR	NR	NR	4
Thailand, Bangkok(134)	2007	FSW	NR	3	ACASI, with/ w/out interviewer	15	15	11.80	1.50	NR	7
Western Pacific											
China, Beijing(135)	2009	MSM	NR	1	ACASI	7	7	4.50	3.00	NR	5
China, Beijing(3)	2004	MSM	NR	1	SA	1	1	None	2.10	NR	3
China, Beijing(3,136)	2005	MSM	NR	1	SA	10	10	None	2.10	NR	4
China, Beijing(3)	2006	MSM	NR	1	SA	8	8	None	2.10	NR	5
China, Beijing(137)	2009	MSM	NR	1	CAPI	7	8	5.00	3.20	NR	5
China, Chongqing(138)	2009	MSM	NR	NR	CASI	7	7	4.50	3.00	NR	5
China, Guangdong(139)	2008	PWID	Yes	1	SA	6	7	7.50	3.00	238	2
China, Guangdong(140)	NR	FSW	NR	1	IA or CASI	4	4	NR	NR	NR	3
China, Guangzhou(141)	2008	MSM	NR	1	SA	13	13	5.00, gift/cash	1.50	NR	3
China, Jinan(142,143)	2007	MSM	Yes	1	SA	9	9	None	NR	428	4
China, Jinan(142,143)	2008	MSM	Yes	1	SA	5	5	None	NR	500	5
China, Jinan(144)	2008	FSW	Yes	1	SA	7	7	7.30	2.90	NR	3
China, Jinan(144)	2009	FSW	Yes	1	SA	4	4	7.30	2.90	NR	4
China, Liuzhou(145,146)	2009-10	FSW	Yes	1	SA	7	8	14.00	7.00	380	5
China, Nanjing(147)	NR	MSM	NR	1	NR	9	9	4.00 phone card	NA	NR	4
China, Shandong(148)	2007-08	Money boys	NR	NR	SA	16	NR	NR	NR	120	1
Indonesia, Bandung(149)	2007	IDU	NR	NR	SA	8	NR	NR	4.00	250	2
Indonesia, Surabaya(149)	2007	IDU	NR	NR	SA	8	NR	NR	4.00	250	2
Vietnam, Cam Ranh(150)	2005	MSM	NR	1	NR	2	NR	1.90	0.95	300	2
Vietnam, Dien Khanh(150)	2005	MSM	NR	1	NR	2	NR	1.90	0.95	300	2
Vietnam, Hai Phong(150)	2004	FSW	NR	NR	SA	20	25	3.00	1.00	200	2
Vietnam, Ho Chi Minh City(150)	2004	FSW	NR	NR	SA	20	24	4.00	1.50	400	4
Vietnam, Nha Trang(150)	2005	MSM	NR	1	NR	2	NR	1.90	0.95	300	2
Vietnam, Ninh Hoa(150)	2005	MSM	NR	1	NR	2	NR	1.90	0.95	300	2
Vietnam, Van Ninh(150)	2005	MSM	NR	1	NR	2	NR	1.90	0.95	300	2

METHODS

Literature search

We examined peer-reviewed literature published in physical or on-line journals that reported using RDS and were either accessible through September, 2013, or were identified from a previously conducted search (22). Searches were conducted using MEDLINE (1997-2013), EMBASE (1997 -2013), and Global Health (1997-2013). Search terms included “respondent driven”, “respondent-driven” or “RDS”. The original extraction included surveys in any country, in any language, and among any study population that reported using RDS (n=4562). Articles excluded in the initial extraction were those that were duplicates (n=2360), irrelevant (e.g., protocols, presentations,

flyers, etc.; n=1716) and either reviews, opinion pieces, editorials, commentaries and papers strictly addressing RDS methodology (n=44, i.e., those not intending to report population based estimates). This resulted in a total of 442 articles and abstracts. We further refined our search by eliminating abstracts (n=58) and publications that were either duplicated (n=3), non-English (n=40), without biological data (n=167), or claimed to, but did not, use RDS (n=23). When there were a number of publications for a single study, all related publications were reviewed to update the extraction sheet. This resulted in 151 articles representing 222 surveys (Figure 1).

Categorizing documents and extraction

We selected and extracted key data from 151 journal articles and entered them into a master table in Excel® into rows specific to the survey(s) described. Journal data entered into the table were organized into seven sub-tables based on WHO categorization of regions: Africa, Eastern Mediterranean (EM), Europe, Latin America and the Caribbean (LAC), North America, South-East Asia (SEA), and Western Pacific. We extracted information considered essential for assessing RDS-specific survey quality as reported in Malekinejad et al (4), Montalegre et al (5) and White et al (21,22) in each publication. The indicators reviewed included those informing survey design and implementation and analysis. Indicators informing survey design and implementation are the survey year, eligibility criteria, specimen type collected for biological testing, whether pre-survey research was conducted, number of recruitment sites, interview method, number of seeds at the start and end (and whether seeds were added or failed during data collection) of the survey, amount or type of primary and secondary incentives (USD), calculated target and final sample size, design effect used for sample

size calculation, maximum number of waves, duration of data collection (in weeks), and maximum number of coupons distributed to each recruiter. Indicators informing analysis are whether equilibrium or convergence was assessed, whether data were adjusted for network size, software used, and the citation and estimator used for adjustment. The rationale for selecting these indicators, including their usefulness in any survey versus specifically for RDS surveys, are provided in Table II.

	Percent of publications reporting information		Values of reported information	Rational for reporting
Indicators	N (222)	%	Median (range)	
Year of survey	219	99	--	Useful for any survey in order to know how current data are, to plan future surveys and to compare data from other surveys.
Eligibility criteria (minimum of behavior description) §	222	97	--	Useful for any survey to determine the denominator being measured, to know measurement for the construction of the social network question needed for

				RDS analysis. Provides readers with possible criteria to use in different populations and settings; allows for comparison of data across countries.
Type of specimen collected for biological testing §	193	87	--	Useful for any survey. Informs readers about the types of testing being conducted in different populations and settings.
Pre-survey research conducted	88	40	--	Useful for any survey. Informs readers about the survey planning process, especially whether attempts were possibly made to assess the underlying network structure of the population.
Number of recruitment sites per survey area	95	43	1 (1-6)	Especially useful in RDS as it alerts readers to the possible violation of the network being one complete component if participants at each site are not connected; informs readers

				about the possible clustering (or diffusion) of sample.
Interview method	210	94	--	Useful for any survey. Provides information about level of confidentiality in the survey (i.e., ACASI may provide more confidentiality) and informs readers about the different types of methods used for interviewing hidden populations in RDS surveys.
Number of seeds at the start of the survey	62	28	7.5 (1-48)	Specifically useful for RDS surveys. Informs readers about whether many seeds were added during data collection and the number of seeds in relation to the sample size and number of waves (too many seeds may result in too few waves needed to reduce seed dependence, adding too many seeds may be an indication that the population is not well
Whether seeds were added during data collection	17	8	5 (1-37)	
Whether seeds failed during data collection	14	6	4 (1-24)	
Number of seeds by the end of the survey	121	54	10 (1-156)	

				networked); provides parameters for readers about seeds needed in different populations and settings.
Amount or type of primary incentive (USD)	186	84	3 (1.9-5) *	Specifically useful for RDS surveys. Provides readers with parameters about amounts used in different populations and settings, provides an indication of potential bias during recruitment (if incentives are too high, more people may enroll who are not eligible).
Having no primary incentive	6	3	--	
Amount or type of secondary incentive (USD)	177	80	10 (0.95-20) †	Useful for any survey. Indicates if an original sample size was calculated and if that sample size was reached in order to ensure sufficient power and confidence for data analysis. Provides readers with parameters about sample sizes used in different populations and settings. Specific for RDS
Having no secondary incentive	18	8	--	
Calculated target sample size	89	40	300 (100-756)	
Final sample size	212	95	325 (100-1056)‡	
Final sample size for multiple cities combined	28	12	--	

				surveys: combining multiple survey sites is often a violation of the network being one network component.
Design effect used for sample size calculation §	50	22	2 (1.3-3)	design effects, currently recommended to be at least 2 (151,152,16), are important for calculating a sufficient sample size to account for RDS not being a traditional random sample
Maximum number of waves	95	43	9 (3-21)	
Duration of data collection (in weeks)	139	63	12 (2-124)	Informs readers of the time needed to gather samples of different sizes from different populations and settings; alerts readers of unusual recruitment lengths that may impact representativeness of the sample.

Maximum number of coupons distributed to each recruiter §	163	73	3 (2-7)	Specifically useful for RDS surveys. The number of coupons used are normally three (7), but some surveys have used more. Analysis does not account for branching induced by the number of coupons provided to each participant so fewer coupons, when possible, is suggested to mimic a random walk process.
Whether equilibrium or convergence was assessed§	44	20		Specifically useful for RDS surveys. Informs readers of seed dependence and is a diagnostic to assess bias.
Whether data were adjusted for network size	157	70	--	Specifically useful for RDS surveys. Informs readers of the extent to which RDS was fully utilized, resulting in the ability to assess whether the survey may represent the network of the population from which the sample was gathered

Software used to adjust data §	162	73	--	Specifically useful for RDS surveys. There are limited software packages available for analyzing RDS data. Analyzing RDS data in more popular, preexisting software (i.e., STATA, SPSS) may not eliminate RDS specific biases.
Citation for adjustment §	59	26	--	Specifically useful for RDS surveys. Given the evolvement of the estimators for the analysis of RDS data, this is useful for providing information about the assumptions supporting the adjustments.
Heckathorn (10,153) §	19	32	--	
Salganik & Heckathorn (12) §	28	47	--	
Heckathorn (11) §	10	17	--	
Volz & Heckathorn (13) §	7	12	--	
Gile (154) §	4	7	--	
Estimator used for adjustment §	10	4	--	
Whether seeds were discarded during analysis	31	14	--	Some studies either did not collect data from seeds or did not include their data their analysis, which could likely

				<p>result in the sample having addition seeds (analysis would assume wave 1 participants are seeds) thereby potentially impacting seed dependence and biasing the final estimate.</p>
--	--	--	--	---

Analysis

Frequencies were used to characterize the surveys and their contents. We conducted robust and logistic regressions of survey start year and pre-survey research, eligibility age, number of seeds at the start and end of survey, survey duration, final sample size, estimated design effect, length of longest recruitment chain, and adjustment of RDS to assess linear trends in the value of these indicators over time (155). Design effects were calculated for surveys that presented a point estimate for HIV prevalence, 95% confidence intervals and the final sample size. The calculation for design effects consisted of dividing the widths of the confidence interval by two, dividing again by 1.96 (the standard normal value corresponding to a central area of 95%), and squaring the final number.

RESULTS

The identified published articles of RDS surveys were conducted in the following WHO regions: 21 from Africa (28 surveys), 12 from EM (11 surveys), 30 from Europe (44 surveys), 17 from LAC (37 surveys), 41 from North America (45 surveys), 12 from SEA (32 surveys), and 18 from the Western Pacific (25 surveys). Extracted surveys included

85 among PWID, 78 among MSM and 38 among FSW. Surveys of other groups included people who use and/or inject drugs (n=2), male sex workers (n=3), high-risk heterosexuals (n=7), transgender (n=2), and youth (n=3). The remaining surveys were of mixed groups such as youth PWID (n=2), people who use and/or inject drugs together (n=1) and MSM who use and/or inject drugs (n=1).

Assessing reports of survey quality

Survey data extracted from published articles included in this review were used to assess whether RDS recruitment and analysis were conducted, but the details provided for these surveys varied across articles. For instance, all published surveys reported basic implementation information such as the city, country and the population sampled, and 99% reported the survey year (Table II). Over 90% of surveys reported the interview technique (e.g., face-to-face questionnaire, computer-assisted self-interviews, etc.) (94%), final sample size (95%) and at least the behavioral component of the eligibility criteria (97%). Eighty four percent reported the primary and 80% reported the secondary incentive amounts or types, and 73% reported the maximum number of coupons given to each recruiter. Sixty-three percent reported the data collection duration, 40% reported whether pre-survey research was conducted, 43% reported the number of recruitment sites used and the maximum number of waves, 40% reported the target sample size, 22% reported the design effect used for calculating the sample size. For those surveys that presented both calculated and final sample sizes (n=77, 35%), the median percentage difference was 1.0 (range 0.2-1.6). There was no significant difference in this measure over time among all populations combined or by population.

Seventy percent reported whether data were adjusted for network size, 73% reported the type of software used to adjust data (74% of which used RDS Analysis Tool [RDSAT]) and 26% cited the statistical adjustment, among which 47% cited Salganik & Heckathorn (12), 32% cited Heckathorn (10) and/or 2002, 17% Heckathorn (11), 12% Volz & Heckathorn (13) and 7% Gile (14,154). Only 20% of surveys reported whether equilibrium or convergence was assessed and 4% reported which estimator was used for their statistical adjustment. Thirty-one surveys (14%) specifically reported discarding seeds from their analysis.

Design effects for HIV

Of the 222 surveys reviewed, 185 reported HIV prevalence point estimates above 0, 136 included 95% confidence bounds, and 210 reported final sample sizes. Ninety-five surveys (42.7%) included all three elements to enable calculation of the estimated design effect for HIV prevalence. Four (4.2%) had a design effect less than 1.0, 28 (29.5%) had a design effect of 1.0, 46 (48.4%) had a design effect of 2. The remaining design effects were as high as 5.9, indicating that a larger sample size was needed to estimate HIV prevalence.

Assessing changes over time

In assessing changes over time (Table III), we found significant decreases in eligibility age, final number of seeds, and final sample size ($p < 0.01$, for all) and significant increases in pre-survey research and using a design effect to calculate the target sample size ($p < 0.01$). There were no significant changes for survey duration even when adjusting for target population and final sample size. Nor were there significant changes by year for survey duration or length of longest recruitment chain.

DISCUSSION

Reporting on details of survey design, implementation, and analysis is essential for assessing the quality of RDS surveys and findings. It is important to adequately describe both the methodological and analytical aspects of RDS in any publication. The preponderance of publications from surveys reported the most essential information such as survey city, country, year, population sampled, interview method, and final sample size. Given that all publications from surveys reported collecting biological specimens, it is surprising that 13% did not provide information about specimen collection and testing methods. Gaps in reporting RDS methodological and analytical information made it difficult to assess survey quality and the strength of results. RDS does not work in all situations and failure to meet assumptions should be noted. For instance, only 43% of surveys reported the maximum number of waves and 20% reported assessment of equilibrium or convergence, information needed to assess potential biases. Among those surveys reporting their maximum number of waves, some reported having only a maximum of three waves, indicating that the survey results were likely biased by the non-randomly selected seeds.

Although pre-survey research should be part of any survey, it is increasingly recognized as an important part of any RDS survey (7,15,22,156), as evidenced by the increase in surveys reporting having conducted formative research. Because RDS samples a social network, formative research is imperative to understand the underlying network structure of the sampled population. If the sampled network is fragmented or has isolated sub-groups, the chances of sampling more than one network are higher, possibly resulting in unstable estimates (15). Furthermore, pre-survey research data

can help investigators plan survey operations and encourage participation by learning about which survey procedures would be most acceptable to the target population. It is possible that the increased use of pre-survey research helped improve recruitment and led to the decrease in the final number of seeds reported over time. Fewer seeds generally occur concomitantly with longer chains, which in turn means decreased bias stemming from seed dependence. We recommend that all surveys using RDS conduct pre-survey research to evaluate social networks, as well as to assess the feasibility of using RDS in a particular population.

Although 70% of surveys reported whether data were adjusted for network size and 73% reported the software used to adjust those data, few cited the adjustment procedure and even fewer reported the estimator used. There are currently at least five different estimators for adjusting RDS data (157). Given that many of the reviewed articles were written before the existence of some estimators, it is understandable that earlier publications did not cite the estimator used for analysis. Forthcoming publications should cite the estimator since knowing this information will allow readers to know how adjustments were made, if they were made properly, and the assumptions supporting those adjustments.

Several publications reported discarding seeds from analysis. While it has been written that “seeds are eliminated from analysis” (13,153), this is not to say that seeds should be manually eliminated from a dataset. The RDS-I and RDS-II estimators (11–13,153) use a matrix of recruits and recruiters whereby data from the recruits are necessary for calculating inclusion probabilities used to derive final estimates. Even though the seeds do not technically show up in the probability matrix since they were

never recruited by their peers, their data are nonetheless necessary for establishing the placement of the seeds' recruits in the matrix.

We found an increase over time in surveys reporting design effects, an element in the sample size calculations to account for RDS not being a simple random sample. Although recent publications have found that design effects of 3 or 4 would be optimal, in most situations, a design effect of 2 is sufficient and recommended (16,151,152). Because operational constraints, such as limited financial resources, often preclude large sample sizes for some RDS surveys, using a design effect greater than 2 may result in unfeasibly large sample sizes. Post-hoc design effects on key variables can help determine if sample sizes were large enough for the analysis and inform sample size calculations for follow-up surveys of the same population. As such it is useful for publications to include point estimates, 95% confidence intervals and final sample size to allow for the post-hoc estimation of design effects.

Equilibrium or convergence was reported in only 20% of the articles reviewed. Equilibrium, the term most often used when referring to RDS surveys, measures the progression of waves to determine when the proportion for a characteristic approaches and remains stable in relation to the final sample statistic (10). Convergence, a more sensitive measurement, measures the progression of enrolling subjects to determine when the proportion for a characteristic approaches and remains stable in relation to the adjusted estimate (15). Nevertheless, the assessment of either equilibrium or convergence is useful for determining seed dependence, a typical bias found in chain referral sampling methods, and should be reported for publications reporting population estimates from RDS surveys (22).

While most surveys reported a minimum eligibility age of 18 years (n=150), we found the minimum age decreased over time. Collecting HIV and other biological and behavioral data from younger key populations is important given they are disproportionately affected by HIV worldwide and are comprising a high percentage of new HIV infections (9,158).

Our review has limitations. As in any systematic review, we are restricted by the completeness of our publication search and whether investigators published their surveys in peer-reviewed journals. Furthermore, we only included surveys that collected biological data leaving room for further evaluation of those surveys that reported using RDS and did not collect biological data. The number of peer-reviewed articles found reporting RDS surveys is far fewer than the actual number of surveys conducted. Although key data were missing from articles, this is in itself an important finding which supports the need to uniformly report results from RDS surveys. It also limited the scope of our analyses and introduced uncertainty into some of our other findings (22). We excluded articles clearly stating they either used RDS 'recruitment' only or did not fulfill necessary features of the method; however, we may have included some surveys that did not incorporate all RDS methodological and analytical features, given their incomplete reporting. In those instances, we classified the surveys as using RDS and included them in the extraction. Several of the 23 articles claiming to use RDS, but did not, reported using a 'modified' or 'mixed methods' RDS. However, they did not provide conclusive evidence such as the collection and use of personal network size data, recruitment ties (who recruited whom) and coupon quotas, and multiple recruitment waves. In several extracted publications, significant limitations were reported, including

unprepared staff, numerous ineligible persons trying to participate, closing or moving survey sites during data collection, overly high incentives (possible indication of enrollment of ineligible participants) or too low, overcrowding at the interview site, failure to recruit important population sub-groups (i.e., females in PWID surveys, older MSM), incorrect or no social network question, and early survey termination due to finances or community disturbances (3,34,41,55,86,127,159). Presenting key limitations is useful for interpreting findings and should be included in all publications presenting data from RDS surveys.

The majority of published surveys were from North American and Europe; it would be useful to see more publications of RDS survey results from other regions. Not only could experiences from these different settings help researchers improve survey methods and analysis, but the results themselves could help policy makers, donors, and service providers to improve responses to HIV and other infection risk. Future publications of biological and behavioral surveys using RDS should provide a minimum set of parameters in order for readers to assess specific methodological and analytical procedures and to make determinations of the overall quality of these surveys.

REFERENCES

1. Des Jarlais DC, Arasteh K, Perlis T, Hagan H, Abdul-Quader A, Heckathorn DD, et al. Convergence of HIV seroprevalence among injecting and non-injecting drug users in New York City. *AIDS*. 2007;21(2):231–5.
2. Johnston LG, Sabin K, Hien MT, Huong PT, Mai TH, Pham TH. Assessment of respondent driven sampling for recruiting female sex workers in two Vietnamese cities: Reaching the unseen sex worker. *J Urban Heal*. 2006;83(6 Suppl):i16–28.
3. Ma X, Zhang Q, He X, Sun W, Yue H, Chen S, et al. Trends in prevalence of HIV, syphilis, hepatitis C, hepatitis B, and sexual risk behavior among men who have sex with men: results of 3 consecutive respondent-driven sampling surveys in Beijing, 2004 through 2006. *J Acquir Immune Defic Syndr*. 2007;45(5):581–7.
4. Malekinejad M, Johnston LG, Kendall C, Kerr LRFS, Rifkin MR, Rutherford GW. Using respondent-driven sampling methodology for HIV biological and behavioral surveillance in international settings: a systematic review. *AIDS Behav*. 2008;12(4 Suppl):S105–30.
5. Montealegre JR, Johnston LG, Murrill C, Monterroso E. Respondent driven sampling for HIV biological and behavioral surveillance in Latin America and the Caribbean. *AIDS Behav*. 2013;17(7):2313–40.
6. Lansky A. Assessing the assumptions of respondent-driven sampling in the national HIV behavioral surveillance system among injecting drug users. *Open AIDS J*. 2012;6:77–82.

7. Johnston LG. Introduction to Respondent-Driven Sampling. Geneva, Switzerland: World Health Organization; 2013.
http://applications.emro.who.int/dsaf/EMRPUB_2013_EN_1539.pdf. Accessed 15 Jun 2015.
8. UNAIDS. Guidelines on surveillance among populations most at risk for HIV. 2011
http://www.unaids.org/sites/default/files/en/media/unaids/contentassets/documents/epidemiology/2011/20110518_Surveillance_among_most_at_risk.pdf. Accessed 15 Jun 2015.
9. UNICEF, UNESCO, UNFPA, UNAIDS. Young Key Populations at Higher Risk of HIV in Asia and the Pacific: Making the Case with Strategic Information. Bangkok, Thailand: UNICEF East Asia and Pacific Regional Office; 2013.
http://www.unicef.org/eapro/Young_key_populations_at_high_risk_of_HIV_in_Asia_Pacific.pdf Accessed 15 Jun 2015.
10. Heckathorn DD. Respondent-driven sampling: a new approach to the study of hidden populations. *Soc Probl.* 1997;44(2):174–99.
11. Heckathorn DD. Extensions of respondent-driven sampling: analyzing continuous variables and controlling for differential recruitment. *Sociol Methodol.* 2007;37(1):151–207.
12. Salganik MJ, Heckathorn DD. Sampling and estimation in hidden populations using respondent-driven sampling. *Sociol Methodol.* 2004;34(1):193–240.
13. Volz E, Heckathorn DD. Probability based estimation theory for respondent driven sampling. *J Off Stat.* 2008;24(1):79–97.
14. Gile KJ. Improved inference for respondent-driven sampling data with application

- to HIV prevalence estimation. *J Am Stat Assoc.* 2011;106(493):135–46.
15. Gile KJ, Johnston LG, Salganik MJ. Diagnostics for respondent-driven sampling. *J R Stat Soc.* 2015 Jan 1;1(1):241–69.
 16. Salganik MJ. Variance estimation, design effects, and sample size calculations for respondent-driven sampling. *J urban Heal.* 2006;83(6 Suppl):i98–112.
 17. Szwarcwald CL, de Souza Júnior PRB, Damacena GN, Junior AB, Kendall C. Analysis of data collected by RDS among sex workers in 10 Brazilian cities, 2009: estimation of the prevalence of HIV, variance, and design effect. *J Acquir Immune Defic Syndr.* 2011;57 Suppl 3:S129–35.
 18. McCreesh N, Frost SDW, Seeley J, Katongole J, Tarsh MN, Ndunguse R, et al. Evaluation of respondent-driven sampling. *Epidemiology.* 2012;23(1):138–47.
 19. Goel S, Salganik MJ. Assessing respondent-driven sampling. *Proc Natl Acad Sci U S A.* 2010;107:6743–7.
 20. Wejnert C. An empirical test of respondent-driven sampling: Point estimates, variance, degree measures, and out-of-equilibrium data. *Sociol Methodol.* 2009;39:73–116.
 21. Hafeez S. A review of the proposed STROBE-RDS reporting checklist as an effective tool for assessing the reporting quality of RDS studies from the developing world. London, United Kingdom. 2012.
 22. White RG, Hakim AJ, Salganik MJ, Spiller MW, Johnston LG, Kerr L, et al. Strengthening the reporting of observational studies in epidemiology for respondent-driven sampling studies: “STROBE-RDS” statement. *J Clin Epidemiol.* 2015;68(12):1463–71.

23. Vandenhoudt HM, Langat L, Menten J, Odongo F, Oswago S, Luttah G, et al. Prevalence of HIV and other sexually transmitted infections among Female Sex Workers in Kisumu, Western Kenya, 1997 and 2008. *PLoS One*. 2013;8(1):1–16.
24. Vu L, Adebajo S, Tun W, Sheehy M, Karlyn A, Njab J, et al. High HIV prevalence among men who have sex with men in Nigeria: implications for combination prevention. *J Acquir Immune Defic Syndr*. 2013;63(2):221–7.
25. Merrigan M, Azeez A, Afolabi B, Chabikuli ON, Onyekwena O, Eluwa G, et al. HIV prevalence and risk behaviours among men having sex with men in Nigeria. *Sex Transm Infect*. 2011;87:65–70.
26. Eluwa GI, Strathdee S a., Adebayo SB, Ahonsi B, Adebajo SB. A profile on HIV prevalence and risk behaviors among injecting drug users in Nigeria: Should we be alarmed? *Drug Alcohol Depend*. 2013;127(1-3):65–71.
27. Adebajo SB, Eluwa GI, Allman D, Myers T, Ahonsi BA. Prevalence of internalized homophobia and HIV associated risks among men who have sex with men in Nigeria. *Afr J Reprod Health*. 2012;16(4):21–8.
28. Johnston L, Saumtally A, Corceal S, Mahadoo I, Oodally F. High HIV and hepatitis C prevalence amongst injecting drug users in Mauritius: findings from a population size estimation and respondent driven sampling survey. *Int J Drug Policy*. 2011;22(4):252–8.
29. Johnston LG, Corceal S. Unexpectedly high injection drug use, HIV and hepatitis C prevalence among female sex workers in the Republic of Mauritius. *AIDS Behav*. 2013;17(2):574–84.
30. Kriitmaa K, Testa A, Osman M, Bozicevic I, Riedner G, Malungu J, et al. HIV

- prevalence and characteristics of sex work among female sex workers in Hargeisa, Somaliland, Somalia. *AIDS*. 2010;24 Suppl 2:S61–7.
31. Rispel LC, Metcalf CA, Cloete A, Reddy V, Lombard C. HIV prevalence and risk practices among men who have sex with men in two South African cities. *J Acquir Immune Defic Syndr*. 2011;57(1):69–76.
 32. Lane T, Raymond HF, Dladla S, Rasethe J, Struthers H, McFarland W, et al. High HIV prevalence among men who have sex with men in Soweto, South Africa: Results from the Soweto men's study. *AIDS Behav*. 2011;15:626–34.
 33. Johnston L, O'Bra H, Chopra M, Mathews C, Townsend L, Sabin K, et al. The associations of voluntary counseling and testing acceptance and the perceived likelihood of being HIV-infected among men with multiple sex partners in a South African township. *AIDS Behav*. 2010;14(4):922–31.
 34. Chopra M, Townsend L, Johnston L, Mathews C, Tomlinson M, Bra HO, et al. Estimating HIV Prevalence and Risk Behaviors Among High-Risk Heterosexual Men With Multiple Sex Partners : Use of Respondent-Driven Sampling. *J Acquir Immune Defic Syndr*. 2009;51(1):72–7.
 35. Townsend L, Rosenthal SR, Parry CDH, Zembe Y, Mathews C, Flisher AJ. Associations between alcohol misuse and risks for HIV infection among men who have multiple female sexual partners in Cape Town, South Africa. *AIDS Care*. 2010;22(12):1544–54.
 36. Townsend L, Johnston LG, Flisher AJ, Mathews C, Zembe Y. Effectiveness of respondent-driven sampling to recruit high risk heterosexual men who have multiple female sexual partners: differences in HIV prevalence and sexual risk

- behaviours measured at two time points. *AIDS Behav.* 2010;14(6):1330–9.
37. Zembe YZ, Townsend L, Thorson A, Ekström AM. Predictors of inconsistent condom use among a hard to reach population of young women with multiple sexual partners in peri-urban south Africa. *PLoS One.* 2012;7(12).
 38. Townsend L, Zembe Y, Mathews C, Mason-Jones A. Estimating HIV Prevalence and HIV-Related Risk Behaviors Among Heterosexual Women Who Have Multiple Sex Partners Using Respondent-Driven Sampling in a High-Risk Community in South Africa. *J Acquir Immune Defic Syndr.* 2013;62(4):457–64.
 39. Abdelrahim MS. HIV prevalence and risk behaviors of female sex workers in Khartoum, north Sudan. *AIDS.* 2010;24 Suppl 2:S55–60.
 40. Johnston LG, Holman A, Dahoma M, Miller LA, Kim E, Mussa M, et al. HIV risk and the overlap of injecting drug use and high-risk sexual behaviours among men who have sex with men in Zanzibar (Unguja), Tanzania. *Int J Drug Policy.* 2010;21(6):485–92.
 41. Dahoma M, Johnston LG, Holman A, Miller LA, Mussa M, Othman A, et al. HIV and related risk behavior among men who have sex with men in Zanzibar, Tanzania: results of a behavioral surveillance survey. *AIDS Behav.* 2011;15(1):186–92.
 42. Hladik W, Barker J, Ssenkusu JM, Opio A, Tappero JW, Hakim A, et al. HIV infection among men who have sex with men in kampala, uganda-a respondent driven sampling survey. *PLoS One.* 2012;7(5):1–9.
 43. Soliman C, Rahman I a, Shawky S, Bahaa T, Elkamhawi S, El Sattar AA, et al. HIV prevalence and risk behaviors of male injection drug users in Cairo, Egypt.

- AIDS. 2010;24 Suppl 2:S33–8.
44. Navadeh S, Mirzazadeh A, Mousavi L, Haghdoost A, Fahimfar N, Sedaghat A. HIV, HSV2 and Syphilis Prevalence in Female Sex Workers in Kerman, South-East Iran; Using Respondent-Driven Sampling. *Iran J Public Health*. 2012;41(12):60–5.
 45. Mahfoud Z, Afifi R, Ramia S, El Khoury D, Kassak K, El Barbir F, et al. HIV/AIDS among female sex workers, injecting drug users and men who have sex with men in Lebanon: results of the first biobehavioral surveys. *AIDS*. 2010;24 Suppl 2:S45–54.
 46. Kassak K, Mahfoud Z, Kreidieh K, Shamra S, Afifi R, Ramia S. Hepatitis B virus and hepatitis C virus infections among female sex workers and men who have sex with men in Lebanon: Prevalence, risk behaviour and immune status. *Sex Health*. 2011;8:229–33.
 47. Mahfoud Z, Kassak K, Kreidieh K, Shamra S, Ramia S. Distribution of hepatitis C virus genotypes among injecting drug users in Lebanon. *Viol J*. 2010;7:96.
 48. Mirzoyan L, Berendes S, Jeffery C, Thomson J, Ben Othman H, Danon L, et al. New evidence on the HIV epidemic in Libya: why countries must implement prevention programs among people who inject drugs. *J Acquir Immune Defic Syndr*. 2013;62(5):577–83.
 49. Valadez JJ, Berendes S, Jeffery C, Thomson J, Ben Othman H, Danon L, et al. Filling the knowledge gap: measuring HIV prevalence and risk factors among men who have sex with men and female sex workers in Tripoli, Libya. *PLoS One*. 2013;8(6).

50. Johnston LG, Alami K, El Rhilani MH, Karkouri M, Mellouk O, Abadie A, et al. HIV, syphilis and sexual risk behaviours among men who have sex with men in Agadir and Marrakesh, Morocco. *Sex Transm Infect.* 2013;89 Suppl 3:iii45–8.
51. Stulhofer A, Chetty A, Rabie RA, Jwehan I, Ramlawi A. The prevalence of HIV, HBV, HCV, and HIV-related risk-taking behaviors among Palestinian injecting drug users in the East Jerusalem Governorate. *J Urban Heal.* 2012;89(4):671–6.
52. Stormer A, Tun W, Guli L, Harxhi A, Bodanovskaia Z, Yakovleva A, et al. An analysis of respondent driven sampling with Injection Drug Users (IDU) in Albania and the Russian Federation. *J urban Heal.* 2006;83(6 Suppl):i73–82. t
53. Elda S, Bani R. An analysis of HIV-Related Risk Behaviors of Men Having Sex with Men (MSM), Using Respondent Driven Sampling (RDS), in Albania. *Int J Med.* 2009 ;2(2):231–5
54. Bozicevic I, Lepej SZ, Rode OD, Grgic I, Jankovic P, Dominkovic Z, et al. Prevalence of HIV and sexually transmitted infections and patterns of recent HIV testing among men who have sex with men in Zagreb, Croatia. *Sex Transm Infect.* 2012;88(7):539–44.
55. Bozicevic I, Rode OD, Lepej SZ, Johnston LG, Stulhofer A, Dominkovic Z, et al. Prevalence of sexually transmitted infections among men who have sex with men in Zagreb, Croatia. *AIDS Behav.* 2009;13(2):303–9.
56. Lepej SZ, Vrakela IB, Poljak M, Bozicevic I, Begovac J. Phylogenetic analysis of HIV sequences obtained in a respondent-driven sampling study of men who have sex with men. *AIDS Res Hum Retroviruses.* 2009;25(12):1335–8.
57. Mills HL, Colijn C, Vickerman P, Leslie D, Hope V, Hickman M. Respondent

- driven sampling and community structure in a population of injecting drug users, Bristol, UK. *Drug Alcohol Depend.* 2012;126(3):324–32.
58. Hope VD, Hickman M, Ngui SL, Jones S, Telfer M, Bizzarri M, et al. Measuring the incidence, prevalence and genetic relatedness of hepatitis C infections among a community recruited sample of injecting drug users, using dried blood spots. *J Viral Hepat.* 2011;18:262–70.
 59. Platt L, Bobrova N, Rhodes T, Uusküla A, Parry J V, Rüütel K, et al. High HIV prevalence among injecting drug users in Estonia: implications for understanding the risk environment. *AIDS.* 2006;20(16):2120–3.
 60. Abel-Ollo K, Rahu M, Rajaleid K, Talu A, Ruutel K, Platt L, et al. Knowledge of HIV serostatus and risk behaviour among injecting drug users in Estonia. *AIDS Care.* 2009;21(7):851–7.
 61. Vorobjov S, Uusküla A, Abel-Ollo K, Talu A, Rüütel K, Des Jarlais DC. Comparison of injecting drug users who obtain syringes from pharmacies and syringe exchange programs in Tallinn, Estonia. *Harm Reduct J.* 2009;6(1):3.
 62. Uusküla A, Des Jarlais DC, Kals M, Rüütel K, Abel-Ollo K, Talu A, et al. Expanded syringe exchange programs and reduced HIV infection among new injection drug users in Tallinn, Estonia. *BMC Public Health.* 2011;11(1):517.
 63. Uusküla A, Johnston LG, Raag M, Trummal A, Talu A, Des Jarlais DC. Evaluating recruitment among female sex workers and injecting drug users at risk for HIV using respondent-driven sampling in Estonia. *J urban Heal.* 2010;87(2):304–17.
 64. Uusküla A, Fischer K, Raudne R, Kilgi H, Krylov R, Salminen M, et al. A study on HIV and hepatitis C virus among commercial sex workers in Tallinn. *Sex Transm*

- Infect. 2008;84(3):189–91.
65. Talu A, Rajaleid K, Abel-Ollo K, Rüütel K, Rahu M, Rhodes T, et al. HIV infection and risk behaviour of primary fentanyl and amphetamine injectors in Tallinn, Estonia: Implications for intervention. *Int J Drug Policy*. 2010;21:56–63.
 66. Berry M, Wirtz AL, Janayeva A, Ragoza V, Terlikbayeva A, Amirov B, et al. Risk factors for HIV and unprotected anal intercourse among men who have sex with men (MSM) in Almaty, Kazakhstan. *PLoS One*. 2012;7(8):e43071.
 67. Zohrabyan L, Johnston LG, Scutelnicuic O, Iovita A, Todirascu L, Costin T, et al. Determinants of HIV infection among female sex workers in two cities in the Republic of Moldova: the role of injection drug use and sexual risk. *AIDS Behav*. 2013;17(8):2588–96.
 68. Busza J, Douthwaite M, Bani R, Scutelnicuic O, Preda M, Simic D. Injecting behaviour and service use among young injectors in Albania, Moldova, Romania and Serbia. *Int J Drug Policy*. 2013;24(5):423–31.
 69. Zohrabyan L, Johnston L, Scutelnicuic O, Iovita a, Todirascu L, Costin T, et al. HIV, hepatitis and syphilis prevalence and correlates of condom use during anal sex among men who have sex with men in the Republic of Moldova. *Int J STD AIDS*. 2013;24:357–64.
 70. Baćak V, Laušević D, Mugoša B, Vratnica Z, Terzić N. Hepatitis C virus infection and related risk factors among injection drug users in montenegro. *Eur Addict Res*. 2013;19:68–73.
 71. Judd A, Rhodes T, Johnston LG, Platt L, Andjelkovic V, Simić D, et al. Improving survey methods in sero-epidemiological studies of injecting drug users: a case

- example of two cross sectional surveys in Serbia and Montenegro. *BMC Infect Dis.* 2009;9:14.
72. Eritsyan KU, Levina OS, White E, Smolskaya TT, Heimer R. HIV prevalence and risk behavior among injection drug users and their sex partners in two Russian cities. *AIDS Res Hum Retroviruses.* 2013;29(4):687–90.
 73. Niccolai LM, Shcherbakova IS, Toussova O V., Kozlov AP, Heimer R. The potential for bridging of HIV transmission in the russian federation: Sex risk behaviors and hiv prevalence among drug users (DUs) and their non-du sex partners. *J Urban Heal.* 2009;86(1):131–43.
 74. Paintsil E, Verevochkin S V., Dukhovlina E, Niccolai L, Barbour R, White E, et al. Hepatitis C virus infection among drug injectors in St Petersburg, Russia: Social and molecular epidemiology of an endemic infection. *Addiction.* 2009;104:1881–90.
 75. Iguchi MY, Ober AJ, Berry SH, Fain T, Heckathorn DD, Gorbach PM, et al. Simultaneous recruitment of drug users and men who have sex with men in the united states and Russia using respondent-driven sampling: Sampling methods and implications. *J Urban Heal.* 2009;86(1):5–31.
 76. Djonic D, Djuric M, Bassioni-Stamenic F, McFarland W, Knezevic T, Nikolic S, et al. HIV-related risk behaviors among roma youth in Serbia: Results of two community-based surveys. *J Adolesc Heal.* 2013;52(2):234–40.
 77. Taran YS, Johnston LG, Pohorila NB, Saliuk TO. Correlates of HIV risk among injecting drug users in sixteen Ukrainian cities. *AIDS Behav.* 2011;15(1):65–74.
 78. Pando M a., Balán IC, Marone R, Dolezal C, Leu CS, Squiquera L, et al. HIV and

other sexually transmitted infections among men who have sex with men recruited by RDS in Buenos Aires, Argentina: High HIV and HPV infection. *PLoS One*. 2012;7(6).

79. Pando M, Marone R, Balan I, Dolezal C, Squiquera L, Picconi A, et al. HIV and STI prevalence among men who have sex with men (MSM) recruited through respondent driven sampling (RDS) in Buenos Aires, Argentina. *Retrovirology*. *BioMed Central*; 2009;6(Suppl 3):P103.
80. Salani Mota RM, Sansigolo Kerr LRF, Kendall C, Pinho A, de Mello MB, Dourado I, et al. Reliability of self-report of HIV status among men who have sex with men in Brazil. *J Acquir Immune Defic Syndr*. 2011;57:S153–6.
81. Kerr LRF, Mota RS, Kendall C, Pinho A de A, Mello MB, Guimarães MDC, et al. HIV among MSM in a large middle-income country. *AIDS*. 2013;27(3):427–35.
82. Bermúdez-Aza EH, Kerr LRF, Kendall C, Pinho A a, Mello MB De, Mota RS, et al. Antiretroviral Drug Resistance in a Respondent-Driven Sample of HIV-Infected Men Who Have Sex With Men in Brazil. *J Acquir Immune Defic Syndr*. 2011;57:S186–92.
83. Damacena GN, Szwarcwald CL, de Souza Júnior PRB, Dourado I. Risk factors associated with HIV prevalence among female sex workers in 10 Brazilian cities. *J Acquir Immune Defic Syndr*. 2011;57 Suppl 3:S144–52.
84. Tun W, de Mello M, Pinho a, Chinaglia M, Diaz J. Sexual risk behaviours and HIV seroprevalence among male sex workers who have sex with men and non-sex workers in Campinas, Brazil. *Sex Transm Infect*. 2008;84:455–7.
85. Martins TA, Kerr LRF, Macena RHM, Mota RS, Carneiro KL, Gondim RC, et al.

- Travestis, an unexplored population at risk of HIV in a large metropolis of northeast Brazil: a respondent-driven sampling survey. *AIDS Care*. 2013;25(5):606–12.
86. Kendall C, Kerr LRFS, Gondim RC, Werneck GL, Macena RHM, Pontes MK, et al. An empirical comparison of respondent-driven sampling, time location sampling, and snowball sampling for behavioral surveillance in men who have sex with men, Fortaleza, Brazil. *AIDS Behav*. 2008;12(4 Suppl):S97–104.
 87. Johnston LG, Vaillant TC, Dolores Y, Vales HM. HIV, hepatitis B/C and syphilis prevalence and risk behaviors among gay, transsexuals and men who have sex with men, Dominican Republic. *Int J STD AIDS*. 2013;24(4):313–21.
 88. Creswell J, Guardado ME, Lee J, Nieto a. I, Kim a. a., Monterroso E, et al. HIV and STI control in El Salvador: results from an integrated behavioural survey among men who have sex with men. *Sex Transm Infect*. 2012;
 89. Dennis AM, Murillo W, de Maria Hernandez F, Guardado ME, Nieto AI, Lorenzana de Rivera I, et al. Social network-based recruitment successfully reveals HIV-1 transmission networks among high-risk individuals in El Salvador. *J Acquir Immune Defic Syndr*. 2013;63(1):135–41. t
 90. Johnston LG, Paz-Bailey G, Morales-Miranda S, Morgan M, Alvarez B, Hickman L, et al. High prevalence of *Mycoplasma genitalium* among female sex workers in Honduras: implications for the spread of HIV and other sexually transmitted infections. *Int J STD AIDS*. 2012. 23(1):5–11.
 91. Silva-Santisteban A, Raymond HF, Salazar X, Villayzan J, Leon S, McFarland W, et al. Understanding the HIV/AIDS epidemic in transgender women of lima, peru:

- Results from a sero-epidemiologic study using respondent driven sampling. *AIDS Behav.* 2012;16:872–81.
92. Moyer LB, Brouwer KC, Brodine SK, Ramos R, Lozada R, Cruz MF, et al. Barriers and missed opportunities to HIV testing among injection drug users in two Mexico--US border cities. *Drug Alcohol Rev.* 2008;27:39–45.
 93. Deiss RG, Brouwer KC, Loza O, Lozada RM, Ramos R, Cruz MAF, et al. High-risk sexual and drug using behaviors among male injection drug users who have sex with men in 2 Mexico-US border cities. *Sex Transm Dis.* 2008;35(3):243–9.
 94. Philbin M, Pollini R a., Ramos R, Lozada R, Brouwer KC, Ramos ME, et al. Shooting gallery attendance among IDUs in Tijuana and Ciudad Juarez, Mexico: correlates, prevention opportunities, and the role of the environment. *AIDS Behav.* 2008;12:552–60.
 95. Frost SDW, Brouwer KC, Firestone Cruz M a., Ramos R, Ramos ME, Lozada RM, et al. Respondent-driven sampling of injection drug users in two U.S.-Mexico border cities: recruitment dynamics and impact on estimates of HIV and syphilis prevalence. *J Urban Heal.* 2006;83.
 96. White EF, Garfein RS, Brouwer KC, Lozada R, Ramos R, Firestone-Cruz M, et al. Prevalence of hepatitis C virus and HIV infection among injection drug users in two Mexican cities bordering the U.S. *Salud Publica Mex.* 2007;49(3):165–72.
 97. Baumbach JP, Foster LN, Mueller M, Cruz MF, Arbona S, Melville S, et al. Seroprevalence of select bloodborne pathogens and associated risk behaviors among injection drug users in the Paso del Norte region of the United States-Mexico border. *Harm Reduct J.* 2008;5(1):33.

98. Rusch ML, Lozada R, Pollini RA, Vera A, Patterson TL, Case P, et al. Polydrug use among IDUs in Tijuana, Mexico: correlates of methamphetamine use and route of administration by gender. *J urban Heal*. 2009;86(5):760–75.
99. Garfein RS, Lozada R, Liu L, Laniado-Laborin R, Rodwell TC, Deiss R, et al. High prevalence of latent tuberculosis infection among injection drug users in Tijuana, Mexico. *Int J Tuberc lung Dis*. 2009;13(5):626–32.
100. Strathdee SA, Lozada R, Ojeda VD, Pollini RA, Brouwer KC, Vera A, et al. Differential effects of migration and deportation on HIV infection among male and female injection drug users in Tijuana, Mexico. *PLoS One*. 2008;3(7):e2690.
101. Brouwer KC, Rusch ML, Weeks JR, Lozada R, Vera A, Magis-Rodríguez C, et al. Spatial Epidemiology of HIV among Injection Drug Users in Tijuana, Mexico. *Ann Assoc Am Geogr*. Taylor & Francis Group; 2012;102(5):1190–9.
102. Abramovitz D, Volz EM, Strathdee SA, Patterson TL, Vera A, Frost SDW. Using respondent-driven sampling in a hidden population at risk of HIV infection: who do HIV-positive recruiters recruit? *Sex Transm Dis*. 2009;36(12):750–6.
103. Stephens DB, Havens JR. Predictors of alcohol use among rural drug users after disclosure of hepatitis C virus status. *J Stud Alcohol Drugs*. 2013;74:386–95.
104. Boodram B, Golub ET, Ouellet LJ. Socio-behavioral and geographic correlates of prevalent hepatitis C virus infection among young injection drug users in metropolitan Baltimore and Chicago. *Drug Alcohol Depend*. 2010;111(1-2):136–45.
105. Villanti AC, German D, Sifakis F, Flynn C, Holtgrave DR. Smoking, self-reported HIV and HIV risk behaviors in a respondent-driven sample of injection drug users

- in Baltimore, Maryland: the BeSure study. *AIDS Educ Prev.* 2012;24(2):132–47.
106. Mimiaga MJ, Reisner SL, Bland S, Skeer M, Cranston K, Isenberg D, et al. Health system and personal barriers resulting in decreased utilization of HIV and STD testing services among at-risk black men who have sex with men in Massachusetts. *AIDS Patient Care STDS.* 2009;23(10):825–35.
 107. Mimiaga MJ, Reisner SL, Cranston K, Isenberg D, Bright D, Daffin G, et al. Sexual mixing patterns and partner characteristics of black msm in massachusetts at increased risk for HIV infection and transmission. *J Urban Heal.* 2009;86(4):602–23.
 108. Risser JMH, Padgett P, Wolverton M, Risser WL. Relationship between heterosexual anal sex, injection drug use and HIV infection among black men and women. *Int J STD AIDS.* 2009;20:310–4.
 109. King WD, Larkins S, Hucks-Ortiz C, Wang P-C, Gorbach PM, Veniegas R, et al. Factors associated with HIV viral load in a respondent driven sample in Los Angeles. *AIDS Behav.* 2009;13(1):145–53.
 110. Shoptaw S, Weiss RE, Munjas B, Hucks-Ortiz C, Young SD, Larkins S, et al. Homonegativity, substance use, sexual risk behaviors, and HIV status in poor and ethnic men who have sex with men in Los Angeles. *J urban Heal.* 2009;86 Suppl 1:77–92.
 111. Gorbach PM, Murphy R, Weiss RE, Hucks-Ortiz C, Shoptaw S. Bridging sexual boundaries: Men who have sex with men and women in a street-based sample in Los Angeles. *J Urban Heal.* 2009;86(1):63–76.
 112. Gelpí-Acosta C, Hagan H, Jenness SM, Wendel T, Neaigus A. Sexual and

- injection-related risks in Puerto Rican-born injection drug users living in New York City: A mixed-methods analysis. *Harm Reduct J*. 2011;8(1):28.
113. Jenness SM, Begier EM, Neaigus A, Murrill CS, Wendel T, Hagan H. Unprotected anal intercourse and sexually transmitted diseases in high-risk heterosexual women. *Am J Public Health*. 2011;101(4):745–50.
 114. Jenness SM, Neaigus A, Murrill CS, Wendel T, Forgione L, Hagan H. Estimated HIV incidence among high-risk heterosexuals in New York City, 2007. *J Acquir Immune Defic Syndr*. 2011;56(2):193–7.
 115. Jenness SM, Murrill CS, Liu K-L, Wendel T, Begier E, Hagan H. Missed opportunities for HIV testing among high-risk heterosexuals. *Sex Transm Dis*. 2009;36(12):704–10.
 116. Jenness SM, Kobrak P, Wendel T, Neaigus A, Murrill CS, Hagan H. Patterns of exchange sex and HIV infection in high-risk heterosexual men and women. *J Urban Heal*. 2011;88(2):329–41.
 117. Jenness SM, Neaigus A, Hagan H, Murrill CS, Wendel T. Heterosexual HIV and sexual partnerships between injection drug users and noninjection drug users. *AIDS Patient Care STDS*. 2010;24(3):175–81.
 118. Hagan H, Jenness SM, Wendel T, Murrill CR, Neaigus A, Gelpi-Acosta C. Herpes simplex virus type 2 associated with HIV infection among New York heterosexuals living in high-risk areas. *Int J STD AIDS*. 2010;21(8):580–3.
 119. Abdul-Quader AS, Heckathorn DD, McKnight C, Bramson H, Nemeth C, Sabin K, et al. Effectiveness of respondent-driven sampling for recruiting drug users in New York City: findings from a pilot study. *J Urban Heal*. 2006;83(3):459–76.

120. McKnight C, Des Jarlais D, Bramson H, Tower L, Abdul-Quader AS, Nemeth C, et al. Respondent-driven sampling in a study of drug users in New York City: notes from the field. *J Urban Heal.* 2006;83(6 Suppl):i54–9.
121. McCoy SI, Shiu K, Martz TE, Smith CD, Mattox L, Gluth DR, et al. Improving the efficiency of HIV testing with peer recruitment, financial incentives, and the involvement of persons living with HIV infection. *J Acquir Immune Defic Syndr.* 2013;63(2):e56–63.
122. Garfein RS, Rondonelli A, Barnes RFW, Cuevas J, Metzner M, Velasquez M, et al. HCV infection prevalence lower than expected among 18-40-year-old injection drug users in San Diego, CA. *J Urban Heal.* 2013;90(3):516–28.
123. Wei C, McFarland W, Colfax GN, Fuqua V, Raymond HF. Reaching black men who have sex with men: a comparison between respondent-driven sampling and time-location sampling. *Sex Transm Infect.* 2012;88(8):622–6.
124. Burt RD, Thiede H. Evaluating consistency in repeat surveys of injection drug users recruited by respondent-driven sampling in the Seattle area: results from the NHBS-IDU1 and NHBS-IDU2 surveys. *Ann Epidemiol.* 2012;22(5):354–63.
125. Magnus M, Kuo I, Shelley K, Rawls A, Peterson J, Montanez L, et al. Risk factors driving the emergence of a generalized heterosexual HIV epidemic in Washington, District of Columbia networks at risk. *AIDS.* 2009;23(November 2008):1277–84.
126. Magnus M, Kuo I, Phillips G, Rawls A, Peterson J, Montanez L, et al. Differing HIV risks and prevention needs among men and women injection drug users (IDU) in the District of Columbia. *J urban Heal.* 2013;90(1):157–66.

127. Johnston LG, Khanam R, Reza M, Khan SI, Banu S, Alam MS, et al. The effectiveness of respondent driven sampling for recruiting males who have sex with males in Dhaka, Bangladesh. *AIDS Behav.* 2008;12(2):294–304.
128. Mahanta J, Medhi GK, Paranjape RS, Roy N, Kohli A, Akoijam BS, et al. Injecting and sexual risk behaviours, sexually transmitted infections and HIV prevalence in injecting drug users in three states in India. *AIDS.* 2008;22 Suppl 5:S59–68.
129. Solomon SS, Srikrishnan AK, Sifakis F, Mehta SH, Vasudevan CK, Balakrishnan P, et al. The emerging HIV epidemic among men who have sex with men in tamil nadu, india: Geographic diffusion and bisexual concurrency. *AIDS Behav.* 2010;14:1001–10.
130. Barua P, Jagadish M, Kumar MG, Jayesh D, Ramesh P, Gay T. Sexual activity as a risk factor for hepatitis C virus (HCV) transmission among the female sex workers in Nagaland. *Indian Journal of Medical Research.* 2012.
131. Shahmanesh M, Cowan F, Wayal S, Copas A, Patel V, Mabey D. The burden and determinants of HIV and sexually transmitted infections in a population-based sample of female sex workers in Goa, India. *Sex Transm Infect.* 2009;85(1):50–9.
132. Shahmanesh M, Wayal S, Copas A, Patel V, Mabey D, Cowan F. A study comparing sexually transmitted infections and HIV among ex-red-light district and non-red-light district sex workers after the demolition of Baina red-light district. *J Acquir Immune Defic Syndr.* 2009;52(2):253–7.
133. Hawkes S, Collumbien M, Platt L, Lalji N, Rizvi N, Andreasen a, et al. HIV and other sexually transmitted infections among men, transgenders and women selling sex in two cities in Pakistan: a cross-sectional prevalence survey. *Sex*

- Transm Infect. 2009;85 Suppl 2:ii8–i16.
134. Manopaiboon C, Prybylski D, Subhachaturas W, Tanpradech S, Suksripanich O, Siangphoe U, et al. Unexpectedly high HIV prevalence among female sex workers in Bangkok, Thailand in a respondent-driven sampling survey. *Int J STD AIDS*. 2013;24(1):34–8.
 135. Li X, Lu H, Raymond HF, Sun Y, Jia Y, He X, et al. Untested and undiagnosed: barriers to HIV testing among men who have sex with men, Beijing, China. *Sex Transm Infect*. 2012;88:187–93.
 136. Ma X-Y, Zhang Q-Y, He X, Zhao J-K, Li Y, Sun W-D, et al. Epidemiological study on the status of HIV/STDs and relative behaviors among MSM in Beijing. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2007;28(9):851–5.
 137. Fan S, Lu H, Ma X, Sun Y, He X, Li C, et al. Behavioral and Serologic Survey of Men Who Have Sex with Men in Beijing, China: Implication for HIV Intervention. *AIDS Patient Care STDS*. 2012;26(3):148–55.
 138. Zhang L, Ding X, Lu R, Feng L, Li X, Xiao Y, et al. Predictors of hiv and syphilis among men who have sex with men in a chinese metropolitan city: Comparison of risks among students and non-students. *PLoS One*. 2012;7(5).
 139. Lin P, Wang M, Li Y, Zhang Q, Yang F, Zhao J. Detoxification center-based sampling missed a subgroup of higher risk drug users, a case from Guangdong, China. *PLoS One*. 2012;7(4):e35189.
 140. Li Y, Detels R, Lin P, Fu X, Deng Z, Liu Y, et al. Prevalence of HIV and STIs and associated risk factors among female sex workers in Guangdong Province, China. *J Acquir Immune Defic Syndr*. 2010;53 Suppl 1:S48–53.

141. Zhong F, Lin P, Xu H, Wang Y, Wang M, He Q, et al. Possible increase in HIV and syphilis prevalence among men who have sex with men in Guangzhou, China: Results from a respondent-driven sampling survey. *AIDS Behav.* 2011;15(1):1058–66.
142. Ruan S, Yang H, Zhu Y, Wang M, Ma Y, Zhao J, et al. Rising HIV prevalence among married and unmarried among men who have sex with men: Jinan, China. *AIDS Behav.* 2009;13:671–6.
143. Ruan S, Yang H, Zhu Y, Ma Y, Li J, Zhao J, et al. HIV prevalence and correlates of unprotected anal intercourse among men who have sex with men, Jinan, China. *AIDS Behav.* 2008 May;12(3):469–75.
144. Liao M, Nie X, Pan R, Wang C, Ruan S, Zhang C, et al. Consistently low prevalence of syphilis among female sex workers in Jinan, China: Findings from two consecutive respondent driven sampling surveys. *PLoS One.* 2012;7(4).
145. Weir SS, Merli MG, Li J, Gandhi AD, Neely WW, Edwards JK, et al. A comparison of respondent-driven and venue-based sampling of female sex workers in Liuzhou, China. *Sex Transm Infect.* 2012;88 Suppl 2:i95–101.
146. Li J, Chen X-S, Merli MG, Weir SS, Henderson GE. Systematic differences in risk behaviors and syphilis prevalence across types of female sex workers: a preliminary study in Liuzhou, China. *Sex Transm Dis.* 2012;39(3):195–200. /
147. Hao C, Yan H, Yang H, Huan X, Guan W, Xu X, et al. The incidence of syphilis, HIV and HCV and associated factors in a cohort of men who have sex with men in Nanjing, China. *Sex Transm Infect.* 2011;87:199–201.
148. Tao X, Gai R, Zhang N, Zheng W, Zhang X, Xu A, et al. HIV infection and mental

- health of “money boys”: A pilot study in Shandong Province, China. *Southeast Asian J Trop Med Public Health*. 2010;41:358–68.
149. Morineau G, Bollen L, Ika Syafitri R, Nurjannah N, Erti Mustikawati D, Magnani R. HIV prevalence and risk behaviours among injecting drug users in six Indonesian cities implications for future HIV prevention programs. *Harm Reduct J. Harm Reduction Journal*; 2012;9(1):37. Available from: Harm Reduction Journal
 150. Colby D, Minh TT, Toan TT. Down on the farm: homosexual behaviour, HIV risk and HIV prevalence in rural communities in Khanh Hoa province, Vietnam. *Sex Transm Infect*. 2008;84:439–43.
 151. Wejnert C, Pham H, Krishna N, Le B, DiNenno E. Estimating design effect and calculating sample size for Respondent-driven sampling studies of injection drug users in the United States. *AIDS Behav*. 2012;16(4):797–806.
 152. Johnston LG, Chen Y-H, Silva-Santisteban A, Raymond HF. An empirical examination of respondent driven sampling design effects among HIV risk groups from studies conducted around the world. *AIDS Behav*. 2013;17(6):2202–10.
 153. Heckathorn DD. Respondent-driven sampling II: deriving valid population estimates from chain-referral samples of hidden populations. *Soc Probl*. 2002;49(1):11–34.
 154. Gile KJ, Handcock MS. Respondent-driven sampling: an assessment of current methodology. *Sociol Methodol*. 2010;40(1):285–327.
 155. Yohai VJ. High Breakdown-Point and High Efficiency Robust Estimates for Regression. *Ann Stat*. 1987;15(2):642–56.
 156. Johnston LG, Whitehead S, Simic-Lawson M, Kendall C. Formative research to

- optimize respondent-driven sampling surveys among hard-to-reach populations in HIV behavioral and biological surveillance: lessons learned from four case studies. *AIDS Care*. 2010;22:784–92.
157. Johnston LG, Luthra RR. Analyzing data in RDS. In: Tyldum G, Johnston L, editors. *Applying Respondent Driven Sampling to Migrant Populations Lessons from the Field*. Palgrave Pivot; 2014
158. UNICEF. *Opportunity in Crisis: Preventing HIV from Early Adolescence to Young Adulthood*. Geneva, Switzerland.2011.. Accessed on 4 April 2016 at: http://www.unicef.org/publications/files/Opportunity_in_Crisis-Report_EN_052711.pdf
159. Johnston LG, Malekinejad M, Kendall C, Iuppa IM, Rutherford GW. Implementation challenges to using respondent-driven sampling methodology for HIV biological and behavioral surveillance: field experiences in international settings. *AIDS Behav*. 2008;12(4 Suppl):S131–41.