

LONDON  
SCHOOL of  
HYGIENE  
& TROPICAL  
MEDICINE



LSHTM Research Online

Tarr, SJ; Moon, RW; Hardege, I; Osborne, AR; (2014) A conserved domain targets exported PHISTb family proteins to the periphery of Plasmodium infected erythrocytes. *Molecular and biochemical parasitology*, 196 (1). pp. 29-40. ISSN 0166-6851 DOI: <https://doi.org/10.1016/j.molbiopara.2014.07.011>

Downloaded from: <http://researchonline.lshtm.ac.uk/2545995/>

DOI: <https://doi.org/10.1016/j.molbiopara.2014.07.011>


**Usage Guidelines:**

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact [researchonline@lshtm.ac.uk](mailto:researchonline@lshtm.ac.uk).


Available under license: <http://creativecommons.org/licenses/by/2.5/>

<https://researchonline.lshtm.ac.uk>

## A PF3D7\_0401800:GFP in Dd2



## B



### Supplementary figure 2

(A) PF3D7\_0401800:GFP (PFD80) expressed in *P. falciparum* Dd2. The left- and right-hand images show GFP localisation and a phase contrast image, respectively. Scale bar, 2  $\mu$ m.

(B) Anti-GFP Western blot of *P. falciparum* Dd2 schizonts expressing PF3D7\_0401800:GFP (PFD80)(lane 2).  $2 \times 10^6$  schizonts were loaded per lane. Black arrow: expressed protein. Lane 1 contains untransfected *P. falciparum* Dd2 parasites.