

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Datta, J; Palmer, MJ; Tanton, C; Gibson, LJ; Jones, KG; Macdowall, W; Glasier, A; Sonnenberg, P; Field, N; Mercer, CH; +2 more... Johnson, AM; Wellings, K; (2016) Prevalence of infertility and help seeking among 15 000 women and men. Human reproduction (Oxford, England). ISSN 0268-1161
DOI: <https://doi.org/10.1093/humrep/dew123>

Downloaded from: <http://researchonline.lshtm.ac.uk/2572522/>

DOI: <https://doi.org/10.1093/humrep/dew123>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

<https://researchonline.lshtm.ac.uk>

Prevalence of infertility and help seeking among 15 000 women and men

J. Datta^{1,*}, M.J. Palmer¹, C. Tanton², L.J. Gibson¹, K.G. Jones²,
W. Macdowall¹, A. Glasier^{3,1}, P. Sonnenberg², N. Field², C.H. Mercer²,
A.M. Johnson², and K. Wellings¹

¹Centre for Sexual and Reproductive Health Research, Department of Social and Environmental Health Research, London School of Hygiene & Tropical Medicine, 15-17 Tavistock Place, London WC1H 9SH, UK ²Research Department of Infection and Population Health, University College London, Mortimer Market Centre, London WC1E 6JB, UK ³MRC Centre for Reproductive Health, Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh EH16 4TY, UK

*Correspondence address. E-mail: jessica.datta@lshtm.ac.uk

Submitted on January 13, 2016; resubmitted on April 25, 2016; accepted on April 29, 2016

STUDY QUESTION: What is the prevalence of infertility and of help seeking among women and men in Britain?

SUMMARY ANSWER: One in eight women and one in ten men aged 16–74 years had experienced infertility, defined by unsuccessfully attempting pregnancy for a year or longer, and little more than half of these people sought medical or professional help.

WHAT IS KNOWN ALREADY: Estimates of infertility and help seeking in Britain vary widely and are not easily comparable because of different definitions and study populations.

STUDY DESIGN, SIZE, DURATION: A cross-sectional population survey was conducted between 2010 and 2012 with a sample of 15 162 women and men aged 16–74 years.

PARTICIPANTS/MATERIALS, SETTING, METHODS: Participants completed the Natsal-3 questionnaire, using computer-assisted personal interviewing (CAPI) and computer-assisted self-interview (CASI).

MAIN RESULTS AND THE ROLE OF CHANCE: The reported prevalence of infertility was 12.5% (CI 95% 11.7–13.3) among women and 10.1% (CI 95% 9.2–11.1) among men. Increased prevalence was associated with later cohabitation with a partner, higher socio-economic status and, for those who had a child, becoming parents at older ages. The reported prevalence of help seeking was 57.3% (CI 95% 53.6–61.0) among women and 53.2% (CI 95% 48.1–58.1) among men. Help seekers were more likely to be better educated and in higher status occupations and, among those who had a child, to have become parents later in life.

LIMITATIONS, REASONS FOR CAUTION: These data are cross-sectional so it is not possible to establish temporality or infer causality. Self-reported data may be subject to recall bias.

WIDER IMPLICATIONS OF THE FINDINGS: The study provides estimates of infertility and help seeking in Britain and the results indicate that the prevalence of infertility is higher among those delaying parenthood. Those with higher educational qualifications and occupational status are more likely to consult with medical professionals for fertility problems than others and these inequalities in help seeking should be considered by clinical practice and public health.

STUDY FUNDING/COMPETING INTEREST(S): Funding was provided by grants from the Medical Research Council and the Wellcome Trust, with support from the Economic and Social Research Council and the Department of Health. AMJ is a Governor of the Wellcome Trust. Other authors have no competing interests.

Key words: infertility / female infertility / male infertility / help seeking / population survey

Introduction

Fertility is a key element of reproductive health and infertility is recognized as a global public health issue by the World Health Organisation (WHO) (Boivin *et al.*, 2007; Macaluso *et al.*, 2010). Infertility is defined by the International Committee for Monitoring Assisted Reproductive Technology (ICMART) and WHO as the 'failure to achieve a pregnancy after 12 months or more of regular unprotected sexual intercourse' (Zegers-Hochschild *et al.*, 2009). The National Institute for Health and Care Excellence (NICE) guideline for England and Wales concurs with this time period, advocating that clinical assessment and investigation should be offered when 'a woman of reproductive age... has not conceived after one year of unprotected vaginal sexual intercourse, in the absence of any known cause of infertility' (National Institute for Health and Clinical Excellence, 2013). The experience of infertility can cause those affected personal distress (Schneider and Forthofer, 2005; Greil *et al.*, 2011), significant treatment costs (Bell, 2010) and, in some contexts, ostracism and discrimination (Cui, 2010).

Infertility is experienced by an estimated 48.5 million couples worldwide (Mascarenhas *et al.*, 2012) and around 1 in 7 couples in the UK (National Institute for Health and Clinical Excellence, 2013). However, prevalence estimates of lifetime infertility vary widely, in part because there is no agreed or consistent definition of infertility (Gurunath *et al.*, 2011) and because study populations vary by age range, unit of measurement and relationship status.

A marked trend for delaying the timing of first birth has been seen in developed and, more recently, developing countries (Gyimah, 2003; Mathews and Hamilton, 2009; Rosero-Bixby *et al.*, 2009; Mills *et al.*, 2011). A consequence of postponement of parenthood is the potential for impaired fertility (Balasch, 2010) and the risk that deferral of parenthood will rule it out (Sobotka, 2006; Leridon, 2008).

Later parenthood, its implications for fertility, and the increased range and availability of fertility treatments are likely to have had an impact on help seeking. There is some evidence of an increase in the reporting of fertility problems to primary care after the introduction of national treatment guidelines, suggesting more awareness of assisted reproductive technology (ART) (Dhalwani *et al.*, 2013). Studies have found that women in later age cohorts are more likely to seek help than women in earlier ones and that women are seeking help at older ages than was previously the case (Oakley *et al.*, 2008; Wilkes *et al.*, 2009). There are few population-based data on experience of infertility and its determinants or on the extent of help seeking, particularly for men. This study uses data from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3) to estimate the prevalence of infertility, and of seeking medical or professional help among women and men in Britain, and to examine associated factors.

Materials and Methods

Study population

Natsal-3 is a survey of 15 162 women (8869) and men (6293) aged 16–74 years. Households were selected using stratified probability sampling from which one eligible individual resident in Britain was selected at random and invited to participate. The sample frame was the Postcode Address File (PAF), a regularly updated list of all addresses in the country. As the PAF excludes those who are homeless or living in institutions, our sample is representative of individuals living in private residential households. Data were

weighted in two stages to correct for participants' unequal probabilities of selection. The first corrected for the selection of one household in multi-household addresses and for the varying probabilities of selection by number of adults within households. The second adjusted for differential non-response by comparing age, sex and region profile of participants with 2011 census data. Although the Natsal-3 sample closely matched those who responded to the census in terms of ethnicity (86.8% in Natsal-3 and 86.7% in the census were white), there was a slight under-representation of Asian women and men in Natsal-3 (6.4%) compared with the census (7.5%). Participants were interviewed between 2010 and 2012 using computer-assisted personal interviewing (CAPI), which included a computer-assisted self-interview (CASI) component for the more sensitive questions. The response rate was 57.7% and the co-operation rate (i.e. of all eligible addresses contacted) was 65.8%. Further details of the methods are described elsewhere (Erens *et al.*, 2014).

Questions about experience of infertility and help seeking for infertility were asked of all Natsal-3 participants who reported ever having heterosexual intercourse (8315 women and 5742 men). Two discrete questions were asked: 'Have you ever had a time, lasting 12 months or longer, when you and a partner were trying for a pregnancy but it didn't happen?' and 'Have you (or a partner) ever sought medical or professional help about infertility?'. In line with the NICE guideline (National Institute for Health and Clinical Excellence, 2013), we considered a participant to have experienced infertility when she or he responded 'yes' to the first question. Participants were also asked questions that comprise the validated Patient Health Questionnaire-2 (PHQ-2) (Kroenke *et al.*, 2003), a composite measure of depression experienced in the 2 weeks prior to interview and questions on satisfaction with sex life and relationship in the past year.

Data are cross-sectional and we did not ask participants for information about when their experience of infertility occurred. Some may have failed to conceive in a 12-month period before becoming parents or between pregnancies while others may never have become parents. Those still of reproductive age could experience a future period of infertility. Natsal-3 data are deposited at the UK Data Service. <https://discover.ukdataservice.ac.uk/catalogue/?sn=7799> (11 May 2016, date last accessed).

Statistical analyses

We used STATA v13.1 (StataCorp, 2013) to undertake complex survey analyses to account for the weighting, clustering and stratification of the Natsal-3 data. We estimated the population prevalence of infertility and help seeking, stratified by gender and age group. We used multivariable logistic regression to explore associations (adjusted for age at interview) between experience of (i) infertility and (ii) help seeking, and a number of socio-demographic, relationship, reproductive and health factors, including: age at interview; relationship status at interview; age at first cohabitation; age at first child; academic attainment; employment status measured using National Statistics Socio-Economic Classification (NS-SEC) and area-level deprivation measured using the Index of Multiple Deprivation (IMD) (Office for National Statistics, 2010). We report results for educational attainment for those aged over 21 years only as younger participants may not have completed full-time education. Finally, using multivariable logistic regression, we present age-adjusted odds ratios to describe associations between experience of infertility and selected health and relationship 'outcome' variables for women aged 50 years or younger. We selected this age group for analysis as their experience of infertility will have occurred more recently than that of older participants.

Ethical approval

The Natsal-3 study was approved by the Oxfordshire Research Ethics Committee A (ref.: 10/H0604/27).

Table 1 Prevalence of and factors associated with infertility by sex.

	Women					Men						
	Pregnancy attempt 12 months or longer		Age-adjusted regression			Pregnancy attempt 12 months or longer		Age-adjusted regression				
	%	95% CI	AOR	95% CI	P-Value	Denominators (unweighted, weighted)	%	95% CI	AOR	95% CI	P-Value	Denominators (unweighted, weighted)
All	12.5%	(11.7–13.3)				8066, 7052	10.1%	(9.2–11.1)				5553, 6811
Age at interview					<0.0001						<0.0001	
16–24	5.3	(4.3–6.6)	1.00			1695, 944	3.7	(2.6–5.1)	1.00			1325, 971
25–34	12.0	(10.7–13.5)	2.43	(1.87–3.17)		2366, 1306	8.5	(7.0–10.2)	2.44	(1.62–3.67)		1421, 1274
35–44	17.7	(15.7–20.0)	3.85	(2.92–5.06)		1173, 1403	14.9	(12.4–17.8)	4.62	(3.10–6.87)		780, 1377
45–54	12.5	(10.6–14.7)	2.56	(1.90–3.45)		1062, 1374	14.5	(11.8–17.7)	4.48	(2.95–6.81)		741, 1328
55–64	13.2	(11.0–15.6)	2.71	(2.00–3.66)		972, 1170	7.8	(5.9–10.3)	2.24	(1.41–3.56)		697, 1094
65–74	11.4	(9.2–14.0)	2.29	(1.65–3.18)		798, 854	8.2	(6.1–11.0)	2.35	(1.47–3.74)		589, 767
Relationship history												
Relationship status at interview					<0.0001						<0.0001	
Married/cohabiting	15.2	(14.1–16.4)	1.00	-		4348, 4669	12.6	(11.3–14.0)	1.00	-		2925, 4640
Non-cohabiting partnership	6.1	(4.8–7.8)	0.39	(0.29–0.51)		1355, 786	4.0	(2.7–5.8)	0.29	(0.18–0.45)		941, 755
No 'steady' partner	7.6	(6.5–9.0)	0.46	(0.38–0.57)		2326, 1573	5.3	(4.2–6.7)	0.39	(0.29–0.51)		1653, 1384
Ever cohabited with a partner					<0.0001						<0.0001	
Never	2.2	(1.5–3.2)	1.00			1499, 855	2.1	(1.3, 3.3)	1.00			1481, 1151
Ever	13.8	(12.9–14.8)	7.50	(5.01–11.25)		6350, 6013	11.8	(10.7, 13.0)	6.91	(4.13–11.55)		3920, 5465
Age at first cohabitation					0.0685						0.8975	
Under 20	12.9	(11.3–14.7)	1.00			1970, 1685	11.6	(9.0–14.9)	1.00			574, 677
20–29	13.8	(12.7–15.1)	1.09	(0.91–1.30)		3972, 3856	11.9	(10.6–13.4)	1.05	(0.77–1.44)		2852, 3982
30 or older	17.6	(13.9–22.0)	1.45	(1.06–1.99)		402, 464	11.1	(8.5–14.5)	0.98	0.65–1.48)		465, 763
Reproductive history												
Ever had a child					0.0880						0.0070	
Yes	13.2	(12.2–14.3)	1.00			5248, 5110	11.7	(10.4–13.2)	1.00			2694, 4031
No	10.5	(9.2–11.9)	0.83	(0.68–1.03)		2808, 1935	7.9	(6.7–9.3)	0.68	(0.52–0.90)		2857, 2775
Age at birth of first child ^b					<0.0001						<0.0001	
<25	9.0	(7.9–10.3)	1.00			3047, 2709	7.6	(5.9–9.7)	1.00			946, 1234
25–29	13.2	(11.3–15.3)	1.54	(1.23–1.92)		1337, 1392	10.2	(8.1–12.7)	1.44	(1.00–2.06)		902, 1363
30–34	20.9	(17.6–24.6)	2.63	(2.03–3.40)		616, 709	14.5	(11.4–18.3)	2.08	(1.41–3.08)		520, 852
35+	35.1	(28.6–42.1)	5.57	(3.97–7.83)		227, 278	19.8	(15.3–25.4)	3.27	(2.16–4.96)		301, 550

Continued

Table I Continued

	Women					Men					
	Pregnancy attempt 12 months or longer		Age-adjusted regression			Pregnancy attempt 12 months or longer		Age-adjusted regression			
	%	95% CI	AOR	95% CI	P-Value	%	95% CI	AOR	95% CI	P-Value	
					Denominators (unweighted, weighted)					Denominators (unweighted, weighted)	
Abortion ever					0.5855						
No	12.4	(11.5–13.3)	1.00	-		6853, 6046					
Yes	12.9	(10.7–15.5)	1.07	(0.84–1.35)		1176, 968					
Socio-economic position											
Educational level ^a					0.0006					0.5684	
Degree	14.5	(12.8–16.3)	1.00			1938, 1755	12.0	(10.1–14.2)	1.00		1375, 1904
A-level/equivalent	13.5	(11.6–15.7)	0.90	(0.72–1.14)		1546, 1420	11.0	(9.1–13.1)	0.90	(0.68–1.19)	1336, 1762
GCSE, O-level or equivalent	13.9	(12.4–15.6)	0.93	(0.76–1.13)		2496, 2294	10.3	(8.6–12.4)	0.84	(0.64–1.12)	1382, 1750
Foreign or other	11.3	(5.8–21.0)	0.76	(0.36, 1.62)		86, 67	12.9	(5.8–26.5)	1.11	(0.44–2.77)	46, 57
None	8.8	(7.1–10.9)	0.53	(0.40, 0.71)		966, 950	9.2	(6.7–12.4)	0.73	(0.48–1.09)	530, 701
Social class (NS-SEC)					<0.0001					0.0003	
Managerial and prof occupations	15.2	(13.5–17.0)	1.00			2341, 2230	12.8	(11.1–14.6)	1.00		1786, 2514
Intermediate occupations	13.5	(11.6–15.7)	0.87	(0.70, 1.09)		1586, 1426	10.0	(7.9–12.6)	0.76	(0.55–1.03)	871, 1159
Semi-routine/routine occupations	11.1	(9.7–12.7)	0.70	(0.58, 0.86)		2282, 1874	9.3	(7.9–11.0)	0.71	(0.55–0.91)	1926, 2235
No job (10+ h/week) or not in the last 10 years	11.7	(9.8–14.0)	0.71	(0.55, 0.93)		1098, 1045	7.4	(4.7–11.4)	0.52	(0.31–0.88)	387, 426
Student in full-time education	2.9	(1.8–4.7)	0.18	(0.11, 0.30)		721, 443	2.3	(0.9–5.8)	0.18	(0.07–0.49)	565, 456
Index of multiple deprivation (IMD)—quintile					0.5362					0.2590	
1–2 (least deprived)	13.2	(11.9–14.6)	1.00			3062, 2909	11.1	(9.6–12.8)	1.00		2197, 2870
3	12.5	(10.7–14.5)	0.96	(0.78–1.18)		1570, 1371	8.7	(6.9–11.0)	0.78	(0.57–1.05)	1080, 1329
4–5 (most deprived)	11.7	(10.5–13.1)	0.91	(0.76–1.08)		3434, 2772	9.8	(8.4–11.4)	0.90	(0.71–1.15)	2276, 2613

Denominator: All those who reported having experience of heterosexual sex and who gave a valid answer to the question on experience of infertility.

AOR, adjusted odds ratio; 95% CI, 95% confidence intervals.

^aParticipants aged 21 and older only.

^bOnly those who had a child.




Figure 1 Prevalence of infertility and help seeking, according to age groups. (A) Women who had attempted pregnancy with a partner for 12 months or longer. (B) Men who had attempted pregnancy with a partner for 12 months or longer.

Results

Prevalence of infertility and associated factors

The proportion of participants aged 16–74 who reported ever having tried unsuccessfully for a year or longer to become pregnant was 12.5% among women and 10.1% among men (Table I). Unsurprisingly, the prevalence of ever experience of infertility was lowest in the youngest women and peaked among those aged 35–44 years (17.7%). A similar pattern was seen for men but the age range in which the prevalence of ever experience of infertility was highest was wider than for women, extending from age 35 to 54 years at interview.

The experience of infertility was associated with relationship factors. It was highest among those who were married or cohabiting at interview and lowest among those in a non-cohabiting partnership. Among women, a positive association was found between older age at first cohabitation and the experience of infertility. Age-adjusted odds of experiencing infertility were significantly higher among women who first gave birth at age 35 or older (AOR 5.57 (3.97–7.83)) compared with those who did so before age 25. A similar, though slightly weaker, association was observed among men (AOR 3.27 (2.16–4.96)).

We found evidence of associations between infertility and indicators of socio-economic status. Experience of infertility was most common among women with a degree and lowest among those with no academic qualifications while no statistically significant association was observed among men. Prevalence was also higher among those in managerial, professional and technical employment compared with those in routine occupations. No association was found between infertility and area-related deprivation at interview among women or men.

Among women, the only health indicator found to be meaningfully associated with infertility was treatment for depression in the previous 12 months and the association was of only borderline significance (Supplementary data, Table SI). Among men, we found associations with ever having been diagnosed with a sexually transmitted infection (men: AOR 1.39 (1.02–1.88)) and, specifically, with chlamydia (AOR 1.81 (1.15–2.84)).

Prevalence of help seeking for infertility and associated factors

A total of 57.3% of women and 53.2% of men who had ever experience of infertility had sought medical or professional help as a result (Table II). Figure 1 shows the proportion of women and men with experience of infertility who sought help for the problem by age group at interview.

Less than one-third (32.6%) of the youngest women (16–24 years) and only 14.1% of the youngest men with experience of infertility reported seeking help. Among both women and men, similar proportions of those aged 35–74 had sought help (women: 58.0–64.9%; men: 53.1–58.1%).

Age-adjusted odds ratios showed help seeking to be higher among those who were currently married or cohabiting compared with those in non-cohabiting or not in 'steady' partnerships. Prevalence of seeking help was highest in women who first cohabited in their twenties while, among men, it was highest in those who first cohabited aged 30 or older.

For women, having had a child and older age at first birth were both associated with seeking help for fertility; help seeking was highest among women who became mothers at 35 or older (74.6% (62.8–83.7)). The association between age at first child and help seeking was less marked among men. Of those who had never had a child, 58.7% (95% CI 51.5–65.4) of women and 54.9% (95% CI 46.6–62.9) of men reported having sought help.

Help seeking was significantly associated with indicators of socio-economic position. Women and men with lower levels of education and lower occupational classifications (as indicated by the NS-SEC) were less likely to have sought help. Associations between help seeking and area-level deprivation at interview were of borderline significance among women only.

Health indicators were not associated with likelihood of having sought help for infertility although there was a borderline association suggesting that men who reported regular heavy drinking were slightly less likely to have sought help (Supplementary data, Table SII).

Associations between experience of infertility and aspects of current well-being and relationship quality among women

Table III presents the associations between experience of infertility and three variables which we have treated as 'outcomes' for the purposes of

Table II Prevalence of and factors associated with help seeking among those with experience of infertility by sex.

	Women					Men						
	%	95% CI	AOR	95% CI	P-Value	Denominators (unweighted, weighted)	%	95% CI	AOR	95% CI	P-Value	Denominators (unweighted, weighted)
All	57.3	(53.6–61.0)				923, 879	53.2	(48.1–58.1)				470, 691
Age group					<0.0001						0.0004	
16–24	32.6	(22.9–44.0)	0.49	0.28–0.86		93, 50	14.1	(6.8–27.1)	0.17	0.07–0.41		47, 36
25–34	49.5	(43.1–55.9)	1.00			278, 157	49.6	(40.1–59.1)	1.00			112, 108
35–44	63.3	(56.0–70.0)	1.76	1.18–2.63		208, 249	56.9	(47.2–66.2)	1.34	0.78–2.33		116, 206
45–54	58.0	(49.0–66.5)	1.41	0.91–2.20		136, 172	56.1	(45.4–66.3)	1.30	0.73–2.30		94, 193
55–64	58.3	(48.8–67.2)	1.43	0.90–2.27		120, 154	58.1	(43.8–71.1)	1.41	0.71–2.80		54, 86
65–74	64.9	(54.0–74.4)	1.89	1.13–3.15		88, 97	53.1	(38.7–67.0)	1.15	0.57–2.30		47, 63
Relationship history												
Relationship status at interview					<0.0001						0.0049	
Married/cohabiting	60.5	(56.3–64.5)	1.00			660, 710	56.1	(50.5–61.6)	1.00			348, 585
Non-cohabiting partnership	47.5	(35.6–59.7)	0.68	0.40–1.13		88, 48	28.9	(15.2–48.1)	0.35	0.15–0.81		34, 30
No 'steady' partner	43.2	(35.1–51.7)	0.43	0.30–0.64		173, 120	39.9	(29.1–51.7)	0.53	0.32–0.89		85, 74
Ever cohabited with a partner					0.0202						0.0017	
Never	26.7	(12.3–48.7)	1.00			37, 19	14.9	(6.1–32.3)	1.00			25, 24
Ever	58.1	(54.3–61.9)	3.10	1.19–8.07		853, 832	54.7	(49.3–59.9)	5.63	1.91–16.58		424, 643
Age at first cohabitation					0.0838						0.0259	
Under 20	51.8	(44.6–59.0)	1.00			253, 217	36.9	(25.3–50.2)	1.00			68, 78
20–29	61.4	(56.7–65.8)	1.44	1.03–2.04		532, 533	56.6	(50.3–62.7)	2.11	1.13–3.93		298, 474
30 or older	53.7	(41.0–66.0)	1.05	0.58–1.91		68, 82	64.0	(49.8–76.1)	2.83	1.26–6.35		55, 85
Reproductive history												
Ever had a child					0.0452						0.3770	
Yes	57.0	(52.7–61.2)	1.00			683, 676	52.3	(46.0–58.6)	1.00			293, 471
No	58.7	(51.5–65.4)	1.14	0.81–1.59		239, 203	54.9	(46.6–62.9)	1.22	0.79–1.88		177, 220
Age at birth of first child ^b					<0.0001						0.0092	
<25	43.0	(36.1–50.1)	1.00			297, 245	32.3	(21.2–45.8)	1.00			74, 94
25–29	59.3	(51.5–66.7)	1.89	1.22–2.92		184, 183	48.9	(37.5–60.3)	1.82	0.86–3.87		89, 139
30–34	65.8	(56.5–74.0)	2.51	1.54–4.09		129, 148	68.0	(54.8–78.8)	4.14	1.85–9.24		68, 129
35 plus	74.6	(62.8–83.7)	3.73	1.99–6.99		71, 98	56.4	(43.0–68.8)	2.42	1.08–5.41		57, 109

Abortion ever					0.954						
No	57.4	(53.4–61.3)	1.00			788, 749					
Yes	56.3	(46.2–66.0)	0.99	0.64–1.52		131, 125					
Socio-economic position											
Educational level ^a					0.0001				0.0158		
Degree	65.5	(58.8–71.7)	1.00			242, 255	66.9	(57.4–75.1)	1.00	131, 228	
A-level/equivalent	58.0	(50.1–65.5)	0.73	0.47–1.12		202, 197	47.0	(37.8–56.5)	0.44	0.25–0.77	131, 198
GCSE, O-level or equivalent	55.6	(49.6–61.5)	0.64	0.44–0.94		360, 330	46.4	(37.1–56.0)	0.45	0.26–0.79	149, 189
Foreign or other	7.1	(0.9–38.1)	0.05	0.01–0.41		11, 8	32.8	(9.27–69.9)	0.27	0.05–1.45	7, 7
None	42.9	(32.9–53.5)	0.33	0.20–0.56		106, 89	45.8	(31.2–61.2)	0.40	0.19–0.86	52, 68
Social class (NS-SEC)					0.0069					0.0136	
Managerial and prof occupations	63.0	(56.9–68.6)	1.00			312, 338	60.2	(52.6–67.4)	1.00		195, 322
Intermediate occupations	62.2	(54.2–69.7)	0.96	0.63–1.46		201, 193	53.8	(41.8–65.4)	0.77	0.42–1.38	80, 116
Semi-routine/routine occupations	47.0	(39.8–54.2)	0.54	0.37–0.81		256, 209	42.8	(34.5–51.6)	0.53	0.33–0.85	161, 208
No job (10+ h/week) or not in the last 10 years	52.3	(42.7–61.8)	0.56	0.35–0.89		131, 123	37.0	(18.7–60.1)	0.34	0.13–0.91	24, 32
Student in full-time education	46.2	(24.0–70.0)	0.64	0.23–1.80		18, 13	78.2	(40.6–94.9)	2.85	0.59–13.90	8, 11
Index of multiple deprivation (IMD)—quintile					0.0583					0.0912	
1–2 (least deprived)	63.3	(57.7–68.6)	1.00			360, 383	60.5	(53.0–67.6)	1.00		193, 319
3	54.6	(46.1–62.8)	0.71	0.47–1.09		175, 171	47.1	(35.2–59.3)	0.60	0.34–1.09	79, 116
4–5 (most deprived)	51.8	(45.7–57.8)	0.67	0.47–0.94		388, 326	46.7	(38.8–54.9)	0.62	0.39–1.00	198, 256

Denominator: all those who reported experience of heterosexual sex and of infertility.

AOR, adjusted odds ratio; 95% CI, 95% confidence intervals.

^aParticipants aged 21 and older only.

^bOnly those who had a child.

Table III Associations between experience of infertility and aspects of current well-being and relationship quality among women aged 50 and younger at interview.^a

Women	Depression ^b			Dissatisfied with sex life ^c			Happy in relationship ^d			
	%	AOR	95% CI	P-Value	Denominators (unweighted, weighted)	%	AOR	95% CI	P-Value	Denominators (unweighted, weighted)
Experience of infertility				0.002					0.689	
No	11.4	1.00			5238, 3964	13.1	1.00			3018, 2561
Yes	15.3	1.63	1.20–2.21		667, 566	16.2	1.37	1.03–1.83		664, 563

Denominator: all women 50 years and younger with experience of heterosexual sex.

AOR, adjusted odds ratio; 95% CI, 95% confidence intervals.

^aadjusted for age at interview, highest educational qualification, relationship status at interview, duration of relationship, age at first cohabitation and whether or not a parent.^bPositive response to composite question on depressive symptoms in two weeks before interview.^cResponded 'disagree' or 'disagree strongly' with statement: 'I feel satisfied with my sex life'.^dResponded 1 or 2 to question: 'On a scale of 1–7, how happy or unhappy are you with your relationship with your partner, all things considered?' (1 = very happy and 7 = very unhappy).

this analysis. We found a positive and significant association between ever experience of infertility and symptoms of depression in the 2 weeks before interview and dissatisfaction with sex life in the past year among women aged 50 and under. These associations remained after controlling for potential confounding factors including age at interview, educational qualification, relationship status at interview, duration of relationship, age at first cohabitation and parental status. We examined whether having children or not moderated the association between experience of infertility and depression and identified no evidence of an interaction ($P = 0.698$). Given the limitations of the data, we do not know whether depression or sexual dissatisfaction was related to the use of fertility treatments. We found no significant association between women's experience of infertility and whether they were happy with their current relationship. Among men, no significant association was observed between the experience of infertility and depression, dissatisfaction with sex life, or happiness with relationship (analysis not shown).

Discussion

In this study, we provide population prevalence estimates of infertility and help seeking among women and men in Britain using national probability survey data. One in eight women and one in ten men experienced infertility defined by unsuccessfully attempting pregnancy for a year or more. Women and men who settled later with a partner, had higher educational attainment and occupational status and, among those who did have a child, became parents at older ages were more likely to have experienced infertility.

Little more than half of women and men who had experienced infertility had sought medical or professional help for the problem. Those who did so were better educated and in higher status occupations and, if they were parents, were more likely to have had children at older ages. These characteristics were more marked among women than men. Women aged 50 or younger who had experience of infertility were more likely to report recent symptoms of depression and dissatisfaction with their sex life. We found that ever experience of infertility and of help seeking were associated with few current health factors for women or men.

A strength of this study is the size of the sample and the fact that it is population-based. Natsal-3 includes data on sexual and reproductive health as well as behavioural and relationship variables less common in health surveys. A limitation is that, although we measure ever experience of infertility, some explanatory variables refer to recent time frames. As data are cross-sectional, it is not possible to establish temporality or infer causality and we cannot establish the age of participants when they experienced infertility or sought help, or the timing of these experiences relative to having children for those who did. Self-reported data may be subject to recall bias. It was not possible to explore associations between infertility and participant ethnicity as numbers were too small.

Our estimates of infertility are broadly in line with those found by previous studies although prevalence estimates differ as a result of diverse definitions and study populations. Earlier estimates of women experiencing infertility range from around 1 in 5 (Bhattacharya *et al.*, 2009; Cabrera-León *et al.*, 2015) to around 1 in 10 (Evers, 2002) with the National Women's Health Study reporting about 1 in 6 (Oakley *et al.*, 2008). Estimates of couples' experience of infertility also vary. Hull *et al.* (1985)

estimated it to be around one in six while NICE cites estimates of one in seven (National Institute for Health and Clinical Excellence, 2013).

A large body of literature describes the trend among women in developed countries to delay having children (Schmidt et al., 2012) and it is proposed that this changing fertility tempo is becoming a global phenomenon (Rosero-Bixby et al., 2009; Sobotka, 2010).

Our finding that infertility was more commonly experienced by married or cohabiting participants probably reflects the fact that those in stable relationships are more likely to have attempted pregnancy and therefore become aware of fertility problems. The much lower prevalence among young people reflects in part that they may never have tried to get pregnant. Previous studies have noted the extended period of transition to adulthood (Stone et al., 2014) and the widening intervals between the key reproductive events of first sex, first cohabitation and first birth (Wellings et al., 2013). Researchers taking a life course perspective (Morgan and Rackin, 2010; Berrington and Pattaro, 2014) have observed discrepancies between individuals' fertility intentions and their subsequent family size, highlighting the complex interaction of childhood socialization with personal and structural factors. Influences on postponement for women include the increase in women's participation in further education (Andersson et al., 2008; Ni Bhrolcháin and Beaujouan, 2012) and in career development (Martin, 2000; Morris et al., 2011), the absence of a 'suitable' partner (Proudfoot et al., 2009), partner's expectations (Iacovou and Tavares, 2011) and perceptions of how parenthood will reduce individual autonomy (Liefbroer, 2005).

Our estimate that 57% of women sought medical help for infertility is close to that reported by an international review which estimated that 56% of women in more developed countries sought help (Boivin et al., 2007), and by a Finnish study (Terävä et al., 2008) which found that 57% of all subfertile women did so. Other studies (Greil and McQuillan, 2004; Morris et al., 2011; Chandra et al., 2014) have reported higher and lower estimates but, because of differences in study groups and outcome measures, comparison is not possible.

Our analyses show that women aged 50 or under who experienced infertility were more likely to report recent symptoms of depression and dissatisfaction with their sex life. Infertility is associated with psychological distress (Cousineau and Domar, 2007; Greil et al., 2011) and patients find the process of undergoing infertility treatment and its uncertain outcome stressful (Sbaragli et al., 2008; Volgsten et al., 2008; Schmidt et al., 2013). The longer term impact of infertility on mental health and sexual well-being is less well known. A Danish study shows a relationship between unsuccessful ART and severe depressive symptoms a year after initiating treatment (Lund et al., 2009) and another, also Danish, found women who did not have a child after fertility treatment were more likely to commit suicide than those who did (Kjaer et al., 2011). Previous research has found associations between undergoing treatment for infertility and sexual dissatisfaction, particularly among women (Millheiser et al., 2012; Wischmann, 2010; Marci et al., 2012).

The large minority of research participants who experienced infertility but did not seek medical help is of concern, as are the marked inequalities in help seeking between those who are well qualified and in high status employment and those who are not. These findings are in line with other studies from Europe, North America and Australia (Terävä et al., 2008; Bushnik et al., 2012; Chambers et al., 2013; Chandra et al., 2014). Several explanations for not seeking (or pursuing) help for infertility have been suggested, including not understanding or acknowledging

that a problem exists (White et al., 2006), fear of being labelled infertile (Bunting and Boivin, 2007), concerns about the cost of treatment (Eisenberg et al., 2010), lack of intent to conceive (Greil and McQuillan, 2004) and the physical and psychological burden of treatment (Verberg et al., 2008). These do not provide a clear rationale for why there should be a distinction between indicators of social status and the likelihood of seeking help, although it has been suggested that acknowledging lack of conception as a problem to be solved is a motivation for seeking treatment and that highly educated women may be better informed about how long conception might typically take (Morris et al., 2011).

Interventions to encourage help seeking include raising public awareness about reproductive risks and strategies to minimise them (Macaluso et al., 2010), general practitioners taking opportunities to discuss fertility with patients (Davies, 2015), greater access to fertility treatments (Bunting and Boivin, 2007) and an acknowledgement of the psychosocial impacts of infertility, including the long-term effects, by health practitioners and the availability of appropriate support (Hinton et al., 2012).

Supplementary data

Supplementary data are available at <http://humrep.oxfordjournals.org/>.

Authors' roles

J.D., K.W., C.T. and M.P. conceived the paper; all authors contributed to data acquisition and interpretation; M.P., L.J.G. and K.G.J. conducted the statistical analysis; J.D. wrote the first draft with contributions from all other authors. All authors approved the final draft.

Funding

The study was funded by grants from the Medical Research Council and the Wellcome Trust with contributions from the Economic and Social Research Council and Department of Health. K.G.J. was supported by a National Institute for Health Research Methods Fellowship. Funding to pay the Open Access publication charges for this article was provided by MRC and Wellcome Trust.

Conflict of interest

AMJ is a Governor of the Wellcome Trust. Other authors declare no conflicts of interest.

References

- Andersson G, Rønsen M, Knudsen L, Lapppegård T, Neyer G, Skrede K, Teschner K, Vikat A. Cohort fertility patterns in the Nordic countries. MPIDR Working Papers. Rostok: Max Planck Institute for Demographic Research, 2008.
- Balash J. Ageing and infertility: an overview. *Gynecol Endocrinol* 2010; **26**:855–860.
- Bell AV. Beyond (financial) accessibility: inequalities within the medicalisation of infertility. *Sociol Health Illn*. 2010;**32**:631–646.
- Berrington A, Pattaro S. Educational differences in fertility desires, intentions and behaviour: a life course perspective. *Adv Life Course Res* 2014; **21**:10–27.

- Bhattacharya S, Porter M, Amalraj E, Templeton A, Hamilton M, Lee AJ, Kurinczuk JJ. The epidemiology of infertility in the North East of Scotland. *Hum Reprod* 2009;**24**:3096–3107.
- Boivin J, Bunting L, Collins JA, Nygren KG. International estimates of infertility prevalence and treatment-seeking: potential need and demand for infertility medical care. *Hum Reprod* 2007;**22**:1506–1512.
- Bunting L, Boivin J. Decision-making about seeking medical advice in an internet sample of women trying to get pregnant. *Hum Reprod* 2007;**22**:1662–1668.
- Bushnik T, Cook J, Hughes E, Tough S. Seeking medical help to conceive. *Health Reports/Statistics Canada, Canadian Centre for Health Information=Rapports sur la Santé/Statistique Canada, Centre Canadien D'information sur la Santé* 2012;**23**:7–13.
- Cabrera-León A, Lopez-Villaverde V, Rueda M, Moya-Garrido MN. Calibrated prevalence of infertility in 30- to 49-year-old women according to different approaches: a cross sectional population-based study. *Hum Reprod* 2015;**30**:2677–2685.
- Chambers GM, Hoang VP, Illingworth PJ. Socioeconomic disparities in access to ART treatment and the differential impact of a policy that increased consumer costs. *Hum Reprod* 2013;**28**:3111–3117.
- Chandra A, Copen CE, Stephen EH. Infertility service use in the United States: data from the National Survey of Family Growth, 1982–2010. *National Health Statistics Reports*. Hyattsville, MD: National Centre for Health Statistics, 2014.
- Cousineau TM, Domar AD. Psychological impact of infertility. *Best Pract Res Clin Obstet Gynaecol* 2007;**21**:293–308.
- Cui W. Mother or nothing: the agony of infertility. *Bull World Health Organ* 2010;**88**:881–882.
- Davies S. *Annual Report of the Chief Medical Officer 2014: the Health of the 51% Women*. London: Department of Health, 2015.
- Dhalwani NN, Fiaschi L, West J, Tata L. Occurrence of fertility problems presenting to primary care: population-level estimates of clinical burden and socioeconomic inequalities across the UK. *Hum Reprod* 2013;**28**:960–968.
- Eisenberg ML, Smith JF, Millstein SG, Nachtigall RD, Adler NE, Pasch LA, Katz PP. Predictors of not pursuing infertility treatment after an infertility diagnosis: examination of a prospective U.S. cohort. *Fertil Steril* 2010;**94**:2369–2371.
- Erens B, Phelps A, Clifton S, Mercer CH, Tanton C, Hussey D, Sonnenberg P, Maccowall W, Field N, Datta J *et al*. Methodology of the third British National Survey of Sexual Attitudes and Lifestyles (Natsal-3). *Sex Transm Infect* 2014;**90**:84–89.
- Evers J. Female subfertility. *Lancet* 2002;**360**:151–159.
- Greil A, McQuillan J. Help-seeking patterns among subfecund women. *J Reprod Infant Psychol* 2004;**22**:305–319.
- Greil AL, Shreffler KM, Schmidt L, McQuillan J. Variation in distress among women with infertility: evidence from a population-based sample. *Hum Reprod* 2011;**26**:2101–2112.
- Gurunath S, Pandian Z, Anderson RA, Bhattacharya S. Defining infertility—a systematic review of prevalence studies. *Hum Reprod Update* 2011;**17**:575–588.
- Gyimah SO. Women's educational attainment and the timing of parenthood in Ghana: a cohort perspective. *PSC Discussion Papers* 2003;**17**:Article 1.
- Hinton L, Kurinczuk JJ, Ziebland S. Reassured or fobbed off? Perspectives on infertility consultations in primary care: a qualitative study. *Br J Gen Pract* 2012. doi:10.3399/bjgp12X649133.
- Hull MGR, Glazener CMA, Kelly NJ, Conway DI, Foster PA, Hinton RA, Coulson C, Lambert PA, Watt EM, Desai KM. Population study of causes, treatment, and outcome of infertility. *BMJ* 1985;**291**:1693–1697.
- Iacovou M, Tavares L. Yearning, learning, and conceding: reasons men and women change their childbearing intentions. *Popul Dev Rev* 2011;**37**:89–123.
- Kjaer TK, Jensen A, Dalton SO, Johansen C, Schmiedel S, Kjaer SK. Suicide in Danish women evaluated for fertility problems. *Hum Reprod* 2011;**26**:2401–2407.
- Kroenke K, Spitzer RL, Williams JBW. The Patient Health Questionnaire-2: validity of a two-item depression screener. *Med Care* 2003;**41**:1284–1292.
- Leridon H. A new estimate of permanent sterility by age: sterility defined as the inability to conceive. *Popul Stud* 2008;**62**:15–24.
- Liefbroer AC. The impact of perceived costs and rewards of childbearing on entry into parenthood: evidence from a panel study. *Eur J Popul* 2005;**21**:367–391.
- Lund R, Sejbaek CS, Christensen U, Schmidt L. The impact of social relations on the incidence of severe depressive symptoms among infertile women and men. *Hum Reprod* 2009;**24**:2810–2820.
- Macaluso M, Wright-Schnapp TJ, Chandra A, Johnson R, Satterwhite CL, Pulver A, Berman SM, Wang RY, Farr SL, Pollack LA. A public health focus on infertility prevention, detection, and management. *Fertil Steril* 2010;**93**:16.e11–16.e10.
- Marci R, Graziano A, Piva I, Lo Monte G, Soave I, Giugliano E, Mazzoni S, Capucci R, Carbonara M, Caracciolo S *et al*. Procreative sex in infertile couples: the decay of pleasure? *Health Qual Life Outcomes* 2012;**10**:140.
- Martin S. Diverging fertility among U.S. women who delay childbearing past age 30. *Demography* 2000;**37**:523–533.
- Mascarenhas MN, Flaxman SR, Boerma T, Vanderpoel S, Stevens GA. National, Regional, and Global Trends in Infertility Prevalence Since 1990: A Systematic Analysis of 277 Health Surveys. *PLoS Med* 2012;**9**:e1001356.
- Mathews T, Hamilton B. Delayed childbearing: more women are having their first child later in life NCHS Data Brief. US Department of Health and Human Services, 2009.
- Millheiser LS, Helmer AE, Quintero RB, Westphal LM, Milki AA, Lathi RB. Is infertility a risk factor for female sexual dysfunction? A case-control study. *Fertil Steril* 2012;**94**:2022–2025.
- Mills M, Rindfuss RR, McDonald P, te Velde E, ESHRE Reproduction and Society Task Force. Why do people postpone parenthood? Reasons and social policy incentives. *Hum Reprod Update* 2011;**17**:848–860.
- Morgan SP, Rackin H. The correspondence between fertility intentions and behaviour in the United States. *Popul Dev Rev* 2010;**36**:91–118.
- Morris M, Oakley L, Maconochie N, Doyle P. An investigation of social inequalities in help-seeking and use of health services for fertility problems in a population-based sample of UK women. *Hum Fertil* 2011;**14**:16–22.
- National Institute for Health and Clinical Excellence. *Fertility: Assessment and Treatment for People with Fertility Problems*. NICE Clinical Guideline. Manchester, 2013.
- Ní Bhrolcháin M, Beaujouan É. Fertility postponement is largely due to rising educational enrolment. *Popul Stud* 2012;**66**:311–327.
- Oakley L, Doyle P, Maconochie N. Lifetime prevalence of infertility and infertility treatment in the UK: results from a population-based survey of reproduction. *Hum Reprod* 2008;**23**:447–450.
- Office for National Statistics. *Standard Occupational Classification 2010*. Basingstoke: Palgrave Macmillan, 2010.
- Proudfoot S, Wellings K, Glasier A. Analysis why nulliparous women over age 33 wish to use contraception. *Contraception* 2009;**79**:98–104.
- Rosero-Bixby L, Castro-Martín T, Martín-García T. Is Latin America starting to retreat from early and universal childbearing? *Demogr Res* 2009;**20**:169–194.
- Sbaragli C, Morgante G, Goracci A, Hofkens T, De Leo V, Castrogiovanni P. Infertility and psychiatric morbidity. *Fertil Steril* 2008;**90**:2107–2111.
- Schmidt L, Sobotka T, Bentzen JG, Nyboe Andersen A. Demographic and medical consequences of the postponement of parenthood. *Hum Reprod Update* 2012;**18**:29–43.
- Schmidt L, Hageman I, Hougaard CØ, Sejbaek CS, Assens M, Ebdrup NH, Pinborg A. Psychiatric disorders among women and men in assisted reproductive technology (ART) treatment. The Danish National

- ART-Couple (DANAC) cohort: protocol for a longitudinal, national register-based cohort study. *BMJ Open* 2013;**3**:e002519. doi:10.1136/bmjopen-2012-002519.
- Schneider MG, Forthofer MS. Associations of psychosocial factors with the stress of infertility treatment. *Health Soc Work* 2005;**30**:183–191.
- Sobotka T. In pursuit of higher education, do we postpone parenthood too long? *Gender Med* 2006;**3**:183–186.
- Sobotka T. Shifting parenthood to advanced reproductive ages: trends, causes and consequences. In: Chet Tremmel J (ed). *A Young Generation Under Pressure*. London, UK: Springer, 2010, pp. 129–154.
- StataCorp. *Stata Statistical Software: Release 13*. College Station: StataCorp LP, 2013.
- Stone J, Berrington A, Falkingham J. Gender, turning points, and boomerangs: returning home in young adulthood in Great Britain. *Demography* 2014;**51**:257–276.
- Terävä A-N, Gissler M, Hemminki E, Luoto R. Infertility and the use of infertility treatments in Finland: Prevalence and socio-demographic determinants 1992–2004. *Eur J Obst Gynecol Reprod Biol* 2008;**136**:61–66.
- Verberg MFG, Eijkemans MJC, Heijnen EMEW, Broekmans FJ, de Klerk C, Fauser BCJM, Macklon NS. Why do couples drop-out from IVF treatment? A prospective cohort study. *Hum Reprod* 2008;**23**:2050–2055.
- Volgsten H, Skoog Svanberg A, Ekselius L, Lundkvist Ö, Sundström Poromaa I. Prevalence of psychiatric disorders in infertile women and men undergoing in vitro fertilization treatment. *Hum Reprod* 2008;**23**:2056–2063.
- Wellings K, Jones KG, Mercer CH, Tanton C, Clifton S, Datta J, Copas AJ, Erens B, Gibson LJ, Macdowall W et al. The prevalence of unplanned pregnancy and associated factors in Britain: findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3). *Lancet* 2013;**382**:1807–1816.
- White L, McQuillan J, Greil AL, Johnson DR. Infertility: testing a helpseeking model. *Soc Sci Med* 2006;**62**:1031–1041.
- Wilkes S, Chinn DJ, Murdoch A, Rubin G. Epidemiology and management of infertility: a population-based study in UK primary care. *Fam Pract* 2009;**26**:269–274.
- Wischmann TH. Couples' sexual dysfunctions: sexual disorders in infertile couples. *J Sex Med* 2010;**7**:1868–1876.
- Zegers-Hochschild F, Adamson G, de Mouzon J, Ishihara O, Mansour R, Nygren K, Sullivan E, Vanderpoel S, for ICMART and WHO. International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) revised glossary of ART terminology, 2009. *Fertil Steril* 2009;**92**:1520–1524.