

Title:

Is cancer survival associated with cancer symptom awareness and barriers to seeking medical help in England? An ecological study

Running title:

Cancer awareness and cancer survival in England

Authors:

Maja Niksic (corresponding author),¹ Bernard Rachet,² Stephen W Duffy,³ Manuela Quaresma,² Henrik Møller,¹ Lindsay JL Forbes¹

Affiliations:

¹ Cancer Epidemiology and Population Health, Cancer Studies Research Division, King's College London, Bermondsey Wing, 3rd floor, Guy's Hospital, Great Maze Pond, London, SE1 9RT, UK

² Cancer Research UK Cancer Survival Group, Department of Non-Communicable Disease Epidemiology, Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT, UK

³ Centre for Cancer Prevention, Wolfson Institute of Preventive Medicine, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, Charterhouse Square, London, EC1M 6BQ, UK

Corresponding author: Maja Niksic (email: maja.niksic@kcl.ac.uk); Cancer Epidemiology and Population Health, Cancer Studies Research Division, King's College London, Bermondsey Wing, 3rd floor, Guy's Hospital, Great Maze Pond, London, SE1 9RT, UK

tel: +44(0) 207 927 2551

fax:+44(0) 207 436 4230

The order of contribution:

- 1) Ms Maja Niksic, PhD candidate, MPhil, MSc Public Health
(<https://kclpure.kcl.ac.uk/portal/en/persons/maja-niksic%28b42d23a5-83b8-4b88-8572-4b76fb64fe27%29.html>)
- 2) Dr Bernard Rachet, MD, PhD, FFPH
email: bernard.rachet@lshtm.ac.uk
(<http://www.lshtm.ac.uk/aboutus/people/rachet.bernard>)
- 3) Dr Lindsay JL Forbes, MD, MSc, MRCP, FFPH
email: lindsay.forbes@kcl.ac.uk
(<https://kclpure.kcl.ac.uk/portal/lindsay.forbes.html>)
- 4) Ms Manuela Quaresma, MSc, PhD student
email: manuela.quaresma@lshtm.ac.uk
(<http://www.lshtm.ac.uk/aboutus/people/quaresma.manuela>)
- 5) Prof Stephen W Duffy, BSc, MSc, CStat
email: s.w.duffy@qmul.ac.uk
(<http://www.wolfson.qmul.ac.uk/a-z-staff-profiles/stephen-duffy>)
- 6) Prof Henrik Møller, PhD
email: henrik.moller@kcl.ac.uk
(<https://kclpure.kcl.ac.uk/portal/henrik.moller.html>)

Abstract

Background: Campaigns aimed at raising cancer awareness and encouraging early presentation have been implemented in England. However, little is known about whether people with low cancer awareness and increased barriers to seeking medical help have worse cancer survival, and whether there is a geographical variation in cancer awareness and barriers in England.

Methods: From population-based surveys (n=35,308), using the Cancer Research UK Cancer Awareness Measure, we calculated the age- and sex -standardised symptom awareness and barriers scores for 52 Primary Care Trusts (PCTs). These measures were evaluated in relation to the sex-, age-, and type of cancer - standardised cancer survival index of the corresponding PCT, from the National Cancer Registry, using linear regression. Breast, lung, and bowel cancer survival were analysed separately.

Results: Cancer symptom awareness and barriers scores varied greatly between geographical regions in England, with the worst scores observed in socio-economically deprived parts of East London. Low cancer awareness score was associated with poor cancer survival at PCT level (estimated slope=1.56, 95%CI: 0.56; 2.57). The barriers score was not associated with overall cancer survival, but it was associated with breast cancer survival (estimated slope=-0.66, 95%CI: -1.20; -0.11). Specific barriers, such as embarrassment and difficulties in arranging transport to the doctor's surgery, were associated with worse breast cancer survival.

Conclusions: Cancer symptom awareness and cancer survival are associated. Campaigns should focus on improving awareness about cancer symptoms, especially in socio-economically deprived areas. Efforts should be made to alleviate barriers to seeking medical help in women with symptoms of breast cancer.

Keywords: cancer survival; cancer awareness; barriers; healthcare seeking; England

INTRODUCTION

Cancer accounts for more than 130,000 deaths each year in England (Office for National Statistics, 2012a). Survival for most types of cancer is lower in England than in comparable Western European countries (Berrino *et al*, 2007), and in other high-income countries, such as Canada and Australia (Coleman *et al*, 2011). The National Health Service (NHS) in England has introduced several cancer plans, highlighting early diagnosis as a key strategy to improve cancer survival (Department of Health, 2000; Department of Health, 2011; Department of Health, 2014). Since *Improving Outcomes: A Strategy for Cancer* in 2011, more than £450 million has been invested in raising cancer awareness and encouraging early diagnosis in England (Department of Health, 2014). A number of 'Be Clear on Cancer' campaigns were developed to improve recognition of early symptoms and encourage prompt visits to the doctor for the most common types of cancer, such as breast, lung and bowel cancer (Cancer Research UK, 2015). The underlying assumption was that survival would be improved if people recognised symptoms early, and sought medical help before developing advanced-stage cancer. Short-term survival is particularly sensitive to stage of disease and therefore to timeliness of cancer diagnosis.

It is not clear what effect, if any, cancer awareness and barriers to help-seeking have on cancer survival. The most appropriate research design to address this question would be a prospective, cohort study examining the predictive value of symptom recognition or barriers to help-seeking on cancer survival. Such a study has not yet been carried out. It is also difficult to assess the effects of intermediary factors, such as time from the onset of symptoms to presentation, time from presentation to starting a treatment, stage of disease, and types of treatment. Ironmonger *et al* (2014) evaluated the first national campaign and a regional campaign in England, both aimed at raising public awareness of a persistent cough as a lung cancer symptom. They reported an increase in cancer awareness, a shift to earlier stage at diagnosis, and a higher surgical resection rate after both campaigns, compared with one year before. Data on survival were available only for the regional campaign, and suggested a statistically non-significant increase in one-year survival in the pilot compared with control areas.

Evidence suggests a wide geographical variation in cancer survival in England, with lower survival in the north compared to the south of the country (Quaresma *et al*, 2011; Walters *et al*, 2011). Quaresma *et al* (2011) showed that the geographical variation in one-year cancer survival at Primary Care Trust (PCT) level is constant year-on-year, from 1996 to 2009. PCTs were the statutory NHS bodies, structured as small geographical areas, responsible for commissioning and providing most health services in England until 2013 (NHS, 2012). Walters

et al (2011) reported the same geographical patterns, with generally lower cancer survival in northern England, for patients diagnosed between 1991 and 2006, using Cancer Network areas. These studies analysed survival from the most common types of cancer, such as lung, breast and bowel.

To explore reasons for variation in cancer survival between countries with similar health care systems, including Australia, Canada, Denmark, Norway, Sweden, and the UK, Forbes *et al* (2013) examined differences in cancer awareness and barriers to help-seeking. While the UK, with the lowest cancer survival, had low average cancer awareness and the highest barriers, there was no consistent pattern across the participating countries. For example, Denmark, also with poor cancer survival, had low levels of barriers. However, it may be difficult to detect any patterns at an international level. The assessment of differences in cancer awareness and barriers by much smaller geographical region in a single country may offer a more salient explanation of cancer survival differences. This study aimed to identify how cancer symptom awareness and barriers to help-seeking vary by small geographical region (PCT) in England, and whether average levels of awareness and barriers are associated with cancer survival at PCT level.

METHODS

Material

Data from population-based surveys, using the Cancer Research UK Cancer Awareness Measure across England (n=49,270), were linked with cancer survival data from the National Cancer Registry at the Office for National Statistics. These two datasets were linked based on matching by the relevant geographical area (PCT) in England. To make the matching procedure more reliable we used cancer survival data of patients diagnosed in 2010 only, because most CAM surveys were conducted in that year (range 2009 to 2011). This meant that both patients and survey participants would have been exposed to the same media health messages and local cancer awareness campaigns, delivered in their area of residence up to 2010.

On one side, we used data from the surveys, described in detail elsewhere (Niksic *et al*, 2015), carried out in 92 PCTs out of the 151 PCTs in England at the time. The response rate was 51%. The surveys questions used to assess cancer symptom awareness (summed up recognition of nine cancer symptoms) and barriers to help-seeking (summed up reporting of ten barriers to help-seeking) are available in Supplement 1.

To allow reliable calculations of awareness and barriers scores, we included only 52 PCTs, which had at least 90 participants in total, with available data in each sex and age category (15-44; 45-54; 55-64; 65-74; and 75+ years). To calculate the symptom awareness score, for each participant we added up the positive responses ('Yes') to the questions about the recognition of nine cancer symptoms (Niksic *et al*, 2015). We then calculated the arithmetic mean of this score for each combination of PCT, sex and age group. We took the same approach with calculating the barriers score, using positive responses ('Yes often' or 'Yes sometimes') to questions about ten barriers to help-seeking. To standardise the awareness and barriers scores by age and sex, we used the age and sex weights developed by the Cancer Research UK Cancer Survival Group, at London School of Hygiene and Tropical Medicine. The same weights were also used for the calculation of cancer survival index (and site-specific indices). The only difference between these weights was that cancer survival indices were also weighted for the type of cancer. This made the comparison between the awareness and barriers score and cancer survival indices more reliable. It also allowed us to avoid using the same weights for the elderly and the young people living in each PCT, who have different risks of being diagnosed with cancer and different cancer survival. The same approach was used with calculating individual symptoms and barriers.

On the other side, we used data from the National Cancer Registry to calculate net cancer survival. Net survival was defined as survival from a cancer of interest in the absence of death from other causes, estimated from population life tables (Perme *et al*, 2012). The life tables were stratified by every year of age, sex and socio-economic deprivation categories for the calendar years 2010 to 2011, and developed by Cancer Research UK Cancer Survival Group, at the London School of Hygiene and Tropical Medicine. The survival index was calculated as one-year net survival for all cancers combined, except non-melanoma skin and prostate cancer, standardised for type of cancer, sex, and age, in five age categories (15-44; 45-54; 55-64; 65-74; and 75+years) for patients diagnosed in 2010 and followed up until 2011. Patients diagnosed with non-melanoma skin cancer were excluded because these malignancies are rarely fatal and extremely rarely linked with metastases (NCIN, 2013). Registries in the UK have incomplete data on non-melanoma skin cancer, which is often underreported, and many patients are treated topically without pathological verification and in the private sector (Goodwin *et al*, 2004; NCIN, 2013). Prostate cancer was excluded because the widespread introduction of prostate-specific antigen (PSA) testing since the early 1990s introduced difficulties in the interpretation of survival trends (Pashayan *et al*, 2006). The exclusion of non-melanoma skin cancer and prostate cancer is the common procedure in calculation of the cancer survival index in England, published annually by the ONS (Office for National Statistics, 2012b).

This index was designed to support the evaluation of the effects of early diagnosis, and to provide a single summary measure of cancer survival, for the purpose of monitoring the effectiveness of the health care system (Quaresma *et al*, 2015). Breast (women only), bowel and lung cancer survival indices were calculated using the same approach. These types of cancer were selected because they are associated with the highest incidence and the highest mortality in Europe (Ferlay *et al*, 2013).

The total number of cancer patients used to construct the survival index and site-specific indices was 133,413, which was the number of residents in PCTs included in our analysis who were diagnosed with cancer in 2010. Out of this number 13,475 (10%) were female breast cancer patients, 21,902 (16%) were bowel cancer patients, 21,510 (n=16%) were lung cancer patients, and 76,526 (57%) of patients diagnosed with other types of cancer. The cancer survival indices at PCT level, were linked to the cancer awareness and barriers measures by the corresponding PCT.

Statistical analysis

The awareness and barriers scores were analysed as continuous variables, and their association with the cancer survival index was assessed using scatter plots and linear regression (at $p < 0.05$ level). The survival index was the dependent variable. Scatter plots were also used to visually assess the association between breast, bowel, and lung cancer survival, and their cancer-specific symptoms, including 'unexplained lump or swelling', 'persistent cough or hoarseness', and 'change in bowel or bladder habits', respectively. Associations between cancer survival indices and individual cancer symptoms, and individual barriers were also assessed using linear regression. In regression models, each PCT was weighted using the inverse variance of the cancer survival estimate. We defined different levels of the barriers score based on the interquartile range (IQR). We considered barriers below 25th percentile as 'low', between 25th and 75th percentile as 'intermediate' barriers, and above 75th percentile as 'high' barriers. The same procedure was used for the awareness score. Maps were also developed using 25th and 75th percentiles to determine the cut-off points for low, intermediate and high scores. We calculated the statistical power to detect the effects in our regression analysis using a post-hoc power calculator (<http://clincalc.com/Stats/Power.aspx>), and reliability of cancer awareness and barriers scores between PCTs, using one-way ANOVA. We analysed data using Stata 14.0 (STATA Corp, 2015) and ArcGIS for Desktop Advanced 10.3 software (Environmental Systems Research Institute, 2014).

RESULTS

Our dataset included 52 PCTs spread across England (Table 1). The mean cancer symptom awareness score was 7.2 (range: 4.8-8.3, the interquartile range (IQR): 6.9, 7.6). The mean barriers to help-seeking score was 1.7 (range: 0.6-3.1, IQR: 1.4, 1.8). One-year cancer survival index had a mean value of 66.9 (range 60.5 to 71.7).

Table 1: (About here)

Geographical variation in cancer awareness and barriers to help-seeking in England

The lowest cancer awareness scores were observed in Tower Hamlets (4.8), Newham (5.5), Redbridge (6.4), Gloucestershire (6.5), and Lambeth (6.5). The highest awareness scores were observed in Peterborough (8.3), Bedfordshire (8.1), Great Yarmouth and Waveney (8.1), and Cambridgeshire (8.0) (Figure 1).

In England different PCT were ranked according to their socio-economic deprivation, based on the income domain of the English indices of Deprivation (Department for Communities and Local Government, 2011). PCTs ranked as some of the most socio-economically deprived were: Newham and Tower Hamlets in East London, Bradford and Airedale and Kirklees in Yorkshire, Lambeth and City and Hackney in London, and Hampshire in Southern coast of England. PCTs ranked as some of the least socio-economically deprived were: Herefordshire in West Midlands, Bedfordshire in the East of England, Richmond and Twickenham in London, and Great Yarmouth and Waveney in the East of England.

Figure 1: (About here)

The highest barriers scores observed in Tower Hamlets (3.1), Kirklees (3.1), Bradford and Airedale (3.0), and Newham (2.7). The lowest barriers scores were observed in North Tyneside (0.6), Northamptonshire (0.8), Newcastle (0.9), and Herefordshire (1.1) (Figure 2).

Figure 2: (About here)

Cancer awareness and barriers scores in relation to cancer survival

There were large differences between survival from different types of cancer – breast cancer survival ranged from 93% to 99%, bowel cancer survival ranged from 67% to 81%, and lung cancer survival ranged from 18% to 42%.

Scatter plots showed the position of each PCT with regard to their cancer survival and cancer awareness score, or recognition of cancer-specific symptoms (Figure 3); and, cancer survival and barriers score (Figure 4). There was a considerable variation in assessed variables across PCT areas. We observed a trend with the best scores in PCTs such as Bedfordshire and Cambridgeshire, and the worst scores in Tower Hamlets and Newham PCTs. The outliers in most scatter plots were Tower Hamlets and Newham PCTs, but even after excluding these outliers, similar trends remained in the scatter plots.

Figure 3: (About here)

Figure 4: (About here)

The association between the cancer awareness score and the cancer survival index was statistically significant (estimated slope=1.56, 95%CI: 0.56; 2.57, $p=0.01$) (Table 2). Each additional symptom recognised was associated with a 1.56% increase in the cancer survival index. Recognition of seven out of nine individual cancer symptoms was statistically significantly associated with a slight increase in cancer survival index (Table 2). For example, for every 1% increase in the recognition of unexplained lump or swallowing among people living in these PCTs there was a 0.15 increase in cancer survival index (Table 2). For every 1% increase in the recognition of unexplained weight loss among people living in these PCTs there was a 0.13 increase in cancer survival index (Table 2). We found no trends (Figure 3) or statistically significant associations (Table 2) between survival from breast, lung, or bowel cancer, and the awareness score or recognition of each cancer symptom.

Table 2: (About here)

The barriers score was not associated with the cancer survival index (estimated slope=-0.14, 95% CI: -1.28; 0.99, $p=0.80$), but it was associated with breast cancer survival: each additional barrier reported was associated with 0.66% decrease in the breast cancer survival (estimated slope=-0.66, 95%CI: -1.20; -0.11, $p=0.02$) (Table 2). Most individual barriers had an inverse association with breast cancer survival. We found no statistically significant associations

between the barriers score or individual barriers, and lung and bowel cancer survival. Similar statistically significant associations were obtained after we excluded from regression models the two outliers, Tower Hamlets and Newham PCTs (data available on request).

Discussion

We found geographical variation in symptom awareness and barriers to help-seeking across England, with the lowest scores observed in socio-economically deprived parts of East London. Low cancer symptom awareness was associated with poor cancer survival index for all cancers combined, and high barriers to seeking medical help were associated with poor survival from breast cancer, at PCT level in England.

Strengths and limitations

This is the first study to examine geographical variation in cancer awareness and barriers to help-seeking in England, and to link these with cancer survival data. All measures, predictors and outcomes, were standardised using the same age- and sex standardised weights. This means that the cancer awareness and barriers score in each PCT were weighted differently depending on the age and sex structure of the survey population in each PCT. The cancer survival indices were standardised by age, sex, and type of cancer in each PCT, to adjust for changes over time in the distribution of cancer patients in these categories. This approach was necessary, because cancer survival varies widely depending on these factors (Quaresma *et al*, 2012). Therefore, the cancer survival indices were not affected by changes over time in the proportion of cancers of different lethality by sex, such as an increase in breast cancer and a reduction of lung cancer incidence. In addition, the survival indices were not affected by different age distribution of the cancer patient population, or a shift in the proportion of men and women diagnosed with a given type of cancer (Office for National Statistics, 2012c; Quaresma *et al*, 2012). We analysed only PCTs with 90+ participants, and available data in all sex and age groups, to allow reliable calculations of the scores. A validated measure of public cancer awareness (Stubbings *et al*, 2009) was used to interview participants included in a large population-based sample.

The matching of surveys with survival data of patients diagnosed in 2010 when most of the surveys were carried out, reduced a possible bias that would happen if patients had more opportunities to improve their cancer awareness than survey participants. For example, several regional campaigns were run in England during 2011, and the first large national campaigns, aimed at improving awareness about lung and bowel cancer symptoms were introduced in 2012 (Ironmonger *et al*, 2014). To the best of our knowledge there were no large-

scale awareness raising campaigns in England prior to 2010. In fact, most of the CAM questionnaires were used by cancer networks as a preliminary assessment prior to launching various cancer awareness campaigns. We used a one-year cancer survival, which is the most commonly used proxy measure for late presentation (Møller *et al*, 2011). This is because most excess deaths in the English patients diagnosed with breast (Møller *et al*, 2010), lung (Cheyne *et al*, 2013) and bowel cancer (Møller *et al*, 2012) occur in the first year after diagnosis. Our study provides some support to the NAEDI hypothesis, which proposed that the main reasons behind late diagnosis, associated with poor survival, are low cancer awareness and increased barriers to help-seeking (Hiom, 2015; Richards, 2009).

As with any ecological study, our study may be subject to what has been called the ecological fallacy, which can occur if we assume that inferences based on group-level analyses are applicable at the individual level (Piantadosi *et al*, 1988). For example, it would be incorrect to assume that by simply increasing awareness about an additional cancer symptom would mean that a person can expect an increase in cancer survival by 1.65 percentage point (if and when cancer is diagnosed). If a woman no longer feels embarrassment in relation to changes in her breasts, following a campaign, it does not mean that her expected survival from breast cancer will be automatically increased by 0.6 percentage points. In addition, the possibility that these associations may be stronger or weaker at the level of individual cancer patients cannot be excluded. Although our findings are not directly applicable to individual people, they outlined an overall pattern of low cancer survival in small geographical areas with low cancer symptom awareness. As such, our findings can be used as a starting point in investigations of the complex relationship between symptom awareness, barriers to help-seeking, and cancer survival. This is essential prior to embarking on a large and complex investigation based on individual-level data. Marmot (1998) argues that ecological analyses are a useful way to explore the relationship between social environment and health, particularly as effective policies are delivered at a community-wide rather than individual-level.

Our results might not be generalised nationally, because we included only a third of all PCTs. However, we included a wide range of environmentally and socio-economically diverse PCTs across the country. We have not adjusted our analyses for socio-economic deprivation or ethnicity, for a number of reasons. First, at the time when we conducted the analysis survival data of different ethnic groups by PCT were not available to us. Therefore, it was not possible to include this factor in our analysis. Second, our units of analysis were PCTs, and there was a wide variation of socio-economic deprivation within each PCT. Calculating a single deprivation score for each PCT may not be valid. This is because on average there are about

100,000 people living in each PCT, and some PCTs had over 250,000 residents such as Brighton and Hove (Bojke *et al*, 2001). Adjusting the PCTs for their average socio-economic status may not accurately capture material or social resources, especially among culturally and ethnically diverse groups. Most large cities in England, such as London, Manchester or Lancaster have some poor people living in affluent areas, and *vice versa*, with contrasting housing types in the same neighbourhood, ranging from Victorian semi-detached houses and maisonettes, to less popular council flats and 'towers' (Cattell, 2001; Popay *et al*, 2003). Therefore, adjusting each geographical area for its average level of socio-economic deprivation is not warranted, especially if there are large gaps between the affluent and poor residents in that area.

Third, our surveys had a substantial proportion of participants with missing data on socio-economic deprivation. If we had decided to use data on socio-economic deprivation to calculate the cancer awareness and barriers scores the number of units in our analysis would be reduced to 22 only. This is because we would have to remove 30 PCTs, because they did not have participants in all five categories of socio-economic deprivation. However, we used population life tables that may in part overcome the fact that socio-economic deprivation was not accounted for. These life tables took into account different mortality rates of a different population groups living in these PCT, during the year 2010 and 2011, and, for example, controlled for the fact that the elderly men, with low socio-economic position, have high mortality rates. In addition, the use of age- and sex-standardised weights also accounted for some of the differences in socio-demographic distribution of people living in these PCTs.

With statistical power of 80% we were able to detect regression coefficients over 1.44 for cancer awareness and over 0.36 for barriers score in relation to the overall cancer survival index. For breast cancer survival, with statistical power of 80% we were able to detect regression coefficients over 0.79 for cancer awareness and over 0.17 for barriers score. For lung cancer survival, with statistical power of 80% we were able to detect regression coefficients over 2.76 for cancer awareness and over 0.62 for barriers score. For bowel cancer survival, with statistical power of 80% we were able to detect regression coefficients over 2.08 for cancer awareness and over 0.50 for barriers score. It is possible that some of our results were not statistically significant because of lack of power, rather than no association. However, it is worth noting that the weighting in the regression model to some extent mitigates the lack of power.

It is important to note that the scores were standardised by age group and sex, which would reduce the within-PCT variability. The reliability of the group-level barriers scores was at around 0.80 for each age and sex combination, with the exception of the 75+ age group (0.23

in men and 0.50 in women). For awareness scores, their reliability was above 0.80 for each age and sex combination, with the exception of elderly men (0.55) and elderly women (0.53). The relatively low reliability among elderly is related to the small number of participants in the 75+ age group. This suggests an overall reliability of the PCT-level scores standardised on age and sex.

Comparison with existing literature

Cancer symptom awareness and barriers to help-seeking varied greatly between geographical areas in England, but we did not observe a north-south divide in these scores. The worst scores were observed in socio-economically deprived parts of East London, such as Tower Hamlets and Newham. There is ample evidence that socio-economically deprived people in England tend to have low cancer awareness and numerous reasons to deter them from seeking medical help, including embarrassment or difficulties in arranging transport to the doctor's surgery (Niksic *et al*, 2015; Robb *et al*, 2009; Waller *et al*, 2009). A recent Danish study corroborated the findings that socio-economic deprivation is strongly associated with barriers to help-seeking (Hvidberg *et al*, 2014). Underprivileged people are also more likely to pursue an unhealthy lifestyle, including smoking and low physical activity (Marmot *et al*, 1991), which act to 'normalise' early symptoms of disease (Dixon-Woods *et al*, 2006). They may also feel embarrassed to discuss their symptoms with their doctor, due to perceived social distance and status differences.

Another possible explanation for our findings is the role of ethnic minority background, which is one of the factors contributing to poor recognition of symptoms and increased barriers to help-seeking (Forbes *et al*, 2011; Niksic *et al*, 2016). A recent study found that ethnic minorities in England had lower cancer symptom awareness and more widespread barriers to seeking medical help than White participants (Niksic *et al*, 2016). Another study carried out in East London also reported that South Asian and Black women had significantly lower breast cancer awareness than White women (Forbes *et al*, 2011). Emotional barriers to seeking medical help, such as embarrassment and lack of confidence to talk to the doctor, were highest among South Asian women. These differences could not be explained by different levels of educational attainment or socio-economic deprivation. Socio-economically deprived areas often have higher proportion of ethnic minorities, and these two factors in combination may have an amplified effect on cancer awareness and barriers to help-seeking.

Low cancer awareness was associated with low cancer survival at PCT level, with the worst results observed in Tower Hamlets and Newham. These boroughs of Inner North-East London, together with Hackney, have been described as 'the unhealthiest place in Britain',

with high levels of unemployment, large ethnic diversity, low levels of social cohesion and the largest population growth in the country (East London NHS Trust, 2013). People living in the same PCT area are likely to share aspects of their social and lifestyle patterns, and access to educational and healthcare facilities, making their health outcomes more similar than those of people from different PCTs (Nur *et al*, 2015). A term 'place effect' was coined to describe the extent of area effects on individuals' health (Macintyre *et al*, 2002). If symptom awareness and cancer survival are to be improved, then it can be argued that place effects need to be taken into account. Campaigns may be more effective if they are culturally sensitive and socially inclusive, with a clear and simple message, translated into languages frequently spoken in the community, and if healthcare provision, in terms of both units and personnel, are increased in areas with rapidly growing population. Socio-economically deprived areas, which have poor cancer survival (Coleman *et al*, 2004; Rachet *et al*, 2010), could benefit most from awareness rising campaigns. A recent NHS Health and Wellbeing Strategy is an example of the area-specific approach to improve health outcomes, which includes reducing the risks of developing cancer (East London NHS Trust, 2013).

We found no strong evidence that recognition of the cancer-specific symptoms were associated with survival from the corresponding cancers. People may perceive these symptoms, such as 'persistent cough', as temporary and attribute them to non-cancer causes like smoking, having a cold or an allergy. As an alternative explanation, the role of 'symptom clusters' has also been implicated. Early symptoms of breast, bowel or lung cancer often occur in clusters rather than in isolation. A symptom cluster is a relatively stable group of two or more related symptoms, occurring simultaneously and independently from other symptom clusters (Kim *et al*, 2005). For example, the cluster of lung cancer symptoms may include thoracic pain, cough, hemoptysis, and vomiting (Hamilton *et al*, 2005a). The cluster of bowel cancer symptoms may include rectal bleeding, abdominal pain, diarrhea, and weight loss (Hamilton *et al*, 2005b). The cluster of breast cancer symptoms may include a lump, fatigue, pain, and insomnia (Banning, 2007; Gaston-Johansson *et al*, 1999). Recognition of a single, cancer-specific symptom may not be sufficient to produce an effect on survival from that cancer. Experiencing a few unexplained symptoms simultaneously, as a cluster, is perhaps more likely to have an effect on cancer survival, because it may prompt people to seek medical help.

The barriers score was not associated with the cancer survival index, but it was associated with breast cancer survival. This could be because breast cancer survival was assessed for women only, and there is evidence that women are more likely to report barriers to help-seeking than men (Niksic *et al*, 2015). Women may perceive numerous reasons to deter them from seeking medical help, imposed by socio-cultural norms. Female breasts have a social symbolism, linked with their maternal and sexual function. Therefore, breast cancer may be

perceived as a violation of femininity, and a source of shame and stigma (Wong & King, 2008). Niksic *et al* (2015) found that women have about 20% higher odds than men to report that they would be put off consulting a doctor because of embarrassment, worry about wasting the doctor's time, and perceived difficulty in talking to the doctor. These barriers were found to have a strong negative association with breast cancer survival.

We also need to acknowledge that the regression coefficients were statistically significant for most cancer symptoms in relation to the cancer survival index, but not for site-specific indices. This may be because of several reasons. First, an attenuation effect due to lower reliability, noting that many of the bowel cancer coefficients are about two thirds the size of the overall coefficients. Second, good matching between the survival index, which included different types of cancer, and awareness score, which included different types of cancer symptoms. For example, it would be surprising if we observed that 'difficulty in swallowing' was associated with breast cancer survival. However, statistically significant association of this symptom with the cancer survival index is not surprising because survival from oesophageal cancer is included in this index. Third, survival from site-specific cancers was not associated with awareness score, as expected, given that this score included a range of different symptoms that were not necessarily relevant for individual cancer sites. Fourth, breast, lung and bowel cancer survival indices may have lower statistical power to detect effects of cancer symptom awareness and barriers on survival.

Furthermore, the coefficients for the barriers score for overall cancer survival index were close to zero, with the coefficients for breast cancer survival generally negative ($p < 0.05$), and for lung and bowel cancer generally positive (although not statistically significant). Therefore, it is possible that there was some cancelling out effect in that the barriers that were harmful for breast cancer survival may well have been protective for the other two cancers. Finally, the combination of positive and negative coefficients between overall cancer survival index and different barriers to help-seeking suggests that the use of the aggregate barriers score may not be warranted. Some barriers may be associated with prompt help-seeking, while other barriers may prolong interval to the first medical consultation. If this is true, different barriers to help-seeking need to be analysed separately, in order to understand the mechanism behind them and their relationship with cancer survival.

Further studies are required to confirm or refute these findings. More specifically, future studies should assess whether the observed associations exist at the individual level, and explore the reasons driving these results, such as educational attainment, ethnicity, number of doctors per PCT, or access to health care services. Researchers should also focus on exploring whether gender differences in cancer awareness and barriers to help-seeking could

account for some of the differences in survival from the most common types of cancer. It is necessary to understand the role of different barriers to help-seeking in cancer survival, and which mechanisms drive these differences. Understanding whether experiencing a cluster of cancer symptoms prompts or hinders help-seeking, and whether it is associated with cancer survival would be useful.

Conclusion and implications

Our study suggests that higher cancer symptom awareness is associated with better cancer survival, which may inform future health campaigns and policy decision makers. We highlighted areas for improvement, and the relative position of each of the 52 PCTs with regard to cancer awareness, barriers to help-seeking and cancer survival. Future campaigns should help people to recognise early symptoms of different types of cancer and encourage preventive health behaviour, especially in socio-economically deprived areas. Efforts should be made to alleviate barriers to seeking medical help in women with symptoms of breast cancer.

References

Banning M (2007) Advanced breast cancer: aetiology, pathogenesis and treatment. *Cancer Nursing Practice* **6**(3): 35-39

Berrino F, De Angelis R, Sant M, Rosso S, Lasota MB, Coebergh JW, Santaquilani M (2007) Survival for eight major cancers and all cancers combined for European adults diagnosed in 1995-99: results of the EURO CARE-4 study. *Lancet Oncology* **8**(9): 773-783

Bojke C, Gravelle H, Wilkin D (2001) Is bigger better for primary care groups and trusts? *BMJ* **322**: 599-602

Cancer Research UK (2015) Be Clear on Cancer. Access date 12/04/2015. Available via: <http://www.cancerresearchuk.org/health-professional/early-diagnosis-activities/be-clear-on-cancer>.

Cattell V (2001) Poor people, poor places, and poor health: the mediating role of social networks and social capital. *Social science & medicine* **52**(10): 1501-1516

Cheyne L, Taylor A, Milton R, Fear J, Callister M (2013) Social deprivation does not affect lung cancer stage at presentation or disease outcome. *Lung Cancer* **81**(2): 247-251

Coleman M, Forman D, Bryant H, Butler J, Rachet B, Maringe C, Nur U, Tracey E, Coory M, Hatcher J, McGahan C, Turner D, Marrett L, Gjerstorff M, Johannesen T, Adolfsson J, Lambe M, Lawrence G, Meechan D, Morris E, Middleton R, Steward J, Richards M, The ICBP Module Working Group (2011) Cancer survival in Australia, Canada, Denmark, Norway, Sweden, and the UK, 1995-2007 (the International Cancer Benchmarking Partnership): an analysis of population-based cancer registry data. *Lancet* **377**: 127-138

Coleman MP, Rachet B, Woods LM, Mitry E, Riga M, Cooper N, Quinn MJ, Brenner H, Esteve J (2004) Trends and socioeconomic inequalities in cancer survival in England and Wales up to 2001. *British Journal of Cancer* **90**(7): 1367-1373

Department for Communities and Local Government (2011) English indices of deprivation 2010: PCT summaries. Accessed on 03/11/2015. Available via <https://www.gov.uk/government/publications/english-indices-of-deprivation-2010> Department for Communities and Local Government

Department of Health (2000) The NHS Cancer Plan: A Plan for Investment, a Plan for Reform. London: Department of Health

Department of Health (2011) *Improving Outcomes: A Strategy for Cancer* London: Department of Health.

Department of Health (2014) *Improving Outcomes: A Strategy for Cancer*. London: Department of Health, NHS England, Public Health England

Dixon-Woods M, Cavers D, Agarwal S, Annandale E, Arthur A, Harvey J, Hsu R, Katbamna S, Olsen R, Smith L (2006) Conducting a critical interpretive synthesis of the literature on access to healthcare by vulnerable groups. *BMC Medical Research Methodology* **6**(1): 35

East London NHS Trust (2013) *Summary of the Health and Wellbeing Strategies for the City of London and the London Boroughs of Hackney, Newham and Tower Hamlets*. Accessed on 15/06/2015. Available via:

<https://www.elft.nhs.uk/uploads/files/1/About/Our%20Board/Trust%20Board%20Meetings/Trust%20Board%20Meeting%20Papers%202013/September%202013/TBD-2013-09-26%20Summary%20of%20Health%20&%20Wellbeing%20Strategies.pdf> London: East London NHS Foundation Trust

Environmental Systems Research Institute (2014) *ArcGIS for Desktop Advanced 10.2*: ESRI

Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, Rosso S, Coebergh J, Comber H, Forman D, Bray F (2013) Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. *European journal of cancer* **49**(6): 1374-1403

Forbes LJJ, Atkins L, Thurnham A, Layburn J, Haste F, Ramirez AJ (2011) Breast cancer awareness and barriers to symptomatic presentation among women from different ethnic groups in East London. *British Journal of Cancer* **105**(10): 1474-1479

Forbes LJJ, Simon AE, Warburton F, Boniface D, Brain KE, Dessaix A, Donnelly C, Haynes K, Hvidberg L, Lagerlund M, Lockwood G, Tishelman C, Vedsted P, Vigmostad MN, Ramirez AJ, Wardle J (2013) Differences in cancer awareness and beliefs between Australia, Canada, Denmark, Norway, Sweden and the UK (the International Cancer Benchmarking Partnership): do they contribute to differences in cancer survival? *British Journal of Cancer* **108**(2): 292-300

Gaston-Johansson F, Fall-Dickson JM, Bakos AB (1999) Fatigue, Pain, and Depression in Pre-Autotransplant Breast Cancer Patients. *Cancer practice* **7**(5): 240-247

Goodwin R, Holme S, Roberts D (2004) Variations in registration of skin cancer in the United Kingdom. *Clinical and experimental dermatology* **29**(3): 328-330

Hamilton W, Peters TJ, Round A, Sharp D (2005a) What are the clinical features of lung cancer before the diagnosis is made? A population based case-control study. *Thorax* **60**: 1059-1065

Hamilton W, Round A, Sharp D, Peters T (2005b) Clinical features of colorectal cancer before diagnosis: a population-based case-control study. *British Journal of Cancer* **93**: 399-405

Hiom S (2015) Diagnosing cancer earlier: reviewing the evidence for improving cancer survival. *British journal of cancer*

Hvidberg L, Wulff CN, Pedersen AF, Vedsted P (2014) Barriers to healthcare seeking, beliefs about cancer and the role of socio-economic position. A Danish population-based study. *Preventive medicine* **71**: 107-113

Ironmonger L, Ohuma E, Ormiston-Smith N, Gildea C, Thomson C, Peake M (2014) An evaluation of the impact of large-scale interventions to raise public awareness of a lung cancer symptom. *British journal of cancer* **112**: 207-216

Kim H-J, McGuire DB, Tulman L, Barsevick AM (2005) Symptom clusters: concept analysis and clinical implications for cancer nursing. *Cancer nursing* **28**(4): 270-282

Macintyre S, Ellaway A, Cummins S (2002) Place effects on health: how can we conceptualise, operationalise and measure them? *Social science & medicine* **55**(1): 125-139

Marmot M (1998) Improvement of social environment to improve health. *Lancet* **351**(9095): 57

Marmot MG, Stansfeld S, Patel C, North F, Head J, White I, Brunner E, Feeney A, Marmot M, Smith GD (1991) Health inequalities among British civil servants: the Whitehall II study. *The Lancet* **337**(8754): 1387-1393

Møller H, Flatt G, Moran A (2011) High cancer mortality rates in the elderly in the UK. *Cancer Epidemiology* **35**(5): 407-412

Møller H, Sandin F, Bray F, Klint A, Linklater KM, Purushotham A, Robinson D, Holmberg L (2010) Breast cancer survival in England, Norway and Sweden: A population-based comparison. *International Journal of Cancer* **127**(11): 2630-2638

Møller H, Sandin F, Robinson D, Bray F, Klint Å, Linklater KM, Lambert PC, Pålman L, Holmberg L, Morris E (2012) Colorectal cancer survival in socioeconomic groups in England: variation is mainly in the short term after diagnosis. *European Journal of Cancer* **48**(1): 46-53

NCIN (2013) *Non-melanoma skin cancer in England, Scotland, Northern Ireland, and Ireland*. NCIN Data Briefing: National Cancer Intelligence Network.

NHS (2012) NHS Commissioning Board: Local area teams. Accessed on 29/04/2013. Available via: <http://www.networks.nhs.uk/nhs-networks/respiratory-leads/documents/NHSCB%20LATs%20-%20Specialised%20Commissioning.pdf>

Niksic M, Rachet B, Warburton F, Forbes L (2016) Ethnic differences in cancer symptom awareness and barriers to seeking medical help in England *British Journal of Cancer* **115**: 136–144

Niksic M, Rachet B, Warburton F, Wardle J, Ramirez A, Forbes L (2015) Cancer symptom awareness and barriers to symptomatic presentation in England - Are we clear on cancer? *British Journal of Cancer* **113**(3): 533–542

Nur U, Quaresma M, De Stavola B, Peake M, Rachet B (2015) Inequalities in non-small cell lung cancer treatment and mortality *Journal of Epidemiology and Community Health* **69**(10): 985–992

Office for National Statistics (2012a) *Cancer Incidence and Mortality in the United Kingdom, 2008-10*. Access date 12/05/2015. Available via: http://webarchive.nationalarchives.gov.uk/20160105160709/http://www.ons.gov.uk/ons/dcp171778_289890.pdf. London, UK Office for National Statistics.

Office for National Statistics (2012b) *Index of Cancer Survival for Primary Care Trusts in England: Patients diagnosed 1996–2010 and followed up to 2011* <http://www.ons.gov.uk/ons/rel/cancer-unit/combined-cancer-survival-by-primary-care-trusts/patients-diagnosed-1996-2010--followed-up-to-2011/stb-index-of-cancer-survival-by-pct-s.html>: Office for National Statistics.

Office for National Statistics (2012c) *Index of Cancer Survival for Primary Care Trusts in England: Patients diagnosed 1996–2010 and followed up to 2011*. Access date: 07/04/2012. Available via: http://www.ons.gov.uk/ons/dcp171778_292580.pdf and <http://www.ons.gov.uk/ons/publications/re-reference-tables.html?edition=tcm%3A77-285389>: London: Office of National Statistics.

Pashayan N, Powles J, Brown C, Duffy S (2006) Excess cases of prostate cancer and estimated overdiagnosis associated with PSA testing in East Anglia. *British journal of cancer* **95**(3): 401-405

Perme MP, Stare J, Estève J (2012) On Estimation in Relative Survival. *Biometrics* **68**(1): 113-120

Piantadosi S, Byar DP, Green SB (1988) The ecological fallacy. *American Journal of Epidemiology* **127**(5): 893-904

Popay J, Thomas C, Williams G, Bennett S, Gatrell A, Bostock L (2003) A proper place to live: health inequalities, agency and the normative dimensions of space. *Social Science & Medicine* **57**(1): 55-69

Quaresma M, Coleman MP, Rachet B (2015) 40-year trends in an index of survival for all cancers combined and survival adjusted for age and sex for each cancer in England and Wales, 1971–2011: a population-based study. *The Lancet* **385**(9974): 1206-1218

Quaresma M, Jakomis N, Gordon E, Carrigan C, Coleman M, Rachet B (2011) *Index of Cancer Survival for Primary Care Trusts in England - Patients diagnosed 1996-2009 and followed up to 2010*.: Office for National Statistics.

Quaresma M, Whitehead S., Coleman MP., Rachet B. (2012) *Index of Cancer Survival for Primary Care Trusts in England: Patients diagnosed 1996-2010 and followed up to 2011*. Access date: 07/04/2012. Available via: http://webarchive.nationalarchives.gov.uk/20160105160709/http://www.ons.gov.uk/ons/dcp171778_292580.pdf. London, UK: Office for National Statistics 2012.

Rachet B, Ellis L, Maringe C, Chu T, Nur U, Quaresma M, Shah A, Walters S, Woods L, Forman D (2010) Socioeconomic inequalities in cancer survival in England after the NHS cancer plan. *British Journal of Cancer* **103**(4): 446-453

Richards M (2009) The National Awareness and Early Diagnosis Initiative in England: assembling the evidence. *British Journal of Cancer* **101 Suppl 2**: s1-s4

Robb KA, Stubbings S, Ramirez A, Macleod U, Austoker J, Waller J, Hiom S, Wardle J (2009) Public awareness of cancer in Britain: A population based survey in adults. *British Journal of Cancer* **101**: s18-s23

STATA Corp (2015) Stata 14.0 Statistical Software. *Texas: College Station*

Stubblings S, Robb KA, Waller J, Ramirez A, Austoker J, Macleod U, Hiom S, Wardle J (2009) Development of a measurement tool to assess public awareness of cancer. *British Journal of Cancer* **101**: s13-s17

Waller J, Robb KA, Stubblings S, Ramirez A, Macleod U, Austoker J, Hiom S, Wardle J (2009) Awareness of cancer symptoms and anticipated help-seeking among ethnic minority groups in England. *British Journal of Cancer* **101**: s24-s30

Walters S, Quaresma M, Coleman MP, Gordon E, Forman D, Rachet B (2011) Geographical variation in cancer survival in England, 1991-2006: an analysis by Cancer Network. *Journal of Epidemiology and Community Health* **65**: 1044-1052

Wong N, King T (2008) The cultural construction of risk understandings through illness narratives. *Journal of Consumer Research* **34**(5): 579-594

Table 1: Primary Care Trusts included in the analysis, with cancer awareness and barriers scores, cancer survival index (all cancers combined, year 2010), weights applied in regression analyses, and number of survey participants and cancer patients per PCT

Region	Primary Care Trust	ID	Awareness score*	Barriers score*	Number of participants	Cancer Survival Index (%)	Variance of cancer survival index	Weight**	Number of patients
Anglia	Bedfordshire	1	8.1	1.4	671	65.8	0.10	6.7	2848
	Cambridgeshire	2	8.0	1.7	679	70.9	0.07	10.0	4603
	Great Yarmouth & Waveney	3	8.1	1.3	518	68.1	0.10	7.2	2143
	Norfolk	4	7.5	1.8	498	68.5	0.03	22.2	7064
	Peterborough	5	8.3	1.5	537	66.1	0.19	3.4	1148
	Suffolk	6	8.0	1.8	624	69.1	0.07	11.0	4960
Central	Hampshire	7	7.8	1.5	387	69.9	0.04	16.4	9190
South Coast	Isle of Wight	8	7.8	1.6	378	66.5	0.17	4.1	1149
	Portsmouth City	9	7.8	1.5	342	65.5	0.16	4.1	1210
	Southampton City	10	7.3	1.6	343	65.5	0.22	2.6	1314
East Midlands	Northamptonshire	11	6.9	0.8	1939	66.5	0.07	9.0	4182
Humber & Yorkshire Coast	East Riding	12	7.7	1.4	1212	69.2	0.09	8.2	2751
	Hartlepool	13	7.3	1.5	94	65.4	0.29	2.4	721
	Hull	14	7.4	1.4	936	65.2	0.12	5.4	1621
	North East Lincolnshire	15	7.7	1.6	1007	67.6	0.17	4.2	1166
	North Lincolnshire	16	7.4	1.6	1081	67.9	0.17	4.4	1172
Kent and Medway	Eastern & Coastal Kent	17	7.4	1.6	1058	65.4	0.07	10.1	5469
	Medway	18	7.2	1.9	1015	61.1	0.14	5.1	1617
	West Kent	19	7.5	1.8	775	66.5	0.07	9.0	4378
Mount Vernon	Hertfordshire	20	7.3	1.2	777	67.7	0.05	12.7	7241
	Luton	21	6.8	1.3	378	63.8	0.19	3.7	1154
North East London	Barking & Dagenham	22	6.9	1.8	245	60.5	0.32	1.7	879
	City & Hackney	23	6.7	2.2	245	65.1	0.25	2.8	933
	Havering	24	7.1	2.1	360	65.5	0.13	4.9	1685
	Newham	25	5.5	2.7	193	61.7	0.21	3.0	954
	Redbridge	26	6.4	1.8	318	64.3	0.16	3.9	1295
	Tower Hamlets	27	4.8	3.1	258	65.1	0.37	1.7	870
	Waltham Forest	28	6.6	1.5	274	63.8	0.24	2.6	1159
North Trent	Barnsley	29	7.7	1.7	795	66.0	0.18	3.7	1727
	Bassetlaw	30	7.7	1.7	811	65.1	0.36	1.9	809
	Doncaster	31	7.4	1.7	1217	66.3	0.12	4.9	2233
	Sheffield	32	7.6	1.7	2993	69.4	0.09	6.9	3583
North of England	Lincolnshire	33	7.9	1.5	95	69.7	0.07	10.0	5762
	Middlesbrough	34	7.0	1.3	298	65.8	0.23	3.1	1009
	Newcastle	35	7.3	1.0	713	67.3	0.11	5.9	1875
	North Tyneside	36	7.5	0.6	242	66.9	0.16	4.1	1538
	Northumberland	37	7.8	1.1	283	69.0	0.11	6.0	2412
South East London	Bexley	38	7.2	2.7	155	67.4	0.13	5.3	1495
	Lambeth	39	6.5	1.2	2144	66.3	0.28	2.3	1159
South West London	Croydon	40	7.3	1.8	824	67.8	0.19	3.2	1834
	Kingston	41	7.1	1.8	830	68.8	0.26	2.5	808
	Richmond & Twickenham	42	7.5	1.9	839	71.7	0.30	2.2	972
	Sutton & Merton	43	7.2	2.0	1644	69.1	0.09	7.2	2146
	Wandsworth	44	6.7	2.4	841	68.7	0.17	4.2	1304
Three Counties	Gloucestershire	45	6.5	2.2	307	68.1	0.07	9.1	4283
	Herefordshire	46	7.1	1.1	290	67.5	0.16	4.3	1333
	Worcestershire	47	6.6	1.3	386	68.1	0.09	7.2	3982
Yorkshire	Bradford & Airedale	48	6.8	3.0	336	68.4	0.08	8.8	2860
	Kirklees	49	7.0	3.1	346	66.8	0.09	7.8	2524
	Leeds	50	7.3	2.4	272	69.1	0.06	10.2	4947
	North Yorkshire & York	51	7.5	1.6	1279	69.9	0.04	18.6	5590
	Wakefield District	52	7.6	2.6	226	67.1	0.10	6.7	2352
Total					35308				133413

Region: a region in England, defined using cancer networks. ID: A sequential PCT identifier.

* Standardised for age and sex. ** Weight = (1/variance(survival)), inverse variance of the cancer survival index, for all cancers combined, in each PCT.

Number of patients: these were patients diagnosed during 2010 in each PCT with breast, lung, bowel or any other type of cancer (except prostate and non-melanoma skin cancer). Number of participants: these were participants who completed the CAM survey.

Figure 1: Cancer symptom awareness score by PCT in England





Figure 2: Barriers to help-seeking score by PCT in England

- 0.00 - 0.00 No data
- 0.59 - 1.43 Low barriers (n=13 PCTs in the lowest quartile)
- 1.44 - 1.87 Intermediate barriers (n=26 PCTs in the middle two quartiles)
- 1.88 - 3.01 High barriers (n=13 PCTs in the highest quartile)




Figure 3: The position of 52 PCTs with regard to (direction: from left to right) the cancer survival index, breast, lung and bowel cancer survival (y-axis) and the cancer awareness score, and the cancer-specific symptoms: an unexplained lump/swelling, persistent cough/hoarseness, and a change in bowel/bladder habits in each PCT (x-axis), standardised by age and sex, and weighted for the inverse variance of appropriate survival estimates.




Figure 4: The position of 52 PCTs with regard to (direction: from left to right) the cancer survival index, breast, lung and bowel cancer survival (y-axis) and barriers score (x-axis), standardised by age and sex, and weighted for the inverse variance of appropriate survival estimates.




Table 2: Linear regression coefficients estimating the relationship between cancer survival indices and cancer awareness (score and individual symptoms) and barriers to help-seeking (score and individual barriers) at PCT level

Independent variables	Overall cancer survival index		Breast cancer survival		Lung cancer survival		Bowel cancer survival	
Awareness score	1.56	(0.56; 2.57)	0.17	(-0.38; 0.71)	0.58	(-1.34; 2.51)	0.86	(-0.58; 2.30)
Unexplained lump/swelling	0.12	(0.01; 0.24)	0.01	(-0.06; 0.08)	0.02	(-0.19; 0.23)	0.05	(-0.12; 0.22)
Unexplained persistent pain	0.06	(-0.04; 0.15)	0.02	(-0.03; 0.07)	0.02	(-0.15; 0.20)	0.01	(-0.12; 0.14)
Unexplained bleeding	0.11	(0.01; 0.21)	0.00	(-0.05; 0.05)	0.04	(-0.14; 0.23)	0.05	(-0.09; 0.19)
Persistent cough/hoarseness	0.09	(0.01; 0.16)	0.00	(-0.04; 0.04)	0.04	(-0.09; 0.18)	0.07	(-0.04; 0.17)
Change in bowel/bladder habits	0.12	(0.04; 0.16)	0.01	(-0.03; 0.05)	0.03	(-0.11; 0.17)	0.09	(-0.02; 0.19)
Difficulty in swallowing	0.08	(0.02; 0.14)	0.00	(-0.03; 0.03)	0.05	(-0.06; 0.17)	0.06	(-0.02; 0.15)
Change in the appearance of a mole	0.14	(0.07; 0.20)	0.02	(-0.02; 0.05)	0.06	(-0.07; 0.20)	0.09	(-0.01; 0.19)
Sore that does not heal	0.05	(-0.00; 0.11)	0.01	(-0.02; 0.04)	0.01	(-0.11; 0.10)	-0.01	(-0.08; 0.07)
Unexplained weight loss	0.12	(0.04; 0.20)	0.02	(-0.02; 0.06)	0.05	(-0.10; 0.21)	0.10	(-0.01; 0.23)
Barriers score	-0.14	(-1.28; 0.99)	-0.66	(-1.20; -0.11)	0.94	(-1.03; 2.91)	0.05	(-1.56; 1.67)
Too embarrassed	-0.06	(-0.14; 0.02)	-0.05	(-0.09; -0.01)	-0.01	(-0.17; 0.14)	-0.07	(-0.19; 0.04)
Too scared	0.00	(-0.08; 0.09)	-0.02	(-0.06; 0.03)	0.00	(-0.15; 0.15)	0.03	(-0.14; 0.09)
Not confident to talk	-0.06	(-0.17; 0.05)	-0.05	(-0.11; 0.00)	0.04	(-0.16; 0.25)	-0.12	(-0.28; 0.04)
Worry about what GP may find	0.03	(-0.04; 0.09)	-0.02	(-0.05; 0.02)	0.06	(-0.06; 0.18)	0.02	(-0.08; 0.11)
Worry about other things	-0.03	(-0.12; 0.05)	-0.07	(-0.11; -0.02)	0.07	(-0.08; 0.23)	0.02	(-0.11; 0.15)
Transport difficulties	-0.09	(-0.24; 0.06)	-0.11	(-0.18; -0.03)	0.03	(-0.22; 0.32)	-0.01	(-0.23; 0.21)
Too busy	0.03	(-0.13; 0.06)	-0.07	(-0.11; -0.02)	0.07	(-0.10; 0.23)	0.04	(-0.09; 0.18)
Worry about wasting GP's time	0.04	(-0.03; 0.10)	-0.04	(-0.07; -0.01)	0.03	(-0.08; 0.14)	0.05	(-0.04; 0.14)
Difficulty talking to GP	-0.08	(-0.18; 0.03)	-0.06	(-0.11; -0.00)	-0.04	(-0.23; 0.16)	-0.15	(-0.30; 0.01)
Difficulty making appointment	-0.01	(-0.09; 0.07)	-0.06	(-0.09; -0.02)	0.07	(-0.07; 0.21)	-0.01	(-0.12; 0.11)

Weights for all cancer survival estimates were used in models where all cancers survival index was an outcome, and cancer-specific weights were used in models where outcomes were breast, lung and bowel cancer, respectively. All awareness and barriers scores (and individual symptoms/barriers) were standardised for age and sex in each PCT.

Estimates in bold were statistically significant at p<0.05 level (95% confidence intervals).

GP-general practitioner/doctor.

