

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Roberts, CH; Shaw, HA; Ferguson, N; Holland, M; Wren, BW; Stabler, RA; (2016) Draft Genome Sequence of *Robinsoniella peoriensis* 6600698, a Confounder of *Clostridium difficile* Diagnosis. *Genome announcements*, 4 (6). ISSN 2169-8287 DOI: <https://doi.org/10.1128/genomeA.01275-16>

Downloaded from: <http://researchonline.lshtm.ac.uk/3131829/>

DOI: <https://doi.org/10.1128/genomeA.01275-16>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

<https://researchonline.lshtm.ac.uk>

Draft Genome Sequence of *Robinsoniella peoriensis* 6600698, a Confounder of *Clostridium difficile* Diagnosis

Chrissy H. Roberts,^a Helen Alexandra Shaw,^a Natalie Ferguson,^b Mark Holland,^b Brendan W. Wren,^a  Richard A. Stabler^a

Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, United Kingdom^a; Microbiology, Maidstone Hospital, Maidstone, Kent, United Kingdom^b

***Robinsoniella peoriensis* is a Gram-positive, strictly anaerobic, spore-forming, rod-shaped organism. Here, we report the draft genome of *R. peoriensis* 6600698, initially classified as *Clostridium difficile* due to growth on selective agar, a fecal *gdh* PCR-positive result, and clinical symptoms. *R. peoriensis* is a potential confounder of *C. difficile* diagnosis.**

Received 21 September 2016 Accepted 27 September 2016 Published 17 November 2016

Citation Roberts CH, Shaw HA, Ferguson N, Holland M, Wren BW, Stabler RA. 2016. Draft genome sequence of *Robinsoniella peoriensis* 6600698, a confounder of *Clostridium difficile* diagnosis. *Genome Announc* 4(6):e01275-16. doi:10.1128/genomeA.01275-16.

Copyright © 2016 Roberts et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Richard A. Stabler, richard.stabler@lshtm.ac.uk.

Robinsoniella peoriensis was first identified from swine manure and manure storage pits (1) as a Gram-positive, strictly anaerobic, spore-forming, rod-shaped organism. Contemporaneously, an isolate from a deep wound on the heel of a 79-year-old person was submitted to the University of Gothenburg Culture Collection (1). Subsequently, there have been reports of clinical disease linked to *R. peoriensis* (2–4). *R. peoriensis* has been isolated from a presumptive *Clostridium difficile* diarrhea sample that tested negative for *C. difficile* toxins (5), and commensal *R. peoriensis* has been isolated from healthy, premature human neonate twins (6).

R. peoriensis 6600698 was originally isolated from an 82-year-old male on ChromID (bioMérieux) *C. difficile* selective agar plates, incubated anaerobically. The patient had elevated levels of fecal lactoferrin (IBD Scan, Techlab, Blacksburg, VA, USA), indicating intestinal inflammation. The bacteria were initially classified as nontoxicogenic *C. difficile* due to clinical symptoms, a positive *C. difficile* glutamate dehydrogenase (GDH) (*C. DIFF* CHEK, TechLab), and negative toxigenic culture (Premier Toxins A&B, Meridian Bioscience, Cincinnati, OH, USA). *C. difficile* PCR ribotyping (7) produced a “sporadic” or uncommon/new PCR ribotype that did not match existing standards. Attempts to identify the multilocus sequence type (MLST) profile (8) failed to amplify a single allele; therefore, whole-genome sequencing was performed.

R. peoriensis 6600698 was sequenced using an Illumina MiSeq (2 × 250 bp) and an Oxford Nanopore Technologies MinION MKI nanosequencer (Oxford Nanopore Technologies, United Kingdom) (ENA accession no. PRJEB15237). Sequence data were analyzed using Metrichor Agent version 2.40.17. The Illumina MiSeq generated 3,500,296 reads and 602 Mb from two runs. The MinION generated 6,245 reads and 22 Mb (maximum read length, 34 kb; median, 3,083 bp; N_{50} , 5,171 bp) from a single run. MiSeq reads were polished using Trimmomatic version 0.33 (9). MinION reads were extracted into FASTA format using poretools version 0.5.1 (10). A draft genome was assembled using SPAdes version 3.6.2 (11). The assembled contigs were further polished using GapFiller version 1.10 (12) and Pilon version 1.16 (13).

Contigs were annotated using Prokka version 1.11 (14). The draft genome consisted of 160 contigs, totaling 7,202,111 bp with 41% G+C; 61 tRNAs, 5,745 coding sequences, and 2 clustered regularly interspaced short palindromic repeats were also present. A BLAST analysis of the assembled 16S ribosomal sequence showed a 1,522/1,524 bp (>99%) identity with *R. peoriensis* PPC44 (AF445283), and the top seven matches were all *R. peoriensis* (29 February 2016).

R. peoriensis 6600698 contains a putative glutamate dehydrogenase (*gdh*) (gene 6600698_02412) that shares low-level homology to the *C. difficile* R20291 *gdh* (137/408 identities; 204/408 positives; bit score, 190; E-value, 1×10^{-55}), which may explain the false *gdh* PCR-positive result. No *C. difficile* pathogenicity locus toxin genes or gene remnants were present in the pathogenicity locus. *R. peoriensis* and *C. difficile* are both members of the order *Clostridiales* and common features of both can result in this emerging human pathogen being misdiagnosed as nontoxicogenic *C. difficile*.

Accession number(s). This whole-genome shotgun project has been deposited at EMBL under the accession numbers [FMJR01000001](https://www.ebi.ac.uk/ena/record/FMJR01000001) to [FMJR01000160](https://www.ebi.ac.uk/ena/record/FMJR01000160). The version described in this paper is the first version.

FUNDING INFORMATION

This work, including the efforts of Brendan W. Wren, was funded by Medical Research Council (MRC).

REFERENCES

- Cotta MA, Whitehead TR, Falsen E, Moore E, Lawson PA. 2009. *Robinsoniella peoriensis* gen. nov., sp. nov., isolated from a swine-manure storage pit and a human clinical source. *Int J Syst Evol Microbiol* 59: 150–155. <http://dx.doi.org/10.1099/ijs.0.65676-0>.
- Shen D, Chen R, Ye L, Luo Y, Tang YW. 2010. *Robinsoniella peoriensis* bacteremia in a patient with pancreatic cancer. *J Clin Microbiol* 48: 3448–3450. <http://dx.doi.org/10.1128/JCM.00477-10>.
- Cassir N, Laget L, Renvoisé A, Gennari JM, Drancourt M. 2012. *Robinsoniella peoriensis* infection following surgery for scoliosis: a case report. *J Med Case Rep* 6: <http://dx.doi.org/10.1186/1752-1947-6-174>.
- Jeon Y, Kim TS, Kim HB, Park KU, Song J, Kim EC. 2012. First Korean

- case of *Robinsoniella peoriensis* bacteremia in a patient with aspiration pneumonia. *Ann Lab Med* 32:370–374. <http://dx.doi.org/10.3343/alm.2012.32.5.370>.
5. Gomez E, Gustafson DR, Colgrove R, Ly T, Santana R, Rosenblatt JE, Patel R. 2011. Isolation of *Robinsoniella peoriensis* from four human specimens. *J Clin Microbiol* 49:458–460. <http://dx.doi.org/10.1128/JCM.01305-10>.
 6. Ferraris L, Aires J, Butel MJ. 2012. Isolation of *Robinsoniella peoriensis* from the feces of premature neonates. *Anaerobe* 18:172–173. <http://dx.doi.org/10.1016/j.anaerobe.2011.11.007>.
 7. Stubbs SL, Brazier JS, O'Neill GL, Duerden BI. 1999. PCR targeted to the 16S-23S rRNA gene intergenic spacer region of *Clostridium difficile* and construction of a library consisting of 116 different PCR ribotypes. *J Clin Microbiol* 37:461–463.
 8. Griffiths D, Fawley W, Kachrimanidou M, Bowden R, Crook DW, Fung R, Golubchik T, Harding RM, Jeffery KJ, Jolley KA, Kirton R, Peto TE, Rees G, Stoesser N, Vaughan A, Walker AS, Young BC, Wilcox M, Dingle KE. 2010. Multilocus sequence typing of *Clostridium difficile*. *J Clin Microbiol* 48:770–778. <http://dx.doi.org/10.1128/JCM.01796-09>.
 9. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <http://dx.doi.org/10.1093/bioinformatics/btu170>.
 10. Loman NJ, Quinlan AR. 2014. Poretools: a toolkit for analyzing nanopore sequence data. *Bioinformatics* 30:3399–3401. <http://dx.doi.org/10.1093/bioinformatics/btu555>.
 11. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
 12. Nadalin F, Vezzi F, Policriti A. 2012. GapFiller: a de novo assembly approach to fill the gap within paired reads. *BMC Bioinformatics* 13(suppl 14):S8. <http://dx.doi.org/10.1186/1471-2105-13-S14-S8>.
 13. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <http://dx.doi.org/10.1371/journal.pone.0112963>.
 14. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.