

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Kositz, C; Butcher, R; Marks, M; (2016) New Diagnostics for Yaws. The American journal of tropical medicine and hygiene, 96 (1). pp. 3-4. ISSN 0002-9637 DOI: <https://doi.org/10.4269/ajtmh.16-0639>

Downloaded from: <http://researchonline.lshtm.ac.uk/3336110/>

DOI: <https://doi.org/10.4269/ajtmh.16-0639>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

Images in Clinical Tropical Medicine New Diagnostics for Yaws

Christian Kositz,¹ Robert Butcher,¹ and Michael Marks^{1,2*}

¹Clinical Research Department, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, United Kingdom; ²The Hospital for Tropical Diseases, London, United Kingdom

Yaws, caused by *Treponema pallidum* subsp. *pertenue*, is an important public health problem in many tropical countries.¹ Like syphilis, the disease manifests in three stages; however, unlike syphilis, its route of transmission is non-genital skin-to-skin contact and not by sexual intercourse. Primary yaws manifests as either a papilloma or a chronic ulcer. Typically, ulcers are painless, with a raised edge and friable base (Figure 1). In secondary yaws, skin manifestations, involvement of the bones and joints including periostitis have been reported. Tertiary yaws develops in a minority of patients causing destructive lesions of the skin and soft tissues. Interest in yaws has been revived by the finding that azithromycin is a highly effective treatment of both primary and secondary yaws.² Clinical diagnosis alone of primary yaws is unreliable, but a point-of-care test has been shown to be of value.³ This test provides a result analogous to a *T. pallidum* particle agglutination assay (Figure 2, line 1) and a rapid plasma reagin (RPR) assay (Figure 2, line 2). In early infection, only the RPR may be positive. Diagnosis has been further complicated by the discovery that *Haemophilus ducreyi* may cause clinically similar ulcers.⁴ New polymerase chain reaction (PCR) assays have been developed for yaws.⁵ DNA suitable for can be extracted directly from swabs collected into dry tubes without the need for transport medium. Figure 3 demonstrates real-time PCR amplification curves of positive and negative controls and a clinical swab from a yaws lesion containing *T. pallidum pertenuae* DNA. Both serological and molecular tests have a major role to play in the World Health Organization yaws eradication campaign.



FIGURE 1. Classical primary ulcer of yaws.

Received August 3, 2016. Accepted for publication September 6, 2016.

Financial support: Michael Marks is supported by a Wellcome Trust Clinical Research Fellowship (102807) and Robert Butcher is supported by the Fred Hollows Foundation (1041).

Authors' addresses: Christian Kositz and Robert Butcher, Clinical Research Department, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, United Kingdom, E-mails: christian.kositz@gmail.com and robert.butcher@lshtm.ac.uk. Michael Marks, Clinical Research Department, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, United Kingdom, and The Hospital for Tropical Diseases, London, United Kingdom, E-mail: michael.marks@lshtm.ac.uk.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




FIGURE 2. Combined treponemal and nontreponemal rapid diagnostic test for yaws.

*Address correspondence to Michael Marks, Clinical Research Department, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, United Kingdom. E-mail: michael.marks@lshtm.ac.uk




FIGURE 3. Quantitative polymerase chain reaction for yaws.

REFERENCES

1. Marks M, Solomon AW, Mabey DC, 2014. Endemic treponemal diseases. *Trans R Soc Trop Med Hyg* 108: 601–607.
2. Mitjà O, Hays R, Ipaï A, Penias M, Paru R, Fagaho D, de Lazzari E, Bassat Q, 2012. Single-dose azithromycin versus benzathine benzylpenicillin for treatment of yaws in children in Papua New Guinea: an open-label, non-inferiority, randomised trial. *Lancet* 379: 342–347.
3. Marks M, Goncalves A, Vahi V, Sokana O, Puiahi E, Zhang Z, Dalipanda T, Bottomley C, Mabey D, Solomon AW, 2014. Evaluation of a rapid diagnostic test for yaws infection in a community surveillance setting. *PLoS Negl Trop Dis* 8: e3156.
4. Marks M, Chi K-H, Vahi V, Pillay A, Sokana O, Pavluck A, Mabey DC, Chen CY, Solomon AW, 2014. *Haemophilus ducreyi* associated with skin ulcers among children, Solomon Islands. *Emerg Infect Dis* 20: 1705–1707.
5. Mitjà O, Lukehart SA, Pokowas G, Moses P, Kapa A, Godornes C, Robson J, Cherian S, Houinei W, Kazadi W, Siba P, de Lazzari E, Bassat Q, 2014. *Haemophilus ducreyi* as a cause of skin ulcers in children from a yaws-endemic area of Papua New Guinea: a prospective cohort study. *Lancet Glob Health* 2: e235–e241.