

**Table 5: Association of depression and suicidality with negative outcomes:**

	<b>Depression</b>		<b>Suicidality</b>	
<b>Outcome</b>	<b><sup>1</sup>aOR (95% CI)</b>	<b>LR P-value</b>	<b><sup>1</sup>aOR (95% CI)</b>	<b>LR P-value</b>
Number of days disrupted in the last month	2.19 (1.45 ; 3.30)	0.0003	1.69 (0.77 ; 3.68)	0.19
Number of visits to a Health Unit in the last month	1.50 (0.97 ; 2.32)	0.071	1.55 (0.68 ; 3.55)	0.31
Number of days admitted to hospital in the last month	3.32 (1.32 ; 8.36)	0.016	1.83 (0.44 ; 7.57)	0.42
Number of risky sexual behaviours in the last month	1.60 (0.89 ; 2.90)	0.13	0.94 (0.29 ; 3.11)	0.92
Any risky sexual behaviour in the last month	1.48 (0.82 ; 2.68)	0.20	1.01 (0.31 ; 3.34)	0.98
Risky sexual behaviour without a condom in the last month	2.44 (0.76 ; 7.78)	0.13	0.19 (0.015 ; 2.43)	0.17
Missed Septrin / Dapsone prophylaxis in the last 3 days	1.63 (1.02 ; 2.61)	0.046	0.45 (0.14 ; 1.43)	0.15

**Note** <sup>1</sup>Adjusted odds ratio: Adjusted for study site, age, sex, educational level and most recent CD4 count (in the past 6 months)