

Table 2: Suicidality profiles of community study and facility detection study participants in PRIME implementation areas in Ethiopia, India, Nepal, South Africa and Uganda, 2013-2014.

	Ethiopia		India		Nepal		South Africa	Uganda
	Community Residents	Facility attendees	Community Residents	Facility attendees	Community Residents	Facility attendees	Facility attendees	Facility attendees
Ideation % (95% CI)	7.0 (5.2-9.4)	10.5 (8.6-12.6)	11.1 (9.6-13.0)	9.3 (7.1-11.9)	3.5 (2.6-4.8)	11.2 (9.3-13.6)	14.8 (9.8-21.7)	5.0 (3.8-6.5)
Planning % (95% CI)	4.6 (3.2-6.4)	7.9 (6.3-9.9)	1.1 (0.7-1.8)	2.0 (1.2-3.2)	0.9 (0.4-2.2)	2.8 (1.8-4.5)	9.5 (5.6-15.8)	1.2 (0.8-1.8)
Planning % (among those with ideation)	65.6	76.6	11.7	32.1	24.3	23.7	58.7	18.3
Attempt % (95% CI)	3.7 (2.4-5.5)	5.6 (4.2-7.4)	0.4 (0.3-0.7)	0.5 (0.1-1.4)	0.7 (0.2-2.1)	1.2 (0.6-2.4)	7.4 (3.8-14.0)	0.3 (0.1-0.6)
Attempt % (among those planning)	72.2	65.2	39.9	24.9	70.4	57.1	55.1	34.7
Disclosure of attempt % (among those with attempt)	45.3	29.2	35.6	35.1	19.6	18.3	51.1	58.3
Help seeking after attempt % (among those with attempt)	26.0	15.7	0.0	0.0	0.0	0.0	21.5	0.0

Note: All percent figures are standardized to the age structure for the INDEPTH 2013 male and female populations of low- and middle-income countries; Percent figures from the Community Survey are also adjusted for the complex sampling design within each site.