

**Table 4.** Utility coefficients from paired conditional logistic regression analysis.

Attribute-level	Coeff.	SE	p-value	95% CI	
<i>Frequency (and setting) of physical (and larynx/pharynx endoscopic) investigations</i>					
Every 2-3 months for 3 years (primary care-based follow-up for 2 more years)	0.041	0.194	0.834	-0.340	0.422
Every 2-3 months for 2 years, every 5-6 months for 3 more years	2.523	0.115	<0.001	2.297	2.749
Every 2-3 months for 5 years	2.155	0.141	<0.001	1.877	2.432
<i>Frequency of MRI/CT scans</i>					
Only at the occurrence of new symptoms	0.526	0.112	<0.001	0.3055	0.746
One examination only at the beginning of follow-up (later only at occurrence of new symptoms)	0.565	0.122	<0.001	0.326	0.804
Once or twice a year	1.885	0.099	<0.001	1.691	2.079
<i>Frequency (and eligibility) of PET scans</i>					
No PET scan during follow-up	-	-	-	-	-
Yearly PET scan only for high-risk patients ( $\geq 50$ years and heavy smokers)	0.962	0.109	<0.001	0.748	1.176
Yearly PET scan for all patients	1.111	0.138	<0.001	0.840	1.382
<i>Telephone calls to monitor occurrence of new symptoms</i>					
No inter-visit calls from the hospital	0.092	0.093	0.322	-0.091	0.276
Inter-visit calls by the nurse	0.087	0.112	0.439	-0.133	0.307
Inter-visit calls by the oncologist	0.601	0.115	<0.001	0.375	0.826

CI: confidence interval; CT: computed tomography; MRI: magnetic resonance imaging; PET: positron emission tomography; SE: standard error.