

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Cowling, TE; Majeed, A; Harris, MJ; (2018) Patient experience of general practice and use of emergency hospital services in England: regression analysis of national cross-sectional time series data. *BMJ quality & safety*. ISSN 2044-5415 DOI: <https://doi.org/10.1136/bmjqs-2017-007174>

Downloaded from: <http://researchonline.lshtm.ac.uk/4646255/>

DOI: <https://doi.org/10.1136/bmjqs-2017-007174>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

Patient experience of general practice and use of emergency hospital services in England: regression analysis of national cross-sectional time series data

Thomas E Cowling^{1 2}

Azeem Majeed¹

Matthew J Harris¹

¹Department of Primary Care and Public Health, Imperial College London

²Department of Health Services Research and Policy, London School of Hygiene and Tropical Medicine

Correspondence to: TE Cowling t.cowling@imperial.ac.uk

Abstract

Background The UK Government has introduced several national policies to improve access to primary care. We examined associations between patient experience of general practice and rates of visits to accident and emergency (A&E) departments and emergency hospital admissions in England.

Methods The study included 8,124 general practices between 2011-12 and 2013-14. Outcome measures were annual rates of A&E visits and emergency admissions by general practice population, according to administrative hospital records. Explanatory variables included three patient experience measures from the General Practice Patient Survey: practice-level means of experience of making an appointment, satisfaction with opening hours, and overall experience (on 0-100 scales). The main analysis used random-effects Poisson regression for cross-sectional time series. Five sensitivity analyses examined changes in model specification.

Results Mean practice-level rates of A&E visits and emergency admissions increased from 2011-12 to 2013-14 (310.3 to 324.4 and 98.8 to 102.9 per 1,000 patients). Each patient experience measure decreased; for example, mean satisfaction with opening hours was 79.4 in 2011-12 and 76.6 in 2013-14. In the adjusted regression analysis, a standard deviation increase in experience of making appointments (equal to nine points) predicted decreases of 1.8% (95% CI: -2.4% to -1.2%) in A&E visit rates and 1.4% (95% CI: -1.9% to -0.9%) in admission rates. This equalled 301,174 fewer A&E visits and 74,610 fewer admissions nationally per year. Satisfaction with opening hours and overall experience were not consistently associated with either outcome measure across the main and sensitivity analyses.

Conclusions Associations between patient experience of general practice and use of emergency hospital services were small or inconsistent. In England, realistic short-term improvements in patient experience of general practice may only have modest effects on A&E visits and emergency admissions.

Introduction

The responsiveness of health systems to the expectations of their populations is a key measure of health system performance internationally. In 2000, the World Health Organization listed this responsiveness as one of three overall goals for health systems to achieve (alongside good health and fair financing).¹ Having high quality and accessible primary care services is central to this goal.² As a result, many countries have experimented with primary care reform to improve patients' experiences of healthcare. For example, reform in the United States has focused on the Primary Care Medical Home model, which includes accessibility as a central principle; operational standards require primary care practices to offer appointments in the evenings and at weekends, for example.³ Part of the interest in this model also comes from the hypothesis that more responsive primary care services will reduce use of emergency hospital services.⁴⁻⁹ This hypothesis has also been studied in European countries, such as in Italy where extended opening hours up to 12 hours each day were found to reduce certain types of emergency department visits in one region.¹⁰

In England, the National Health Service (NHS) has planned several changes to general practice services.¹¹ One widely debated policy is for general practices to offer more appointments in the evenings and at weekends.¹²⁻¹³ The UK Government's mandate to the NHS states '100% of population has access to weekend/evening routine GP appointments' as a goal to achieve by 2020.¹⁴ Other interventions promoted to improve patient experience include greater use of telephone and video consultations and online appointment booking systems.¹⁵ One goal of these reforms, as in other countries, is to reduce use of emergency hospital services.¹⁵ The number of visits to accident and emergency (A&E) departments in England increased from 18.9 to 22.9 million from 2006-07 to 2015-16.¹⁶ The number of emergency hospital admissions increased from 4.7 to 5.8 million over the same period.¹⁷¹⁸ However, the relationships between patient experience of general practice and rates of A&E visits and emergency admissions, and their policy implications, remain unclear.

Several observational studies¹⁹⁻³⁴ have investigated associations between use of emergency hospital services and patient experience of access to general practice by linking administrative records from English hospitals to a national patient survey—the General Practice Patient Survey (GPPS). However, these studies have focused on individual English regions,¹⁹⁻²¹ demographic groups (e.g. children),²² or conditions (e.g. asthma)²³⁻³¹; only analysed A&E visits or a subset of them³²⁻³⁴; and have often not translated abstract statistical results into absolute numbers that are actionable. Consequently, past research does not suggest what the effect of improving patient experience of general practice on total use of emergency hospital services, across the whole population, is likely to be. This is arguably what is most relevant to national policy makers. It is not possible to synthesise results across the past studies because of substantial differences in their methods. These studies often have further limitations such as analysing only one data year cross-sectionally;^{19-23 31-34} cross-sectional analyses cannot examine temporal relationships between variables and can be more susceptible to residual confounding.

In this study, we use national data to assess relationships between all A&E visits and emergency admissions in England with three measures of patient experience from the GPPS. Two of these measures, relating to experiences

of making appointments and overall experiences, are monitored in the national outcomes framework for the NHS.³⁵ The third measure—satisfaction with opening hours—is directly relevant to plans to extend the opening hours of general practices. Each measure is plausibly linked to use of emergency hospital services. For example, patients with poor experiences overall or of making appointments in the past may be more likely to visit an A&E department instead. Many patients report visiting A&E departments after being unable to get suitable general practice appointments,³⁶ which may be affected by opening hours.³⁷ Moreover, patients with acute illnesses who do not have prompt access to general practice may deteriorate and be at greater risks of requiring unplanned hospital treatment. We estimate annual differences in national numbers and costs of A&E visits and emergency admissions with changes in patient experience of general practice.

In England, almost all residents are registered with a general practice that is free at the point of use as part of the universal coverage provided by the healthcare system.

Methods

Our study was a regression analysis of cross-sectional time series data from the 2011-12, 2012-13, and 2013-14 financial years. The GPPS included 8,289 English general practices (24,392 practice-years) during this period. We excluded practice-years where practices had 1,000 or fewer registered patients or had 50 or fewer patients respond to the GPPS. This removed practices serving the smallest populations (which are likely to be atypical) with unreliable survey results at the practice level. The vast majority of practices in the GPPS remained in the study population (8,124 practices, 98% of survey sample); 23,875 practice-years was the final sample size. Most practices were present in the final sample in each year of the study period (7,802 practices, 96% of final sample).

A&E visits and emergency admissions

We used routine administrative records from Hospital Episode Statistics (HES) data to count numbers of A&E visits and emergency admissions by practice-year. HES is a data warehouse containing records of admissions, outpatient visits, and A&E visits to all NHS-funded hospitals in England. This study used the HES Inpatient and A&E datasets for the 2011-12, 2012-13, and 2013-14 financial years, as provided by the Health and Social Care Information Centre (now NHS Digital). Both datasets contained a variable recording the general practice that each patient was registered with. This allowed data to be aggregated by practice-year and linked to the GPPS.

Each record in HES A&E data corresponds to a single visit. Our main outcome variable for A&E visits was a simple count of these records by practice-year. This variable included visits to each of the three types of A&E department in England: consultant-led general services (known as emergency departments); consultant-led single specialty services (such as ophthalmology); and doctor or nurse-led minor injury and illness services (such as walk-in centres). Record coverage for emergency departments was around 100% (99% in 2012-13), but it was less for single specialty (77%) and minor injury and illness services (46%); coverage across all A&E types was

83% in 2012-13.³⁸ In this year, emergency department visits accounted for 66% of all A&E visits, but 79% of records in HES A&E data. We conducted a sensitivity analysis with only emergency departments visits included in the outcome variable. We discuss possible implications of the missing data for the results in the ‘Discussion’ section of the article.

Each record in HES Inpatient data corresponds to a continuous period of hospital care under the same consultant doctor. Some hospital stays generate several records where patients are transferred between consultants within the same hospital or between hospitals. To avoid counting several admissions in these instances, the analysis was based on ‘superspells’ which are continuous periods of care from initial admission to final discharge after any hospital transfers; we refer to superspells as admissions in the rest of the article.³⁹ The outcome variable for emergency admissions included all superspells where the method of admission indicated an emergency admission; this included emergency admissions via A&E departments (73% in 2012-13) as well as other methods such as general practitioners (GPs) and hospital outpatient clinics (27%).¹⁶ The outcome variable recorded numbers of emergency admissions by practice-year.

The outcome variables in the main analysis included all age ranges of patients, including children. A sensitivity analysis included A&E visits and emergency admissions for adults only (see ‘Sensitivity analyses’ below).

Patient experience of general practice

The GPPS is a quantitative postal survey conducted annually for the NHS. Patients aged at least 18 years old who have valid NHS numbers and have been registered with an English general practice continuously for the last six months are eligible to participate.⁴⁰ The sampling frame includes all practices with eligible patients. Questionnaires are sent to stratified (by age group, gender, and practice) random samples of eligible patients in each practice, with 2,912,535 respondents from 8,289 practices between 2011-12 and 2013-14 (36% of 8,134,705 questionnaires sent).⁴⁰⁻⁴² The mean number of responses per practice-year was 119 (standard deviation [SD]=23), which provides most patient experience measures with practice-level reliability that is ‘excellent’ (≥ 0.9) or ‘good’ (≥ 0.8) in a given year.⁴³ NHS England provided us with respondent-level survey data.

We analysed three patient experience measures—experience of making an appointment, satisfaction with opening hours, and overall experience. Each measure was generated from survey questions with five response options: satisfaction with opening hours was recorded as ‘very satisfied’ to ‘very dissatisfied’, while experience of making an appointment and overall experience were recorded as ‘very good’ to ‘very poor’ (Appendix 1). We treated responses as lying on an interval scale: 100 (most favourable), 75, 50, 25, 0 (least favourable), as is the common method in previous research.⁴⁴⁻⁴⁶ The survey asked all respondents to complete the questions analysed. We chose these measures because of their policy relevance, plausible causal mechanisms, and because they represent three distinct constructs within the GPPS; overall experience is most strongly associated with the interpersonal quality of care provided by GPs,⁴⁷ whereas experience of making an appointment is most strongly associated with the accessibility of services, and satisfaction with opening hours is only weakly associated with other GPPS measures.⁴⁸

We calculated practice-level measures of patient experience as the means of respondent-level values for each practice-year. When calculating these means, we weighted responses using the weights given in the survey datasets; these weights account for differential response probabilities (based on patient age, gender, region of England, and area-based demographic and socioeconomic indicators) and the probabilities of eligible patients being sent questionnaires in each practice.⁴⁰⁻⁴² Practice-level measures had hypothetical ranges of 0 (when all respondents chose the most negative option) to 100 (when all chose the most positive option). Some variation in patient experience measures between practices is explained by differences in respondent characteristics between practices.⁴⁴⁻⁴⁶ This variation may not reflect true differences in patient experience; it could be influenced by how different patient groups interpret questionnaires, for example. We therefore conducted a sensitivity analysis with practice-level measures adjusted for respondent case-mix (see ‘Sensitivity analyses’ below). The weighted measures represent the values of patient experience expected if all patients registered at each general practice responded, whereas the case-mix adjusted measures represent the values practices would have if they had similar patient populations.

General practice characteristics

We analysed eight characteristics of general practices as potential confounding variables for associations between patient experience and use of emergency hospital services.

The number and age-gender distribution of patients registered to each practice were provided by the Health and Social Care Information Centre (by email or online).⁴⁹ Data were for 30th September in each financial year of the study period. We coded the age-gender distributions as percentages for 14 age-gender bands⁵⁰ (male/female; 0-4 years, 5-14 years, 15-44 years, 45-64 years, 65-74 years, 75-84 years, 85 years or more), as coded in the original datasets. Ethnicity profiles for lower layer super output areas (LSOAs; small areas with mean populations of 1500) were available from the 2011 national census.⁵¹ We calculated the percentages of each practice’s population of various ethnicities (white, mixed, Asian, black, and other) as weighted sums of the percentages for each LSOA in which a practice had patients.⁵¹⁻⁵² The weights were the proportions of each practice’s population living in each LSOA.⁵² We also used this method to calculate seven measures of socioeconomic status at the practice level based on each domain of the 2010 Index of Multiple Deprivation.⁵³

We analysed several intermediate clinical outcome measures from the UK Quality and Outcomes Framework, using data provided by the Health and Social Care Information Centre for each year of the study period (Appendix 2).⁵⁴⁻⁵⁶ We focused on intermediate outcome (rather than process) measures as these show the greatest correlations with patient experience measures, though the correlations are still weak (Spearman rank correlations ≤ 0.18).⁵⁷ We used a composite measure of performance calculated as the weighted sum of achievement on intermediate outcome measures, using the relative number of points for each measure as the weights.⁵⁷ The type of general practice contract (‘General’, ‘Personal’, ‘Alternative Provider’, or ‘Primary Care Trust’ Medical Services) was also available from the Health and Social Care Information Centre in each study year.⁴⁹⁻⁵⁸ We also analysed whether a general practice was located in an urban area (population of 10,000 or more⁵⁹) and which Clinical Commissioning Group area it was in (out of 211 areas in England).

General practice variables for ethnicity, socioeconomic status, urban classification, and Clinical Commissioning Group area were treated as unchanging over the study period. Other general practice characteristics could change between financial years.

Statistical methods

We linked GPPS data with HES and other data sources at the practice level using the unique identifiers of general practices in England. Only a few practice-years in the GPPS data were not present in the other datasets (0.2%). Descriptive statistics included all eligible practice-years (23,875 for 8,124 practices). When estimating associations between variables, each model only included practice-years without missing data for any of the variables included in that model (as indicated in the results tables).

We estimated associations between the outcome and patient experience measures using Poisson regression. The outcome variable in the regression models was either the number of A&E visits or the number of emergency admissions by practice-year. Each model was estimated, in turn, for A&E visits and emergency admissions. Models included the natural logarithm of the number of patients registered to each practice as an offset term, with its coefficient equal to 1, so that the estimated coefficients represented associations with rates of the outcomes (rather than numbers). We present exponentiated coefficients, referred to as rate ratios (RRs), with $100 \times (RR - 1)$ equalling the percentage change in the rate of A&E visits or emergency admissions resulting from a specified change in a patient experience measure.

Our models included a random intercept at the general practice level to account for repeat observations on the same practices. Associations were estimated using both variation in measures between practices and within practices between financial years. This partially adjusted results for potential confounders that were unobserved and were constant within practices over the study period. Models also included either the mean A&E visit rate or emergency admission rate for each practice in the three financial years before the study period (2008-09 to 2010-11).^{50,60} These pre-sample mean rates provide extra adjustment for unobserved characteristics of practices that are constant in time and affect use of emergency hospital services. We adjusted for financial year using indicator variables, which accounted for national time trends in rates of A&E visits and emergency admissions. We calculated 95% confidence intervals from Huber-White standard errors to account for possible overdispersion and heteroscedasticity.⁶¹

In multivariable analyses, we entered patient experience measures into models as explanatory variables simultaneously, along with the measures of general practice characteristics. Each model was estimated four times: once with all three experience measures coded as continuous variables and three times with only one of these measures coded as categorical fifths. All practice characteristics given above were adjusted for (regardless of their statistical significance) as their purpose was to reduce confounding. We omitted two variables to prevent perfect collinearity among general practice characteristics: the percentages of practice populations that were male aged 15-44 years old and the percentages of white ethnicity. Remaining collinearity between the age-gender and ethnicity variables was unimportant as they were only being used as control variables (so their associations were

not of interest). We generated indicator variables for practice contract type (three indicators), urban location (one indicator), and Clinical Commissioning Group (210 indicators).

We standardised patient experience measures to have means equal to zero and SDs equal to one. This allowed associations to be interpreted in terms of a one SD change in these measures, which helped to compare the sizes of associations across measures. We used the associations to calculate absolute differences in numbers of A&E visits and emergency admissions with 0.5 and one SD increases in any patient experience measures consistently associated with the outcomes. We translated these differences into provider costs using NHS reference costs.⁶²

Sensitivity analyses

We tested the robustness of our results in five sensitivity analyses. First, we included visits to emergency departments only (rather than all A&E departments such as walk-in centres or minor injuries units) in the outcome. Second, we included visits to all types of A&E department but only included patients aged 18 years old or more, as only adults participate in the GPPS.

Third, we created practice-level measures of patient experience adjusted for the case-mix of survey respondents. To do this, we estimated random-effects linear regression models with the patient experience measures (at the respondent level) as the outcome variables and respondent characteristics (age, gender, ethnicity, socioeconomic status, ability to take time off work to see a GP, and confidence in managing own health⁶³) as explanatory variables. We then predicted the practice-level random intercepts to give empirical Bayes case-mix adjusted measures of patient experience at the practice level in each study year.⁵⁷

Fourth, we added further explanatory variables. These were: the prevalences of 21 conditions in each practice as recorded for the UK Quality and Outcomes Framework,⁵⁴⁻⁵⁶ the number of full-time equivalent GPs per 1,000 patients, the weighted percentages of GPPS respondents who knew how to contact an out-of-hours GP service, and a measure of relational continuity of care from the GPPS; this measure, detailed elsewhere,⁶⁴ assesses how often respondents who have a preferred GP consult that GP.

Fifth, we estimated associations using variation in measures within practices between years only (using practice-level fixed effects). This removed possible confounding from variables that affect use of emergency hospital services and do not change within practices over time. Data management and analysis used Stata MP V.13.⁶⁵

Results

Table 1 describes rates of A&E visits and emergency admissions and the patient experience measures at the practice level. Mean values of the patient experience measures decreased year-on-year, while use of emergency hospital services increased over the study period. Figure 1 shows the distribution of the experience of making an appointment measure at the practice level; the distributions of the other two experience measures were similar (table 1). Most survey respondents reported favourable experiences of their practices; for example, 42% and 45% of weighted respondents described their overall experiences as ‘good’ and ‘very good’ respectively (Appendix 3). The mean number of patients registered to each practice was 6,940 (SD=4,220) and most practices were located in urban areas (85.7% of practice years) (Appendix 4).

Table 2 shows that general practices with increased values for the three patient experience measures reduced rates of A&E visits on average in the unadjusted analyses. After adjusting for other variables, this association remained for experience of making an appointment only (table 2). A one SD increase in this measure predicted a 1.8% decrease in the rate of A&E visits (95% CI -2.4% to -1.2%). There was no evidence of an association for satisfaction with opening hours (RR=1.001, 95% CI 0.997 to 1.006, P=0.519) or overall experience (RR=1.004, 95% CI 0.998 to 1.010, P=0.182). The adjusted rate of A&E visits in the fifth of practices with the greatest mean values of experience of making an appointment was 2.1% lower than in the fifth with the lowest mean values (95% CI -3.4% to -0.7%) (table 2).

Table 3 shows that practices with improved experiences of making appointments also reduced mean rates of emergency admissions in both unadjusted and adjusted analyses. After adjusting for other variables, a one SD increase in this experience measure predicted a 1.4% decrease in the admission rate (95% CI -1.9% to -0.9%). The adjusted rate of emergency admissions in the fifth of practices with the greatest mean values of experience of making an appointment was 3.9% lower than in the fifth with the lowest mean values (95% CI -4.9% to -2.9%) (table 3). Table 3 also shows that emergency admission rates did not vary systematically over categories of satisfaction with opening hours or overall experience.

Table 4 presents the results of the sensitivity analyses. It shows adjusted rate ratios for A&E visits and emergency admissions comparing the fifths of practices with the greatest patient experience values to the fifths of practices with the lowest values. The results were generally consistent with the main analysis; better experiences of making appointments were associated with lower rates of A&E visits and emergency admissions in all analyses.

Table 5 provides estimates of the predicted annual differences in numbers of A&E visits and emergency admissions with 0.5 and one SD increases in experience of making an appointment. The total short-term cost saving, for both A&E visits and emergency admissions, from a one SD increase in experience of making an appointment was estimated to be £206,359,098 (\$278,025,549, €233,802,805) per year.

Discussion

General practices with improved experiences of making appointments on average reduced rates of A&E visits and emergency admissions. This association existed before and after adjusting results for several characteristics of practices and remained in all sensitivity analyses. The results suggest 301,174 fewer A&E visits and -74,610 fewer emergency admissions per year if values of this patient experience measure were one SD greater than they were during the study period. This equals a total annual cost saving to A&E departments and hospitals of approximately £206.4 million (\$278.0 million, €233.8 million). Variation between general practices in the measure of experience of making an appointment is largely explained by four other measures in the GPPS: ease of telephone contact, helpfulness of receptionists, ability to get an appointment, and appointment convenience.⁴⁸ These measures are all relevant to patients' access to services. Possible mechanisms for the above results, under a causal interpretation, include that: some patients who experience or expect problems with accessing their general practices visit an A&E department instead^{36,37}; and difficulties with access may lead to worsened health and increased risks of requiring unplanned hospital care.

However, the size of the association was modest. NHS England has committed at least £175 million (\$236 million, €198 million) to fund general practices to test new ways of improving access since April 2014.⁶⁶ The effect of these interventions on patient experience is unknown, but an increase in the experience of making an appointment measure by one SD is ambitious (nine points on the 0-100 scale); it is equal to 84% of practices having greater values of this measure than the mean value before the intervention period. An increase of 0.5 SD is more realistic in the short-term which we estimated as producing annual cost savings of £103.6 million (\$139.5 million, €117.3 million) to A&E departments and hospitals. It is therefore uncertain whether current plans to improve patient experience of general practice will provide net cost savings to the NHS. Previous research suggests that many A&E visits in England currently occur after a patient is unable to get a suitable general practice appointment,³⁶ but the extent to which this can be changed in the short-term is another matter. General practices, A&E departments, and hospitals are embedded in complex systems that may act to resist change and reinforce the status quo, while patient behaviour may change in unexpected ways.⁶⁷

Our results provide little evidence that improvements in satisfaction with opening hours or overall experience are independently associated with reduced A&E visit or emergency admission rates. Patient experience of making appointments was the only experience measure consistently associated with use of emergency hospital services, though the estimated associations were relatively modest.

Relation to existing literature

Three other studies³²⁻³⁴ have investigated associations between patient experience measures from the GPPS and rates of A&E visits across England. The first of these studies³² examined emergency department visits where patients had referred themselves and were discharged (39% of visits); practices where patients were more able to see a GP within two weekdays had lower A&E visit rates in 2010-11.³² Tammes *et al.*³³ examined data from 2012-

13 and reported similar findings for the above outcome variable and also for an outcome including additional types of A&E department. Scantlebury *et al.*³⁴ investigated all A&E visits in 2011-12 and reported that practices where patients were more likely to be able to get an appointment had lower A&E visit rates. Cecil *et al.*²² also found this association in 2011-12 for children aged less than 15 years old.²² Zhou *et al.*⁶⁸ estimated that worse experiences of in-hours general practice services were associated with greater use of out-of-hours services.

Other studies²⁴⁻³¹ have examined relationships between patient experience of accessing general practice and rates of emergency admissions in England. These studies focused on specific conditions such as asthma,³¹ chronic obstructive pulmonary disease,²⁵ and heart failure.²⁶ Practices with higher scores on GPPS measures related to access had lower adjusted admission rates in all but one study, though the exact measure(s) showing this association is inconsistent across studies. Bottle *et al.*²³ found that patients had lower risks of an emergency admission (versus elective admission) for cancer if they were more likely to be able to see a GP within two weekdays. Another study found that patients are more likely to have an emergency admission via an A&E department, rather than via their GP, if they are registered with a less accessible practice.⁶⁹

Our study develops this existing literature in several important ways. First, it included the whole population of England and was not limited to individual demographic or clinical groups. Second, it included all A&E visits and emergency admissions to assess the total, or overall, relationship between acute hospital services and patient experience of general practice. Third, it translated results into absolute reductions in A&E visits and emergency admissions and their associated costs. Fourth, it used several years of data to estimate associations with more robust methods than previous studies. This novel approach now suggests that, overall, realistic short-term improvements in patient experience of general practice will likely only have modest effects on A&E visits and emergency admissions and not provide large financial benefits to the health service nationally.

Previous studies, which have often focused on particular populations or subsets of A&E visits and admissions, may mislead national policy makers if the overall associations in the whole population are much smaller than in the study population. For example, a quasi-experimental study of interventions to improve access to general practice in Greater Manchester, England, estimated a 26% reduction in 'minor' A&E visits but a statistically insignificant reduction of 3% across all A&E visits.⁷⁰ A similar analysis in London, England, estimated a 10% decrease in A&E visits in four practices that extended opening times, but it found no effect in 30 neighbouring practices whose patients could use the practices with longer opening hours.⁷¹ It is important to consider the magnitudes of associations in whole populations when assessing this research area.

Strengths and limitations

A strength of our study is that we used national data including all general practices in England. The findings are therefore externally valid and directly relevant to national policy makers, in contrast to previous local analyses.^{19-21 70 71} We also used several years of data which allowed us to test the sensitivity of results to different assumptions about unmeasured confounders at the practice level. Most previous national analyses cited above have often used just one data year or assumed that there is no confounding by time-constant practice level variables which is

unlikely. We used three data years because GPPS methods changed substantially from 2010-11 to 2011-12, and the most recent HES data we could obtain was for 2013-14. Though we adjusted results for possible confounding factors that are constant within practices over time, the results may still be biased by unobserved confounders that vary within practices over time. We were unable to account for this form of confounding using our chosen study design. Data on the quality of mental health services and social care, for example, would have been useful to incorporate. Since our analysis was conducted at the practice level, associations should not be inferred at the patient level (to avoid ecological fallacy). We could not conduct analysis at the patient level as individual GPPS responses cannot be linked to HES data.

HES recorded 83% of visits across all types of A&E department in 2012-13.³⁸ If the percentages of visits missing for each practice were not associated with the patient experience measures, associations estimated for these measures will not be biased but the 95% confidence intervals will be wider than they would otherwise have been (as the standard errors will be biased upwards). This is not a critical concern here because the sample size was large so the confidence intervals remained small. Incomplete recording of visits by an A&E department should affect all practices within the local area relatively equally, so adjusting for practice location (as we did) should reduce the chances of bias being introduced by missing data. Moreover, incomplete recording may affect the same practice equally in each year of the study period, so the analysis is unlikely to be affected by this potential source of bias. The sensitivity analysis that included visits to emergency departments only (where record coverage is close to 100%³⁸) produced results that do not alter our overall conclusions.

We estimated potential changes in costs associated with A&E visits and emergency admissions using average reference costs. This may not accurately reflect the true change in costs if, for example, changes in patient experience of general practice disproportionately affected simpler, and therefore cheaper, use of services.

Though we acknowledge the potential problems associated with the survey response rate of 36%, any non-response bias that may result should also be fairly constant within practices over time (as the patient populations do not change quickly) and therefore have little effect on the results. Moreover, a previous analysis of GPPS data suggested minimal associations between response rates and the outcome measures studied.⁷² Meta-analysis of survey literature indicates that associations between response rates and non-response bias are generally weak when probability sampling methods like those in the GPPS are used.⁷³

Changes in patient experience within practices between years are measured with error due to sampling variation in the GPPS. This measurement error will have attenuated associations between patient experience measures and rates of A&E visits and emergency admissions; the analysis could therefore underestimate the expected reductions in A&E visits and emergency admissions, and associated costs, with improvements in experiences of making appointments, for example.

Implications for research and policy

Our results suggest that only patient experience of making appointments at general practices was consistently associated with use of emergency hospital services, and this association was modest. These results cast doubt on interventions that are expected to substantially reduce use of emergency hospital services through improving patient experience of general practice. However, interventions may affect A&E visit and admission rates independently of patient experience as reported in the GPPS; the effects and causal mechanisms of any intervention are likely to depend largely on its unique design and context. Policies may also be implemented primarily for other reasons, such as improving patient experience of general practice in itself (which declined across several measures from 2011-12 to 2014-15).⁴⁸ Moreover, the results from trialling new policies now, despite an uncertain evidence base, may be better than the results of other policies introduced later on after waiting for more evidence to form. To conclude, we recommend caution in expectations that current plans to improve patient experience of general practice can substantially reduce use of emergency hospital services and provide net cost savings to the NHS in England.

References

1. World Health Organization. The World Health Report 2000 - Health Systems: Improving Performance. 2000. http://www.who.int/whr/2000/en/whr00_en.pdf?ua=1.
2. World Health Organization. The World Health Report 2008 - Primary Health Care: Now More Than Ever. 2008. http://www.who.int/whr/2008/whr08_en.pdf.
3. Aysola J, Rhodes KV, Polsky D. Patient-centered Medical Homes and Access to Services for New Primary Care Patients. *Med Care* 2015;53:857-62.
4. David G, Gunnarsson C, Saynisch PA, Chawla R, Nigam S. Do patient-centered medical homes reduce emergency department visits? *Health Serv Res* 2015;50:418-39.
5. Rosenthal MB, Friedberg MW, Singer SJ, Eastman D, Li Z, Schneider EC. Effect of a multipayer patient-centered medical home on health care utilization and quality: the Rhode Island chronic care sustainability initiative pilot program. *JAMA Intern Med* 2013;173:1907-13.
6. Friedberg MW, Rosenthal MB, Werner RM, Volpp KG, Schneider EC. Effects of a Medical Home and Shared Savings Intervention on Quality and Utilization of Care. *JAMA Intern Med* 2015;175:1362-8.
7. Friedberg MW, Schneider EC, Rosenthal MB, Volpp KG, Werner RM. Association between participation in a multipayer medical home intervention and changes in quality, utilization, and costs of care. *JAMA* 2014;311:815-25.
8. Pines JM, Keyes V, van Hasselt M, McCall N. Emergency department and inpatient hospital use by Medicare beneficiaries in patient-centered medical homes. *Ann Emerg Med* 2015;65:652-60.
9. Rosenthal MB, Alidina S, Friedberg MW, Singer SJ, Eastman D, Li Z, et al. A Difference-in-Difference Analysis of Changes in Quality, Utilization and Cost Following the Colorado Multi-Payer Patient-Centered Medical Home Pilot. *J Gen Intern Med* 2016;31:289-96.
10. Lippi Bruni M, Mammi I, Ugolini C. Does the extension of primary care practice opening hours reduce the use of emergency services? *J Health Econ* 2016;50:144-55.
11. NHS England. General Practice Forward View. 2016. <https://www.england.nhs.uk/wp-content/uploads/2016/04/gpfv.pdf>.
12. Cowling TE, Gunning EJ. Access to general practice in England: political, theoretical, and empirical considerations. *Br J Gen Pract* 2016;66:e680-2. doi: 10.3399/bjgp16X686977.
13. Cowling TE, Harris MJ, Majeed A. Evidence and rhetoric about access to UK primary care. *BMJ* 2015;350:h1513. doi: 10.1136/bmj.h1513.
14. Department of Health. NHS Mandate 2016 to 2017. 2015. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/494485/NHSE_mandate_16-17_22_Jan.pdf.
15. Mott MacDonald. Prime Minister's Challenge Fund: Improving Access to General Practice - First Evaluation Report. 2015. <https://www.england.nhs.uk/wp-content/uploads/2015/10/pmcf-wv-one-eval-report.pdf>.
16. NHS England. A&E Attendances and Emergency Admissions 2016-17. 2016. <https://www.england.nhs.uk/statistics/statistical-work-areas/ae-waiting-times-and-activity/statistical-work-areasae-waiting-times-and-activityae-attendances-and-emergency-admissions-2016-17/>.

17. NHS Digital. Hospital Admitted Patient Care Activity, 2015-16. 2016.
<http://content.digital.nhs.uk/article/2021/Website-Search?productid=23488&q=admitted+patient+care&sort=Relevance&size=10&page=1#top>.
18. NHS Digital. Hospital Episode Statistics, Admitted Patient Care - England, 2014-15. 2015.
<http://content.digital.nhs.uk/article/2021/Website-Search?productid=19420&q=hospital+episode+statistics+admitted+patient+care&sort=Relevance&size=10&page=2#top>.
19. Harris MJ, Patel B, Bowen S. Primary care access and its relationship with emergency department utilisation: an observational, cross-sectional, ecological study. *Br J Gen Pract* 2011;61:e787-93. doi: 10.3399/bjgp11X613124.
20. Baker R, Bankart MJ, Rashid A, Banerjee J, Conroy S, Habiba M, et al. Characteristics of general practices associated with emergency-department attendance rates: a cross-sectional study. *BMJ Qual Saf* 2011;20:953-8.
21. Bankart MJ, Baker R, Rashid A, Habiba M, Banerjee J, Hsu R, et al. Characteristics of general practices associated with emergency admission rates to hospital: a cross-sectional study. *Emerg Med J* 2011;28:558-63.
22. Cecil E, Bottle A, Cowling TE, Majeed A, Wolfe I, Saxena S. Primary Care Access, Emergency Department Visits, and Unplanned Short Hospitalizations in the UK. *Pediatrics* 2016;137. doi: 10.1542/peds.2015-1492.
23. Bottle A, Tsang C, Parsons C, Majeed A, Soljak M, Aylin P. Association between patient and general practice characteristics and unplanned first-time admissions for cancer: observational study. *Br J Cancer* 2012;107:1213-9.
24. Calderón-Larrañaga A, Soljak M, Cowling TE, Gaitatzis A, Majeed A. Association of primary care factors with hospital admissions for epilepsy in England, 2004-10: national observational study. *Seizure* 2014;23:657-61.
25. Calderon-Larranaga A, Carney L, Soljak M, Bottle A, Partridge M, Bell D, et al. Association of population and primary healthcare factors with hospital admission rates for chronic obstructive pulmonary disease in England: national cross-sectional study. *Thorax* 2011;66:191-6.
26. Brettell R, Soljak M, Cecil E, Cowie MR, Tuppin P, Majeed A. Reducing heart failure admission rates in England 2004-2011 are not related to changes in primary care quality: national observational study. *Eur J Heart Fail* 2013;15:1335-42.
27. Calderon-Larranaga A, Soljak M, Cecil E, Valabhji J, Bell D, Prados Torres A, et al. Does higher quality of primary healthcare reduce hospital admissions for diabetes complications? A national observational study. *Diabet Med* 2014;31:657-65.
28. Soljak M, Calderon-Larranaga A, Sharma P, Cecil E, Bell D, Abi-Aad G, et al. Does higher quality primary health care reduce stroke admissions? A national cross-sectional study. *Br J Gen Pract* 2011;61:e801-7. doi: 10.3399/bjgp11X613142.
29. Kasteridis P, Mason AR, Goddard MK, Jacobs R, Santos R, McGonigal G. The influence of primary care quality on hospital admissions for people with dementia in England: a regression analysis. *PLoS One* 2015;10:e0121506. doi: 10.1371/journal.pone.0121506.
30. McKay AJ, Newson RB, Soljak M, Riboli E, Car J, Majeed A. Are primary care factors associated with hospital episodes for adverse drug reactions? A national observational study. *BMJ Open* 2015;5:e008130. doi: 10.1136/bmjopen-2015-008130.
31. Fleetcroft R, Noble M, Martin A, Coombes E, Ford J, Steel N. Emergency hospital admissions for asthma and access to primary care: cross-sectional analysis. *Br J Gen Pract* 2016. doi: 10.3399/bjgp16X686089.

32. Cowling TE, Cecil EV, Soljak MA, Lee JT, Millett C, Majeed A, et al. Access to primary care and visits to emergency departments in England: a cross-sectional, population-based study. *PLoS One* 2013;8:e66699. doi: 10.1371/journal.pone.0066699.
33. Tammes P, Morris RW, Brangan E, Checkland K, England H, Huntley A, et al. Exploring the relationship between general practice characteristics, and attendance at walk-in centres, minor injuries units and EDs in England 2012/2013: a cross-sectional study. *Emerg Med J* 2016. doi: 10.1136/emered-2015-205339.
34. Scantlebury R, Rowlands G, Durbaba S, Schofield P, Sidhu K, Ashworth M. Socioeconomic deprivation and accident and emergency attendances: cross-sectional analysis of general practices in England. *British Journal of General Practice* 2015;65:e649-54. doi: 10.3399/bjgp15X686893.
35. Department of Health. NHS Outcomes Framework 2016 to 2017. 2016. <https://www.gov.uk/government/publications/nhs-outcomes-framework-2016-to-2017>.
36. Cowling TE, Harris MJ, Watt HC, Gibbons DC, Majeed A. Access to general practice and visits to accident and emergency (A&E) departments in England: cross-sectional analysis of a national patient survey. *Br J Gen Pract* 2014;64:e434-9. doi: 10.3399/bjgp14X680533.
37. Agarwal S, Banerjee J, Baker R, Conroy S, Hsu R, Rashid A, et al. Potentially avoidable emergency department attendance: interview study of patients' reasons for attendance. *Emerg Med J* 2012;29:e3. doi: 10.1136/emered-2011-200585.
38. NHS Digital. Accident and Emergency Attendances in England - 2013-14. 2015. <http://content.digital.nhs.uk/searchcatalogue?productid=17200&q=Accident+and+Emergency+Attendances+in+England&sort=Relevance&size=10&page=1#top>.
39. Aylin P, Bottle A, Middleton S. HSMR Mortality Indicators Full Methodology. 2011. http://www.drfooster.com/wp-content/uploads/2014/09/HSMR_Toolkit_Version_9_July_2014.pdf.
40. Ipsos MORI. GP Patient Survey - Technical annex: 2013-14 annual report. 2014. http://gp-survey-production.s3.amazonaws.com/archive/2014/July/1301375001_Technical%20Annex%202013-2014_FINAL%20v1.pdf.
41. Ipsos MORI. Technical Annex for the GP Patient Survey - 2011-2012 Annual Report. 2012. <http://gp-survey-production.s3.amazonaws.com/archive/2012/June/June%202012%20Technical%20Annex.pdf>.
42. Ipsos MORI. Technical Annex for the GP Patient Survey - 2012-2013 Annual Report. 2013. <http://gp-survey-production.s3.amazonaws.com/archive/2013/June/June%202013%20Technical%20Annex.pdf>.
43. Lyratzopoulos G, Elliott MN, Barbiere JM, Staetsky L, Paddison CA, Campbell J, et al. How can health care organizations be reliably compared?: Lessons from a national survey of patient experience. *Med Care* 2011;49:724-33.
44. Lyratzopoulos G, Elliott M, Barbiere JM, Henderson A, Staetsky L, Paddison C, et al. Understanding ethnic and other socio-demographic differences in patient experience of primary care: evidence from the English General Practice Patient Survey. *BMJ Qual Saf* 2012;21:21-9.
45. Warren FC, Abel G, Lyratzopoulos G, Elliott MN, Richards S, Barry HE, et al. Characteristics of service users and provider organisations associated with experience of out of hours general practitioner care in England: population based cross sectional postal questionnaire survey. *BMJ* 2015;350:h2040. doi: 10.1136/bmj.h2040.
46. Paddison C, Elliott M, Parker R, Staetsky L, Lyratzopoulos G, Campbell JL, et al. Should measures of patient experience in primary care be adjusted for case mix? Evidence from the English General Practice Patient Survey. *BMJ Qual Saf* 2012;21:634-40.

47. Paddison CA, Abel GA, Roland MO, Elliott MN, Lyratzopoulos G, Campbell JL. Drivers of overall satisfaction with primary care: evidence from the English General Practice Patient Survey. *Health Expect* 2013;18:1081-92.
48. Cowling TE. *Access to Primary Health Care in England: Policy, Theory, and Evidence*. PhD thesis. Imperial College London; 2016. <https://spiral.imperial.ac.uk/handle/10044/1/42497>.
49. NHS Digital. General and Personal Medical Services, England - 2003-13, As at 30 September. 2014. <http://content.digital.nhs.uk/searchcatalogue?productid=14458&q=general+and+personal+medical+services&sort=Relevance&size=10&page=2#top>.
50. Dusheiko M, Doran T, Gravelle H, Fullwood C, Roland M. Does higher quality of diabetes management in family practice reduce unplanned hospital admissions? *Health Serv Res* 2011;46:27-46.
51. Office for National Statistics. 2011 Census: QS211EW Ethnic group (detailed), Middle Layer Super Output Areas (MSOAs) and Lower Layer Super Output Areas (LSOAs) in England and Wales. 2013. <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/datasets/2011censuskeystatisticsandquickstatisticsforwardsandoutputareasinenglandandwales>.
52. NHS Digital. Numbers of Patients Registered at a GP Practice - January 2014. 2014. <http://content.digital.nhs.uk/searchcatalogue?productid=14010&q=Numbers+of+Patients+Registered+at+a+GP+Practice++April+2014&sort=Title&size=100&page=1#top>.
53. Department for Communities and Local Government. English indices of deprivation 2010. 2011. <https://www.gov.uk/government/statistics/english-indices-of-deprivation-2010>.
54. NHS Digital. Quality and Outcomes Framework (QOF) - 2013-14 2014. <http://content.digital.nhs.uk/searchcatalogue?productid=16273&q=quality+and+outcomes+framework&topics=1%2fPrimary+care+services%2fQuality+Outcomes+Framework&sort=Title&size=100&page=1#top>.
55. NHS Digital. Quality and Outcomes Framework - 2012-13. 2013. <http://content.digital.nhs.uk/searchcatalogue?productid=12972&q=quality+and+outcomes+framework&topics=1%2fPrimary+care+services%2fQuality+Outcomes+Framework&sort=Title&size=100&page=1#top>.
56. NHS Digital. Quality and Outcomes Framework - 2011-12, Practice level. 2012. <http://content.digital.nhs.uk/searchcatalogue?productid=9587&q=quality+and+outcomes+framework&topics=1%2fPrimary+care+services%2fQuality+Outcomes+Framework&sort=Title&size=100&page=1#top>.
57. Llanwarne NR, Abel GA, Elliott MN, Paddison CA, Lyratzopoulos G, Campbell JL, et al. Relationship between clinical quality and patient experience: analysis of data from the English Quality and Outcomes Framework and the National GP Patient Survey. *Ann Fam Med* 2013;11:467-72.
58. NHS Digital. General Practice Contracts 2011. 2012. <https://indicators.ic.nhs.uk/webview/>.
59. Office for National Statistics. 2011 rural/urban classification. 2015. <http://www.ons.gov.uk/ons/guide-method/geography/products/area-classifications/2011-rural-urban/index.html>.
60. Gutacker N, Mason AR, Kendrick T, Goddard M, Gravelle H, Gilbody S, et al. Does the quality and outcomes framework reduce psychiatric admissions in people with serious mental illness? A regression analysis. *BMJ Open* 2015;5:e007342. doi: 10.1136/bmjopen-2014-007342.
61. Wooldridge J. *Econometric analysis of cross section and panel data*. 2nd ed. Cambridge, Massachusetts: MIT Press, 2010.
62. Department of Health. NHS reference costs 2014 to 2015. 2015. <https://www.gov.uk/government/publications/nhs-reference-costs-2014-to-2015>.

63. Cowling TE, Harris MJ, Majeed A. Extended opening hours and patient experience of general practice in England: multilevel regression analysis of a national patient survey. *BMJ Qual Saf* 2016. doi: 10.1136/bmjqs-2016-005233.
64. Aboulghate A, Abel G, Elliott MN, Parker RA, Campbell J, Lyratzopoulos G, et al. Do English patients want continuity of care, and do they receive it? *Br J Gen Pract* 2012;62:e567-75. doi: 10.3399/bjgp12X653624.
65. Stata Statistical Software: Release 13 [program]. MP 13.1 version. College Station, TX: StataCorp LP, 2013.
66. National Audit Office. Improving patient access to general practice. 2017. <https://www.nao.org.uk/wp-content/uploads/2017/01/Improving-patient-access-general-practice.pdf>.
67. Plsek PE, Greenhalgh T. Complexity science: The challenge of complexity in health care. *BMJ* 2001;323:625-8.
68. Zhou Y, Abel G, Warren F, Roland M, Campbell J, Lyratzopoulos G. Do difficulties in accessing in-hours primary care predict higher use of out-of-hours GP services? Evidence from an English National Patient Survey. *Emerg Med J* 2015;32:373-8.
69. Cowling TE, Harris M, Watt H, Soljak M, Richards E, Gunning E, et al. Access to primary care and the route of emergency admission to hospital: retrospective analysis of national hospital administrative data. *BMJ Qual Saf* 2015;25:432-40.
70. Whittaker W, Anselmi L, Kristensen SR, Lau Y-S, Bailey S, Bower P, et al. Associations between Extending Access to Primary Care and Emergency Department Visits: A Difference-In-Differences Analysis. *PLoS Med* 2016;13:e1002113. doi: 10.1371/journal.pmed.1002113.
71. Dolton P, Pathania V. Can Increased Primary Care Access Reduce Demand for Emergency Care? Evidence from England's 7-Day GP Opening. *Journal of Health Economics* 2016. doi: 10.1016/j.jhealeco.2016.05.002.
72. Roland M, Elliott M, Lyratzopoulos G, Barbiere J, Parker RA, Smith P, et al. Reliability of patient responses in pay for performance schemes: analysis of national General Practitioner Patient Survey data in England. *BMJ* 2009;339:b3851. doi: 10.1136/bmj.b3851.
73. Groves RM, Peytcheva E. The Impact of Nonresponse Rates on Nonresponse Bias: A Meta-Analysis. *Public Opinion Quarterly* 2008;72:167-89.

Figure 1. Distribution of mean experiences of making appointments for general practices in England, 2011-14 (n=23,875 practice years)

54 practice-years (0.2%) with values less than 45 (minimum=30.3) not charted.




Table 1. Descriptive statistics for patient experience measures and rates of A&E visits and emergency admissions for general practices in England, 2011-14

	Mean	SD	Minimum	Maximum
2011-12 (n=8,075)				
Rate of A&E visits	310.3	107.2	26.6	1194.3
Rate of emergency admissions	98.8	27.5	15.0	413.2
Experience of making an appointment	77.8	8.5	40.2	98.2
Satisfaction with opening hours	79.4	6.1	46.8	98.7
Overall experience	82.8	6.1	55.5	98.3
2012-13 (n=7,950)				
Rate of A&E visits	324.4	115.0	36.3	1318.2
Rate of emergency admissions	100.3	26.9	3.0	405.9
Experience of making an appointment	76.0	9.2	34.4	99.2
Satisfaction with opening hours	78.5	6.3	47.1	98.5
Overall experience	81.7	6.5	49.1	99.1
2013-14 (n=7,850)				
Rate of A&E visits	324.4	109.7	19.4	1157.0
Rate of emergency admissions	102.9	26.8	3.8	339.5
Experience of making an appointment	74.8	9.5	30.3	99.0
Satisfaction with opening hours	76.6	6.6	46.0	98.5
Overall experience	81.1	6.7	46.9	98.7

A&E=accident and emergency, SD=standard deviation.

Total number of practices across period was 8,124.

Rates of A&E visits and emergency admissions are numbers per 1,000 registered patients.

Table 2. Associations between rates of A&E visits and fifths of experience of making an appointment, satisfaction with opening hours, and overall experience for general practices in England, 2011-14

	Unadjusted RR (95% CI)	Adjusted RR (95% CI)
Experience of making an appointment		
30.3 to 68.8	1 (ref.)	1 (ref.)
68.8 to 74.5	0.980 (0.972 to 0.987)***	0.996 (0.988 to 1.004)
74.5 to 79.2	0.965 (0.957 to 0.974)***	0.991 (0.981 to 1.001)
79.2 to 84.1	0.948 (0.939 to 0.957)***	0.985 (0.973 to 0.996)*
84.1 to 99.2	0.933 (0.922 to 0.943)***	0.979 (0.966 to 0.993)**
Satisfaction with opening hours		
46.0 to 73.0	1 (ref.)	1 (ref.)
73.0 to 77.0	0.984 (0.978 to 0.990)***	0.999 (0.993 to 1.006)
77.0 to 80.1	0.972 (0.966 to 0.978)***	0.999 (0.992 to 1.006)
80.1 to 83.6	0.966 (0.959 to 0.973)***	1.003 (0.995 to 1.011)
83.6 to 98.7	0.954 (0.944 to 0.963)***	1.002 (0.991 to 1.012)
Overall experience		
46.9 to 76.7	1 (ref.)	1 (ref.)
76.7 to 80.9	0.984 (0.977 to 0.992)***	1.004 (0.996 to 1.012)
80.9 to 84.1	0.974 (0.966 to 0.982)***	1.007 (0.999 to 1.016)
84.1 to 87.4	0.962 (0.953 to 0.971)***	1.007 (0.997 to 1.018)
87.4 to 99.1	0.951 (0.941 to 0.961)***	1.009 (0.996 to 1.021)

A&E=accident and emergency, CI=confidence interval, ref.=reference category, RR=rate ratio.

Unadjusted rate ratios estimated using random-effects Poisson regression with no other explanatory variables.

Adjusted rate ratios estimated using random-effects Poisson regression with all explanatory variables included.

In adjusted analyses, the number of included practice-years was 23,334 (97.7% of eligible practice-years).

* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Table 3. Associations between rates of emergency admissions rates and fifths of experience of making an appointment, satisfaction with opening hours, and overall experience for general practices in England, 2011-14

	Unadjusted RR (95% CI)	Adjusted RR (95% CI)
Experience of making an appointment		
30.3 to 68.8	1 (ref.)	1 (ref.)
68.8 to 74.5	0.978 (0.973 to 0.983) ^{***}	0.989 (0.983 to 0.994) ^{***}
74.5 to 79.2	0.970 (0.964 to 0.977) ^{***}	0.984 (0.977 to 0.991) ^{***}
79.2 to 84.1	0.955 (0.948 to 0.963) ^{***}	0.976 (0.968 to 0.984) ^{***}
84.1 to 99.2	0.939 (0.929 to 0.948) ^{***}	0.961 (0.951 to 0.971) ^{***}
Satisfaction with opening hours		
46.0 to 73.0	1 (ref.)	1 (ref.)
73.0 to 77.0	0.983 (0.978 to 0.988) ^{***}	0.998 (0.992 to 1.003)
77.0 to 80.1	0.974 (0.969 to 0.979) ^{***}	0.998 (0.992 to 1.004)
80.1 to 83.6	0.965 (0.959 to 0.971) ^{***}	0.997 (0.990 to 1.004)
83.6 to 98.7	0.953 (0.946 to 0.960) ^{***}	0.994 (0.986 to 1.003)
Overall experience		
46.9 to 76.7	1 (ref.)	1 (ref.)
76.7 to 80.9	0.987 (0.982 to 0.993) ^{***}	1.002 (0.997 to 1.008)
80.9 to 84.1	0.983 (0.977 to 0.990) ^{***}	1.007 (1.000 to 1.014) [*]
84.1 to 87.4	0.973 (0.965 to 0.980) ^{***}	1.005 (0.997 to 1.013)
87.4 to 99.1	0.963 (0.955 to 0.972) ^{***}	1.004 (0.994 to 1.015)

CI=confidence interval, RR=rate ratio.

Unadjusted rate ratios estimated using random-effects Poisson regression with no other explanatory variables.

Adjusted rate ratios estimated using random-effects Poisson regression with all explanatory variables included.

In adjusted analyses, the number of included practice-years was 23,334 (97.7% of eligible practice-years).

* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Table 4. Sensitivity analyses: adjusted rate ratios for A&E visits and emergency admissions comparing English general practices in the highest and lowest performing fifths of patient experience measures, 2011-14

	A&E visits			Emergency admissions		
	RR	95% CI		RR	95% CI	
Main analysis						
Experience of making an appointment	0.979	0.966	0.993	0.961	0.951	0.971
Satisfaction with opening hours	1.002	0.991	1.012	0.994	0.986	1.003
Overall experience	1.009	0.996	1.021	1.004	0.994	1.015
Sensitivity analysis 1*						
Experience of making an appointment	0.977	0.956	0.998	-	-	-
Satisfaction with opening hours	0.998	0.982	1.014	-	-	-
Overall experience	1.029	1.005	1.052	-	-	-
Sensitivity analysis 2†						
Experience of making an appointment	0.977	0.964	0.990	0.958	0.948	0.969
Satisfaction with opening hours	1.002	0.992	1.013	0.996	0.987	1.005
Overall experience	1.006	0.994	1.019	1.008	0.997	1.019
Sensitivity analysis 3‡						
Experience of making an appointment	0.981	0.967	0.994	0.967	0.956	0.977
Satisfaction with opening hours	1.004	0.992	1.015	0.999	0.990	1.008
Overall experience	1.020	1.008	1.033	1.001	0.992	1.011
Sensitivity analysis 4§						
Experience of making an appointment	0.979	0.966	0.993	0.962	0.952	0.972
Satisfaction with opening hours	1.000	0.990	1.011	0.994	0.986	1.002
Overall experience	1.006	0.994	1.019	1.005	0.995	1.015
Sensitivity analysis 5**						
Experience of making an appointment	0.981	0.968	0.995	0.967	0.956	0.978
Satisfaction with opening hours	1.002	0.991	1.013	0.993	0.985	1.002
Overall experience	1.010	0.997	1.022	1.006	0.995	1.017

CI=confidence interval, RR=rate ratio.

Results given in bold are statistically significant at the 5% level.

*Includes only visits to emergency departments (type 1 A&E departments) in the outcome variable.

†Includes only A&E visits and emergency admissions for patients aged 18 years old or over.

‡Used shrunken case-mix adjusted patient experience measures as explanatory variables.

§Adjusted for additional practice characteristics.

**Estimated using variation within practices between financial years only.

Table 5. Estimated changes in numbers of A&E visits and emergency admissions, and associated provider costs, with 0.5 and 1 SD increases in practice mean experiences of making appointments

	Number of A&E visits	Difference in number of A&E visits (95% CI)	Difference in cost (£) (95% CI)
No change	16,543,210	-	-
0.5 SD increase	16,391,931	-151,279 (-203,492 to -98,898)	-19,968,828 (-26,860,944 to -13,054,536)
1 SD increase	16,242,036	-301,174 (-404,481 to -197,205)	-39,754,968 (-53,391,492 to -26,031,060)
	Number of emergency admissions	Difference in number of emergency admissions (95% CI)	Difference in cost (£) (95% CI)
No change	5,423,838	-	-
0.5 SD increase	5,386,404	-37,434 (-50,410 to -24,427)	-83,590,122 (-112,565,530 to -54,545,491)
1 SD increase	5,349,228	-74,610 (-100,351 to -48,744)	-166,604,130 (-224,083,783 to -108,845,352)

A&E=accident and emergency, CI=confidence interval, SD=standard deviation.

Appendix 1. Definition of patient experience measures at the respondent level in the General Practice Patient Survey 2011-14

General Practice Patient Survey question	Answer value*
Overall, how would you describe your experience of making an appointment?	
Very good	100
Fairly good	75
Neither good nor poor	50
Fairly poor	25
Very poor	0
How satisfied are you with the hours that your GP surgery is open?†	
Very satisfied	100
Fairly satisfied	75
Neither satisfied nor dissatisfied	50
Fairly dissatisfied	25
Very dissatisfied	0
Overall, how would you describe your experience of your GP surgery?	
Very good	100
Fairly good	75
Neither good nor poor	50
Fairly poor	25
Very poor	0

GP=general practitioner.

All respondents were asked to complete each of the tabulated questions.

*Values were those assigned after linearly rescaling each measure.

†Responses of 'I'm not sure when my GP surgery is open' were excluded from analysis (n=80 636, 2.8%).

Appendix 2. Quality and Outcomes Framework indicators used to generate a composite measure of performance across intermediate clinical outcomes.

	2011-12	2012-13	2013-14	Points
Coronary heart disease				
The percentage of patients with coronary heart disease in whom the last blood pressure reading is 150/90 mmHg or less*	CHD06	CHD06	CHD002	17
The percentage of patients with coronary heart disease whose last measured total cholesterol is 5 mmol/l or less*	CHD08	CHD08	CHD003	17
Stroke and transient ischaemic attack (TIA)				
The percentage of patients with a history of stroke or TIA in whom the last blood pressure reading is 150/90 mmHg or less*	STROKE06	STROKE06	STIA003	5
The percentage of patients with stroke or TIA whose last measured total cholesterol is 5 mmol/l or less*†	STROKE08	STROKE08	STIA005	5
Hypertension				
The percentage of patients with hypertension in whom the last blood pressure reading is 150/90 mmHg or less	BP05	BP05	HYP002	10
Diabetes				
The percentage of patients with diabetes whose last measured total cholesterol is 5mmol/l or less*	DM17	DM17	DM004	6
The percentage of patients with diabetes in whom the last IFCC-HbA1c reading is 59 mmol/mol or less*	DM26	DM26	DM007	17
The percentage of patients with diabetes in whom the last IFCC-HbA1c reading is 64 mmol/mol or less*	DM27	DM27	DM008	8
The percentage of patients with diabetes in whom the last IFCC-HbA1c reading is 75 mmol/mol or less*	DM28	DM28	DM009	10
The percentage of patients with diabetes in whom the last blood pressure reading is 150/90 mmHg or less*	DM30	DM30	DM002	8
The percentage of patients with diabetes in whom the last blood pressure reading is 140/80 mmHg or less*	DM31	DM31	DM003	10
Epilepsy				
The percentage of patients aged 18 years or over on drug treatment for epilepsy who have been seizure free for the last 12 months*	EPILEPSY 08	EPILEPSY 08	EP002	6
Chronic kidney disease				
The percentage of patients on the CKD register in whom the last blood pressure reading is 140/85 mmHg or less*	CKD03	CKD03	CKD002	11

Intermediate outcome measures for peripheral arterial disease (PAD002 and PAD003) were introduced to the QOF in 2012-13, and an additional measure for hypertension (HYP003) was introduced in 2013-14 (not used in analysis). An intermediate outcome measure for mental health (MH18/MH010) was not included, as it applied to few patients in each practice.

Points are those available in QOF 2013-14; 57 and 55 points were available for BP05 in 2011-12 and 2012-13 respectively.

*Last reading recorded in the preceding 15 months in QOF 2011-12 and 2012-13 and in the preceding 12 months in QOF 2013-14.

†Measure changed for QOF 2013-14 to apply only to patients with a non-haemorrhagic stroke or a history of TIA.

Appendix 3. Descriptive statistics for patient experience measures from the General Practice Patient Survey, 2011-14

	Number of responses	Unweighted percentage of respondents (%)	Weighted percentage of respondents (%)
Satisfaction with opening hours*			
Very satisfied	1,235,576	44.8	40.0
Fairly satisfied	1,109,522	40.2	42.3
Neither satisfied nor dissatisfied	224,494	8.1	9.3
Fairly dissatisfied	132,747	4.8	5.9
Very dissatisfied	55,309	2.0	2.5
<i>Total</i>	<i>2,757,648</i>		
Experience of making an appointment			
Very good	1,176,083	42.4	35.7
Fairly good	1,080,176	38.9	41.2
Neither good nor poor	301,154	10.9	13.2
Fairly poor	145,114	5.2	6.6
Very poor	74,139	2.7	3.5
<i>Total</i>	<i>2,776,666</i>		
Overall experience			
Very good	1,452,265	51.2	44.8
Fairly good	1,080,961	38.1	42.2
Neither good nor poor	208,637	7.4	8.8
Fairly poor	71,511	2.5	3.2
Very poor	23,300	0.8	1.0
<i>Total</i>	<i>2,836,674</i>		

2,912,535 survey respondents from 8,289 general practices; data presented where available for each variable.

Weighted percentages account for survey design and non-response probabilities.

*Responses of 'I'm not sure when my GP surgery is open' were excluded (n=80 636, 2.8%).

Appendix 4. Descriptive statistics for characteristics of general practices in England measured as continuous variables, 2011-14, and pre-sample rates of A&E visits and emergency admissions

	Mean	SD	Minimum	Maximum
Rate of A&E visits, 2008-11	275.6	107.1	15.9	3,769.2
Rate of emergency admissions, 2008-11	98.6	30.3	21.5	846.2
Number of registered patients	6,940.3	4,219.9	1,001	52,609
Percentages of patients by age-gender (%):				
Male, 0 to 4	3.1	0.9	0.2	9.2
5 to 14	5.8	1.2	0.2	14.4
15 to 44*	20.9	4.7	8.8	60.5
45 to 64	13.0	2.2	1.7	25.4
65 to 74	4.2	1.6	0.1	11.7
75 to 84	2.4	1.0	0	8.5
85 or over	0.7	0.4	0	3.7
Female, 0 to 4	3.0	0.9	0.1	8.7
5 to 14	5.5	1.2	0.2	14.8
15 to 44	20.1	4.1	8.9	54.6
45 to 64	12.4	2.5	1.2	19.2
65 to 74	4.4	1.6	0.1	13.2
75 to 84	3.0	1.2	0	9.8
85 or over	1.4	0.7	0	8.2
Percentages of patients by ethnicity (%):				
White*	83.3	19.7	9.7	99.6
Mixed	2.4	1.8	0.2	11.3
Asian	9.1	13.3	0.1	76.3
Black	4.0	6.7	0.0	46.1
Other	1.2	1.8	0.0	18.8
QOF achievement	74.6	5.1	39.8	97.2

Data are for 23 875 practice-years and 8124 practices.

A&E=accident and emergency, QOF=Quality and Outcomes Framework, SD=standard deviation.

Rates of A&E visits and emergency admissions are numbers per 1000 registered patients.

Other variables adjusted for in regression models were: Index of Multiple Deprivation domain scores, practice contract type, urban/rural location, and Clinical Commissioning Group.

Statistics for Index of Multiple Deprivation domains have no natural interpretation (mean, SD): Income (0.16, 0.09); Employment (0.11, 0.05); Health and Disability (0.09, 0.72); Education, Skills, and Training (23.08, 13.79); Barriers to Housing and Services (22.33, 8.73); Crime (0.08, 0.62); Living Environment (23.77, 13.36).

*Variables used as reference categories in regression models and therefore omitted in model estimation.