

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Nalwoga, A; Cose, S; Nash, S; Miley, W; Asiki, G; Kusemererwa, S; Yarchoan, R; Labo, N; Whitby, D; Newton, R; (2018) Relationship between Anaemia, Malaria Co-infection and Kaposi Sarcoma-associated Herpesvirus (KSHV) Seropositivity in a Population-based Study in Rural Uganda. The Journal of infectious diseases. ISSN 0022-1899 DOI: <https://doi.org/10.1093/infdis/jiy274>

Downloaded from: <http://researchonline.lshtm.ac.uk/id/eprint/4647716/>

DOI: <https://doi.org/10.1093/infdis/jiy274>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

1 Relationship between anaemia, malaria co-infection and Kaposi Sarcoma-associated Herpesvirus
2 (KSHV) seropositivity in a population-based study in rural Uganda

3 Running title: Anaemia, malaria co-infection and KSHV seropositivity

4 *Angela Nalwoga^{1,2}, Stephen Cose^{1,2}, Stephen Nash², Wendell Miley³, Gershim Asiki⁴, Sylvia*
5 *Kusemererwa¹, Robert Yarchoan⁵, Nazzarena Labo³, Denise Whitby³, Robert Newton^{1,6}*

6 1. *MRC/UVRI and LSHTM Uganda Research Unit, Entebbe; Uganda*

7 2. *London School of Hygiene & Tropical Medicine, London; United Kingdom*

8 3. *Viral Oncology Section, AIDS and Cancer Virus Program, Leidos Biomedical Research, Inc.,*
9 *Frederick National Laboratory for Cancer Research, Frederick, MD; United States of America*

10 4. *African Population and Health Research Centre, Nairobi, Kenya*

11 5. *HIV and AIDS Malignancy Branch, National Cancer Institute, National Institutes of Health,*
12 *Bethesda, MD; United States of America*

13 6. *University of York, York; United Kingdom*

14 **Summary**

15 KSHV seroprevalence is very high in rural Uganda. Malaria infection and anaemia are the risk factors
16 associated with KSHV seroprevalence and antibody levels in this study. These factors might cause
17 reactivation of the virus and hence lead to increased transmission.

18

19 **Abstract**

20 We examined anaemia and malaria as risk factors for KSHV seropositivity and antibody levels in a
21 long-standing rural Ugandan cohort, in which KSHV is prevalent. Samples from 4134 children, aged
22 1-17 years, with a sex ratio of 1:1 and 3149 adults aged 18-103 years, 41% of whom were males,
23 were analysed. Among children, malaria infection was associated with higher KSHV prevalence (61%
24 versus 41% prevalence among malaria infected and uninfected respectively); malaria was not
25 assessed in adults. Additionally, lower haemoglobin level was associated with an increased
26 prevalence of KSHV seropositivity, both in children and in adults.

27 Word count: 97

28 **Keywords: Kaposi's sarcoma herpesvirus antibodies, rural population, anaemia, malaria**

29 **Introduction**

30 We recently reported KSHV prevalence of 95% among adults in a rural population cohort (the
31 General Population Cohort (GPC) in Uganda [1], the highest prevalence of KSHV ever reported, in
32 addition to high KSHV antibody levels. We propose that the very high prevalence in this population
33 may be driven by frequent KSHV reactivation, viral shedding and transmission rates and that the
34 high antibody levels also reflect frequent reactivation. It is important therefore to study potential co-
35 factors for reactivation in relation to KSHV prevalence and antibody levels. We have previously
36 reported an association between malaria and hookworm infections and KSHV seropositivity in an
37 urban population in Uganda [2, 3]. Since both malaria and hookworm are associated with anaemia,
38 we hypothesized that anaemia may have a role in KSHV transmission via viral reactivation. This
39 hypothesis is supported by data from *in vitro* experiments showing reactivation of KSHV in
40 conditions of hypoxia [4]. In this study we aimed to confirm the high prevalence of KSHV in the GPC
41 in recent years, with higher ART coverage, and determine the role of anaemia and malaria co-
42 infection as risk factors for KSHV prevalence and antibody levels in a highly endemic population.

43

44 **Methods**

45 **Study population and socio-demographic data collection**

46 The General Population Cohort (GPC) is located in south-western Uganda in Kyamulibwa sub-
47 community of Kalungu district with an altitude of approximately 1200m above sea level. It is
48 community-based open cohort of about 22,000 people in 25 adjacent villages [5]. This cross
49 sectional study analysed plasma samples collected from two surveys, the adult survey and the
50 children survey. The adults were surveyed in 2014/2015 and the children in 2016. Adults without
51 haematological parameter data and children without either haematological parameter data or
52 malaria parasitaemia status data were excluded in the laboratory analysis. Children less than 1 year
53 of age were excluded from the statistical analysis and children less than 2 years were not tested for
54 HIV serostatus, due to the potential for maternal IgG to be present, which could affect antibody

55 measurement. Socio-economic scores were generated for adults using Principal Component Analysis
56 of various household indicators during the previous survey.

57

58 **Ethical approval**

59 The study was approved by the Research and Ethics Committee Uganda Virus Research Institute and
60 the Uganda National Council for Science and Technology.

61

62 **Haematological and serological analysis**

63 During these two surveys, blood was collected from study participants and tested immediately after
64 collection for HIV; a smaller proportion of samples were also tested for malaria parasitaemia and
65 haemoglobin levels, using point-of care assays and rapid tests. HIV serostatus was determined using
66 rapid diagnostic tests. Malaria parasitaemia was measured in children only, using malaria Rapid
67 Diagnostic Tests (ONE STEP Malaria HRP-II (P.f) and pLDH (Pan) Antigen Rapid Test). Haemoglobin
68 levels in g/dL were obtained from the Haemocue 201 analyzer.

69

70 Stored plasma samples for both children and adults were retrieved and tested for KSHV antibodies
71 using an in-house ELISA as previously described [6]. Samples from the two surveys were tested
72 separately after simple randomisation onto ELISA plates. Antibodies to both K8.1 and ORF73
73 proteins were measured as optical density. Each ELISA plate contained three negative and positive
74 control wells; negative controls were used to calculate a cut-off value for every plate as previously
75 described [2, 7]. Seropositivity was defined as reactivity to either K8.1 or ORF73 proteins.

76

77 **Statistical analysis**

78 Statistical analysis was carried out using STATA13 (Statacorp, College Station, Texas USA). Children's
79 and adults' results were analysed separately. Haemoglobin levels were mean centred for easier
80 interpretation. Anaemia was defined using haemoglobin levels in g/dL after altitude adjustment

81 following WHO guidelines [8]. A constant value 0.5 was subtracted from haemoglobin levels for
82 altitude adjustment, the results were then categorised into normal and anaemic using the following
83 cut-off values: 11.0 for pregnant females and children below 5 years, 11.5 for children 5 to 11 years,
84 12.0 for children 12 to 14 years and other females 15 years and above, 13.0 for males 15 years and
85 above. These WHO haemoglobin reference ranges used to define anaemia may not be
86 representative of African populations, as previously reported [9, 10], because they are based on
87 western population data. We therefore analysed haemoglobin both as a continuous variable and as
88 categorised into normal and anaemic using separate regression models.

89

90 Linear regression with bootstrapped confidence intervals was used for antibody levels analysis,
91 because they were severely skewed. Logistic regression was used for seroprevalence analysis,
92 furthermore, we adjusted for clustering at the village level using survey commands. We assessed
93 interaction between age and haemoglobin levels, as well as between age and anaemia in relation to
94 anti-K8.1 antibody levels, anti-ORF73 antibody levels and KSHV prevalence based on *a priori*
95 suspicions of interaction, using likelihood ratio tests.

96

97 **Results**

98 The characteristics of the individuals analysed are shown in supplementary Table 1. We analysed
99 results from 3149 adults and 4134 children. This analysis included children aged 1 to 17 years and
100 adults aged 18 to 103 years (supplementary Table 1).

101

102 **Risk factors for KSHV prevalence and antibody levels among adults**

103 KSHV prevalence was 91% in all adults (2871/3149) (supplementary Figure 1). Every 1g/dL decrease
104 in haemoglobin values was associated with increased odds of being KSHV seropositive (OR=0.86
105 (0.77, 0.96), P=0.006 and anaemic individuals were more likely to be KSHV seropositive compared to

106 people with normal haemoglobin values, but this association was not statistically significant OR=1.25
107 (0.87, 1.79), P=0.229 (Table 1A).

108

109 We then analysed antibody levels to K8.1 and ORF73 proteins as continuous variables without
110 categorising participants as seropositive or seronegative. Anaemic adults had higher antibodies to
111 ORF73 protein compared to individuals with normal haemoglobin values (coef. 0.28 (0.16, 0.39),
112 $P < 0.0001$). Similarly, every 1g/dL decrease in haemoglobin was associated with an increase in ORF73
113 antibody ODs (Table 1B). The association between haemoglobin and antibodies to ORF73 protein
114 was strongest among older people (Table 1C). Conversely, anti-K8.1 antibody levels were not
115 significantly associated with either haemoglobin levels or anaemia (Table 1B). This might be
116 attributed to the relative abundance of LANA compared to late lytic proteins such as K8.1, even
117 during KS disease [11].

118

119 Compared to HIV negative adults, HIV positive adults had lower antibodies to KSHV, especially those
120 with CD4 counts of 500cell/ μ L or less (Table 1B). This may be due to B cell dysfunction caused by HIV
121 infection, and consequent decreased antibody responses [12, 13]. On the other hand, HIV positive
122 adults with CD4 counts above 500cells/ μ L, compared to HIV negative adults were more likely to be
123 KSHV seropositive (Table 1A), which may be due to antiretroviral treatment.

124

125 **Risk factors for KSHV prevalence and antibody levels among children**

126 We then investigated associations between KSHV prevalence and antibody levels and risk factors
127 among children. Overall, KSHV prevalence was 51% (2117/4134) in the children, the prevalence
128 increased with age, rising from 31% among 1-5 year olds, to 53% among 6-12 year olds, to 73%
129 among 13-17 year olds (supplementary Figure 1). We first adjusted for HIV status, age and sex, then
130 malaria parasitaemia and anaemia/haemoglobin levels were added in the full models. In the first
131 analysis, haemoglobin levels, malaria parasitaemia and age were strongly associated with KSHV

132 prevalence (Table 2A). Every 1 g/dL decrease in haemoglobin levels increased the odds of being
133 KSHV seropositive by 11% ($P<0.0001$) and the odds of being KSHV positive if anaemic compared to
134 with normal haemoglobin levels was 1.42 (1.18, 1.71) ($p<0.0001$). The odds of being KSHV
135 seropositive, if malaria infected, compared to the uninfected were 2.26 (1.85, 2.77) ($P<0.0001$) and
136 every annual increase in age was associated with a 17% increased risk of being KSHV seropositive
137 (Table 2A).

138

139 After adjusting for malaria parasitaemia, the risk of being KSHV seropositive for every 1 g/dL
140 decrease in haemoglobin reduced to 7% ($p=0.005$). Similarly, the odds of being KSHV seropositive in
141 comparing anaemic individuals to people with normal haemoglobin levels reduced to 1.23 (1.01,
142 1.49) $p=0.037$. After adjusting for haemoglobin levels, the odds of being KSHV seropositive
143 comparing people with and without malaria parasitaemia changed little (OR=2.12 (1.75, 2.57),
144 $p<0.0001$) (Table 2A). Every annual increase in age remained strongly associated with increased
145 KSHV prevalence risk, OR=1.18 ($P<0.0001$) even after adjusting for malaria parasitaemia and
146 haemoglobin (Table 2A).

147

148 We then finally investigated associations between the same risk factors and KSHV antibody levels
149 (OD) as continuous variables without categorising participants as seropositive or seronegative. Only
150 malaria parasitaemia was associated with both anti-K8.1 and anti-ORF73 antibody levels in the fully
151 multivariate analysis (Table 2B).

152

153 **Discussion**

154 We observed a significant association between haemoglobin levels and KSHV prevalence among
155 children and adults, where people with low levels of haemoglobin were more likely to be KSHV
156 seropositive. As a categorical variable, anaemia was associated with KSHV prevalence among
157 children. Reduction in haemoglobin has been shown to cause hypoxia/low tissue oxygen, while

158 hypoxia has been shown to reactivate KSHV *in-vitro* [4]. We therefore hypothesise that low
159 haemoglobin levels leads to reactivation of KSHV through hypoxia. Increased reactivation may help
160 spread the virus during initial infection. Alternately, hypoxia may enhance initial infection of cells,
161 possibly through upregulation of the replication and transcription activator [14]. In this cross
162 sectional study, we did not directly measure KSHV reactivation or KSHV viral load in blood or plasma,
163 although antibody levels may be viewed as a surrogate marker for frequent reactivation. The
164 connection between KSHV reactivation, hypoxia and anaemia requires further investigation.

165

166 We showed that children infected with malaria are more likely to be KSHV seropositive. Additionally,
167 the effect of anaemia and/or haemoglobin levels on KSHV prevalence and antibody levels reduced to
168 about 50% after adjusting for malaria infection. Malaria causes anaemia, and in part the anaemia
169 effect in children could be explained (confounded) by malaria infection. The consistent association
170 between malaria infection and KSHV prevalence suggests malaria may be driving KSHV transmission
171 in malaria endemic areas. This might imply that exposure to malaria significantly impacts on KSHV
172 reactivation, which might also have long lasting effects. The mechanism through which malaria may
173 reactivate KSHV requires further investigation.

174

175 **Conclusion**

176 Findings from this study suggest malaria infection as a risk factor for KSHV prevalence. Malaria
177 associated anaemia is one mechanism that likely contributes to this association but cannot entirely
178 explain it. In KSHV and malaria endemic areas, a number of other parasite co-infections such as
179 helminths, which cause anaemia and/or immunomodulation are common. The role of multiple
180 parasitic infections and KSHV transmission and pathogenesis warrants further careful study.

181

182

183 Word count: 1976

184 **References**

- 185 1. Newton, R., et al., *Kaposi's sarcoma associated herpesvirus in a rural Ugandan cohort: 1992-*
186 *2008*. J Infect Dis, 2017.
- 187 2. Wakeham, K., et al., *Parasite infection is associated with Kaposi's sarcoma associated*
188 *herpesvirus (KSHV) in Ugandan women*. Infect Agent Cancer, 2011. **6**(1): p. 15.
- 189 3. Wakeham, K., et al., *Risk factors for seropositivity to Kaposi sarcoma-associated herpesvirus*
190 *among children in Uganda*. J Acquir Immune Defic Syndr, 2013. **63**(2): p. 228-33.
- 191 4. Davis, D.A., et al., *Hypoxia induces lytic replication of Kaposi sarcoma-associated herpesvirus*.
192 *Blood*, 2001. **97**(10): p. 3244-50.
- 193 5. Asiki, G., et al., *The general population cohort in rural south-western Uganda: a platform for*
194 *communicable and non-communicable disease studies*. Int J Epidemiol, 2013. **42**(1): p. 129-
195 41.
- 196 6. Mbisa, G.L., et al., *Detection of antibodies to Kaposi's sarcoma-associated herpesvirus: a new*
197 *approach using K8.1 ELISA and a newly developed recombinant LANA ELISA*. J Immunol
198 *Methods*, 2010. **356**(1-2): p. 39-46.
- 199 7. Nalwoga, A., et al., *Association between malaria exposure and Kaposi's sarcoma-associated*
200 *herpes virus seropositivity in Uganda*. Trop Med Int Health, 2015. **20**(5): p. 665-672.
- 201 8. Organization, W.H., *Haemoglobin concentrations for the diagnosis of anaemia and*
202 *assessment of severity*. 2011. Download from: [http://www.who.](http://www.who.int/vmnis/indicators/haemoglobin.pdf)
203 [int/vmnis/indicators/haemoglobin.pdf](http://www.who.int/vmnis/indicators/haemoglobin.pdf), 2015.
- 204 9. Lugada, E.S., et al., *Population-based hematologic and immunologic reference values for a*
205 *healthy Ugandan population*. Clin Diagn Lab Immunol, 2004. **11**(1): p. 29-34.
- 206 10. Mugisha, J.O., J. Seeley, and H. Kuper, *Population based haematology reference ranges for*
207 *old people in rural South-West Uganda*. BMC Res Notes, 2016. **9**(1): p. 433.
- 208 11. Dupin, N., et al., *Distribution of human herpesvirus-8 latently infected cells in Kaposi's*
209 *sarcoma, multicentric Castleman's disease, and primary effusion lymphoma*. Proc Natl Acad
210 *Sci U S A*, 1999. **96**(8): p. 4546-51.
- 211 12. Abudulai, L.N., et al., *Chronic HIV-1 Infection Induces B-Cell Dysfunction That Is Incompletely*
212 *Resolved by Long-Term Antiretroviral Therapy*. J Acquir Immune Defic Syndr, 2016. **71**(4): p.
213 381-9.
- 214 13. Falconer, O., M.L. Newell, and C.E. Jones, *The Effect of Human Immunodeficiency Virus and*
215 *Cytomegalovirus Infection on Infant Responses to Vaccines: A Review*. Front Immunol, 2018.
216 **9**: p. 328.
- 217 14. Krishnan, H.H., et al., *Concurrent expression of latent and a limited number of lytic genes*
218 *with immune modulation and antiapoptotic function by Kaposi's sarcoma-associated*
219 *herpesvirus early during infection of primary endothelial and fibroblast cells and subsequent*
220 *decline of lytic gene expression*. J Virol, 2004. **78**(7): p. 3601-20.

221

222

223

224 Supplementary Figure 1 title: KSHV prevalence and 95% confidence intervals across ages 1 to 103
225 years among individuals from the General Population Cohort in south-western Uganda
226 Supplementary Figure 1 legend: KSHV seroprevalence defined as antibody reactivity to either K8.1 or
227 ORF73 antigens. Anti-K8.1 and anti-ORF antibodies detected using ELISA.
228

229 Footnote page

230 All authors have declared that they have no association which may cause any conflict of interest.

231 Funding

232 This research is partially funded by the African Partnership for Chronic Disease Research, University
233 of Cambridge, United Kingdom. It has been funded in part with federal funds from the National
234 Cancer Institute, National Institutes of Health, under Contract No. HHSN261200800001E. The
235 content of this publication does not necessarily reflect the views or policies of the Department of
236 Health and Human Services, nor does mention of trade names, commercial products, or
237 organizations imply endorsement by the U.S. Government. This research was supported in part by
238 the Intramural Research Program of the NIH, National Cancer Institute. The Ugandan General
239 Population Cohort study is jointly funded by the UK Medical Research Council (MRC) and the UK
240 Department for International Development (DFID) under the MRC/DFID Concordat agreement.

241 This work was presented at the 16th International Conference on Malignancies in HIV/AIDS, NIH Main
242 Campus, October 23-34, 2017

243 Corresponding author

244 Angela Nalwoga

245 MRC/UVRI and LSHTM Uganda Research Unit

246 P.O Box 49 Entebbe Uganda

247 Angela.nalwoga@mrcuganda.org; angelanalwoga@gmail.com

248 Alternative corresponding author

249 Robert Newton

- 250 MRC/UVRI and LSHTM Uganda Research Unit
- 251 P.O Box 49 Entebbe Uganda
- 252 rob.newton@mrcuganda.org; robert.newton@york.ac.uk
- 253
- 254