

1  
2  
3  
4  
5  
6  
7 **Comparing the clinical severity of disease caused by enteroviruses and human**  
8 **parechoviruses in neonates and infants**

9  
10 Sally Black<sup>1</sup>, Carina Bradley<sup>2</sup>, Florence Y Lai<sup>3</sup>, Savitha Shenoy<sup>1</sup>, Srinu Bandi<sup>1</sup>, David J  
11 Allen<sup>4,5</sup>, Julian W Tang<sup>2,6</sup>

12  
13  
14 <sup>1</sup>Leicester Children's Hospital, University Hospitals of Leicester NHS Trust, Leicester, UK

15  
16 <sup>2</sup>Clinical Microbiology, University Hospitals of Leicester NHS Trust, Leicester, UK

17  
18 <sup>3</sup>Department of Cardiovascular Science, University of Leicester, Leicester, UK

19  
20 <sup>4</sup>Virus Reference Department, National Infections Service, Public Health England, London,  
21 UK

22  
23 <sup>5</sup>Pathogen Molecular Biology Department, Faculty of Infectious and Tropical Diseases,  
24 London School of Hygiene and Tropical Medicine, UK

25  
26 <sup>6</sup>Infection, Immunity, Inflammation, University of Leicester, Leicester, UK

27  
28  
29  
30  
31  
32  
33  
34  
35  
36 **Correspondence to: Dr Julian W Tang**

37  
38 Clinical Microbiology, University Hospitals of Leicester NHS Trust

39  
40 Level 5 Sandringham Building, Leicester Royal Infirmary

41  
42 Infirmary Square, Leicester LE1 5WW, UK.

43  
44 Email: [julian.tang@uhl-tr.nhs.uk](mailto:julian.tang@uhl-tr.nhs.uk); [jwtang49@hotmail.com](mailto:jwtang49@hotmail.com)

45  
46  
47  
48 Tel: 0116 258 6516. Fax: 0116 255 1949

**ABSTRACT**

Comparison of children hospitalised with enterovirus (EV) or human parechovirus (HPeV) infections of their cerebrospinal fluid (CSF) revealed that HPeV infections presented with more persistent fever, irritability and feeding problems, more frequent leukopenia and lymphopenia, and higher admission rates to high dependency or intensive care units. As very few HPeV cases were followed-up, further studies on long-term outcomes are needed.

**Word count 60**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## HIGHLIGHTS

Children infected with HPeV had more persistent symptoms than those infected with EV

Children infected with HPeV had higher HDU/ICU admissions than those infected with EV

Children infected with HPeV had more leukopenia than those infected with EV

Children infected with EV had higher CSF white cells than those infected with EV

Further studies are needed to characterise any longer-term EV and HPeV complications

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## INTRODUCTION

1  
2 Enteroviruses are well-known causes of sepsis in neonates and infants. In recent  
3  
4 years, the extent to which parechoviruses may be contributing to neonatal and infant  
5  
6 morbidity and mortality has begun to emerge.<sup>1-3</sup>  
7  
8

9  
10 Enteroviruses (EV) and human parechoviruses (HPeV) are non-enveloped, single-  
11  
12 stranded, positive-sense RNA viruses and members of the Picornavirus family. They are  
13  
14 common causes of neonatal and infant sepsis, worldwide.  
15  
16

17 Enteroviruses exist as multiple serotypes, subdivided into various genus, including  
18  
19 echoviruses, Coxsackie A and B viruses, and the numbered enteroviruses. Human  
20  
21 parechoviruses exist in at least 17 genotypes, of which genotypes 1-6 are most commonly  
22  
23 found in humans, with genotype 3 being most commonly responsible for sepsis in neonates  
24  
25 and infants.  
26  
27

28  
29 Whilst most episodes of EV and HPeV neonatal and infant sepsis are self-limiting,  
30  
31 more severe illness can occur and there are current concerns about longer-term sequelae,  
32  
33 particularly in HPeV infections where there is more significant neurological involvement.  
34  
35 Previous studies have found that the clinical presentation of the two viruses are often  
36  
37 indistinguishable.<sup>1,3</sup>  
38  
39  
40

41 Our diagnostic virology laboratory has only relatively recently (since mid-2014)  
42  
43 introduced routine testing for parechoviruses as part of our neonatal and infant septic work-  
44  
45 up. We examined the demographics, laboratory results and clinical notes for paediatric  
46  
47 patients admitted with sepsis with laboratory-confirmed human enterovirus (EV) or  
48  
49 parechovirus (HPeV) infections of the cerebrospinal fluid (CSF), during Feb 2014 to Aug  
50  
51 2017.  
52  
53  
54  
55  
56  
57

## METHODS

58  
59  
60  
61  
62  
63  
64  
65

1 All cases from a 3.5 year period (Feb 2014 to Aug 2017) were selected on the basis of  
2 a positive cerebrospinal fluid (CSF) polymerase chain reaction (PCR) result for either EV or  
3 HPeV RNA, using assays previously described elsewhere,<sup>2,4</sup> which were performed as part of  
4 the routine workup for neonates or infants admitted with suspected sepsis. Additional  
5 samples were taken depending on the degree of clinical illness, including: EDTA blood,  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

This study was performed as part of a paediatric departmental audit which aimed to ensure that all EV and HPeV-infected paediatric patients had received appropriate follow-up after discharge for sepsis, where there was laboratory confirmed EV or HPeV infection of the CSF. Therefore formal ethics approval was not required.

Clinical parameters examined included: age at presentation, length of stay, fever, rash, seizures, respiratory difficulty, feeding problems, antimicrobial use, and admission to high dependency or intensive care units. Laboratory parameters compared included: C-reactive protein (CRP), white cell counts (WCC), liver function tests (LFT), CSF profile (glucose, protein, and cell counts), and radiological investigations, where available.

Clinical and laboratory characteristics were compared between patients with HPeV and EV infection. Continuous variables were presented as mean and standard deviation (or median and interquartile range if not normally distributed) and compared with Student t-test (or Wilcoxon test). Binary variables were presented as frequency and percentages, and compared with the Fisher Exact test. Multivariable analysis for risk ratios (RR) comparing HPeV infection to EV infection as the reference, were estimated using log-Binomial regression model.

## RESULTS

There were no statistically significant differences in age or sex of the children affected by EV vs HPeV CNS infections but there appeared to be a difference in range, with EV often affecting older children than HPeV (IQR 29-102 days for EV and 25.5-61 days for PeV).

Out of a total of 163 cases, there were 131 EV (i.e. 7 Coxsackie A, 18 Coxsackie B, 46 echoviruses, with 60 enteroviruses that could not be typed further) and 32 HPeV infections (**Table 1**). All HPeV infections were caused by HPeV genotype 3 (HPeV-3). Of the EV cases, 73% (95 cases) were in children younger than 90 days (3 months), whereas over 90% (30 cases) of HPeV cases were in children younger than 90 days (3 months).

Cases of enterovirus meningitis showed three peaks of activity each year with the most significant being in the Nov-Dec period. In contrast, HPeV had only one significant outbreak over two months in summer 2016 (**Figure S1**). There was no difference in the mean age or sex of the children affected by EV or PeV, although there was a difference in range, with EV meningitis affecting some much older children.

A greater number of abnormal parameters were found with HPeV than for enteroviruses, with a greater likelihood of admission to high dependency unit (HDU)/intensive care unit (ICU) ( $p=0.004$ ) and a higher rate of persistent symptoms (i.e. fever, irritability, and feeding problems,  $p<0.05$ ) (**Table 2**). Compared with children infected with EV, children with HPeV were more likely to have an abnormally low WCC (leukopenia) (56% HPeV vs. 14% EV,  $p<0.001$ ), and an abnormally low lymphocyte count (lymphopenia) (91% HPeV vs. 39% EV,  $p<0.001$ ) (**Table S1**).

In contrast, EV cases were more likely to have a high white cell count in the CSF (6% HPeV vs. 50% EV,  $p<0.001$ ) (**Table S1**). In the adjusted (log-Binomial regression) analysis, the HPeV cases were over 5 more times more likely to have lymphopenia than EV cases

1 (RR=5.11, 95% CI 1.53-17.05, p=0.008), with EV cases being marginally more likely to have  
2 a higher CSF WCC (RR=0.22, 95% CI 0.05-0.92, p=0.038) (**Table S2**).  
3

4 Other laboratory and clinical parameters, including overall length of stay (LOS) did  
5 not differ significantly between the EV or HPeV cases, however. There was no significant  
6 difference in whether or not EV or HPeV cases received antibiotic (98.4% vs. 100%,  
7 respectively, p=0.999) and/or acyclovir (37.1% vs. 34.4%, respectively, p=0.839) treatment  
8  
9  
10  
11  
12  
13  
14 (**Table 2**).  
15

16 Finally, relatively few patients were deemed to require longer-term followup. Of the  
17 total number of cases, 80% of children did not require any follow-up at one year post-  
18 infection. At one year post-infection 3% of children were under follow-up by ophthalmology,  
19 with no abnormalities detected. Sixteen children (~10%) attended a routine hearing check but  
20 none had any detectable sensorineural hearing loss. Only 3 patients (<2%) were reported as  
21 having had any developmental delay problems on admission: 1 child had delayed speech and  
22 manipulative skills – both of which resolved by one year post-infection. Another child still  
23 had some speech delay at one year followup, and one child had gross motor delay (despite a  
24 normal MRI). As 2 of the 3 children with developmental problems had uncomplicated, short  
25 inpatient stays, this might suggest that their viral infection were not direct causes of this.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41 However, this does not completely exclude this aetiological possibility.  
42  
43  
44  
45

## 46 **DISCUSSION**

47 Infections by EVs and HPeVs are well-documented causes of neonatal and infant  
48 sepsis. However relatively few studies have compared the severity of clinical illness caused  
49 by these viruses within the same paediatric population within the same season.<sup>5</sup>  
50  
51  
52  
53

54 Here we demonstrate differences in presentation and severity of these two viruses,  
55  
56 with HPeV cases having a higher likelihood of having persistent fevers (p<0.05), irritability  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 or feeding problems ( $p < 0.05$ ), leukopenia, lymphopenia, and requiring admission to HDU or  
2 PICU than children with EV infections. These findings are consistent with those reported  
3 from other studies.<sup>6-8</sup> In addition, more specifically, Cabrerizo and colleagues<sup>9</sup> also noted a  
4 higher CSF pleocytosis in EV vs. HPeV infections, as found in this study.  
5  
6  
7  
8

9 In our population more children aged 30-90 days ( $n = 19$ , 58%) were infected with HPeV  
10 than neonates ( $n = 11$ , 33%). Some studies have found children over the age of 2 months<sup>8</sup> or 3  
11 months<sup>9</sup> were unaffected by HPeV, whereas 21% of our cases (7 patients) were diagnosed in  
12 children aged over 2 months, with 2 cases being in a 4-month and 6-month-old, respectively.  
13 Thus, routine testing for HPeV in all children with febrile rash illness and sepsis may reveal a  
14 higher number of older children infected with HPeV.  
15  
16  
17  
18  
19  
20  
21  
22  
23

24 Although some previous studies have found pediatric HPeV and EV infections  
25 clinically indistinguishable,<sup>1,3</sup> anecdotally, in our pediatric population, nurses who worked  
26 with children involved in our recent HPeV outbreak,<sup>2</sup> reported that they were able to  
27 distinguish which children had HPeV rather than EV, prior to any laboratory confirmation, on  
28 their clinical presentation alone. These HPeV cases were noted to be generally more irritable  
29 and persistently inconsolable, tachycardic and pyrexial than the more frequently  
30 encountered annual, seasonal EV cases with which the nurses were very familiar.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 The main limitation of this study is related to the infrequent and sporadic approach to  
42 the longer-term followup of these EV and HPeV-infected patients, as individual clinical  
43 teams were left to decide on whether patients being admitted under them warranted such  
44 followup. This was mostly based on the individual patient's clinical course during their  
45 admission, as well as prior experiences of the lead paediatrician concerned, rather than any  
46 local consensus guidelines.  
47  
48  
49  
50  
51  
52  
53  
54

55 At present clinical guidelines do not differentiate between the management of  
56 children presenting with EV versus HPeV infections.<sup>3</sup> This study demonstrates that  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 differences in the severity of clinical illness can be seen between the HPeV and EV CNS  
2 infections, with a greater degree of severity in HPeV cases. Further studies are required to  
3  
4 clarify and confirm these findings, which may then lead to more practical clinical guidelines  
5  
6  
7 for the immediate and longer-term management and followup of these patients.  
8  
9

10  
11  
12  
13  
14 **Word count 1493**  
15

16  
17  
18  
19 **REFERENCES**  
20

- 21  
22 1. Verboon-Maciolek MA, Krediet TG, Gerards LJ, et al. Severe neonatal parechovirus  
23 infection and similarity with enterovirus infection. *Pediatr Infect Dis J.* 2008;27:241-  
24 245.  
25  
26  
27  
28 2. Tang JW, Holmes CW, Elsanousi FA, et al. Cluster of human parechovirus infections  
29 as the predominant cause of sepsis in neonates and infants, Leicester, United  
30 Kingdom, 8 May to 2 August 2016. *Euro Surveill.* 2016;21:pii=30326.  
31  
32  
33  
34 3. Drysdale SB, Kelly DF. Fifteen-minute consultation: enterovirus meningitis and  
35 encephalitis—when can we stop the antibiotics? *Arch Dis Child Educ Pract Ed.*  
36 2017;102:66–71.  
37  
38  
39 4. Holmes CW, Koo SS, Osman H, et al. Predominance of enterovirus B and echovirus  
40 30 as cause of viral meningitis in a UK population. *J Clin Virol.* 2016;81:90-93.  
41  
42  
43  
44 5. Walters B, Peñaranda SW, Nix AM et al. Detection of human parechovirus (HPeV)-3  
45 in spinal fluid specimens from pediatric patients in the Chicago area. *J Clin Virol.*  
46 2011;52:187-191.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
6. Sharp J, Harrison CJ, Puckett K, et al. Characteristics of young infants in whom human parechovirus, enterovirus or neither were detected in cerebrospinal fluid during sepsis evaluations. *Pediatr Infect Dis J.* 2013;32:213-216.
7. Sano K, Hamada H, Hirose S, et al. Prevalence and characteristics of human parechovirus and enterovirus infections in febrile infants. *Pediatr Int.* 2017 Dec 5. Doi:10.1111/ped.13467. [Epub ahead of print]
8. Harvala H, Calvert J, Van Nguyen D, et al. Comparison of diagnostic clinical samples and environmental sampling for enterovirus and parechovirus surveillance in Scotland, 2010 to 2012. *Euro Surveill.* 2014;19:pii=20772.
9. Cabrerizo M, Trallero G, Pena MJ, Cilla A, et al. Comparison of epidemiology and clinical characteristics of infections by human parechovirus vs. those by enterovirus during the first month of life. *Eur J Pediatr.* 2015;174:1511-1516.

**Table 1.** Specific human enteroviruses (EVs) by type and proportion, identified in the CSF of these patients

<b>EVs</b>	<b>Number</b>	<b>Percentage (%)</b>
CA16	1	0.76
CA6	2	1.53
CA9	4	3.05
CB1	1	0.76
CB4	7	5.34
CB5	10	7.63
E11	2	1.53
E16	3	2.29
E18	7	5.34
E21	1	0.76
E25	3	2.29
E3	1	0.76
E30	3	2.29
E5	5	3.82
E6	4	3.05
E7	5	3.82
E71	1	0.76
E9	10	7.63
EV7	1	0.76
Untypeable	60	45.80
<b>Total</b>	<b>131</b>	<b>100.00</b>

**TABLE 2.** Clinical and laboratory characteristics of the EV and HPeV cases

	<b>Enterovirus (EV) (n=131)</b>	<b>Parechovirus (HPeV) (n=32)</b>	<b>p-value</b>
<b><i>Demographics</i></b>			
Age (days)	50 (29 - 102)	39.5 (25.5 - 61)	0.069
Sex (female)	42.0% (55/131)	40.6% (13/32)	0.999
<b><i>Symptoms</i></b>			
Fever	94.6% (123/130)	100% (32/32)	0.347
Peak temperature (°C)	38.5 (0.7)	38.7 (0.7)	0.293
Feeding problems	56.6% (73/129)	68.8% (22/32)	0.234
Rash in history	25.2% (32/127)	25.0% (8/32)	0.999
Seizure	4.7% (6/127)	3.1% (1/32)	0.999
Respiratory symptoms (coryzal symptoms, grunting, cough, wheeze).	35.2% (45/128)	37.5% (12/32)	0.838
<b><i>Blood and CSF results</i></b>			
C-reactive protein (CRP, mg/L) (normal range: 0 - 10)	10 (3 - 24)	6 (3 - 13.5)	0.073
White cell count (WCC, x10 <sup>9</sup> /L) (normal range: 6.0 - 17.0)	10.5 (4.1)	6.4 (2.9)	<b>&lt;0.001</b>
Neutrophils (x10 <sup>9</sup> /L) (normal range: 1.50 - 8.50)	5.5 (3.4)	3.7 (2.5)	<b>&lt;0.001</b>
Lymphocytes (x10 <sup>9</sup> /L) (normal range: 4.00 - 13.50)	4.0 (2.1)	2.0 (0.8)	<b>&lt;0.001</b>
Platelets (x10 <sup>9</sup> /L) (normal range: 140 - 400)	390.1 (130.0)	353.7 (139.1)	0.165
Alanine transferase (ALT, IU/L)	23 (18 - 30)	24 (20 - 31)	0.465

(normal range: 2 - 53)			
Total bilirubin ( $\mu\text{mol/L}$ )			
(normal range: 0 - 21)	10.5 (5 - 22)	12 (8 - 32)	0.284
CSF glucose (mmol/L)			
	2.9 (2.6 - 3.2)	3.1 (2.75 - 3.2)	0.123
CSF protein (g/L)			
(normal range: 0.10 - 0.45)	0.45 (0.32 - 0.65)	0.39 (0.31 - 0.67)	0.338
CSF RBC ( $\times 10^6/\text{L}$ )			
(normal range: 0)	8 (2 - 500)	4 (1 - 685)	0.481
CSF WCC ( $\times 10^6/\text{L}$ )			
(normal range: 0-20)	5 (1 - 75)	1 (0 - 2)	<b>&lt;0.001</b>
CSF %polymorphs*			
(not applicable)	12 (5 - 40)		
CSF %lymphocytes*			
(not applicable)	76.5 (54 - 88)		
CSF taken before antibiotics given	61.9% (78/126)	43.8% (14/32)	0.073
<b><i>Treatment and outcome</i></b>			
Antibiotics given during admission	98.4% (127/129)	100% (32/32)	0.999
Acyclovir given during admission	37.1% (46/124)	34.4% (11/32)	0.839
Persistent** pyrexia (first 24-48 hours)	33.1% (42/127)	53.1% (17/32)	<b>0.042</b>
Persistent** irritability/feeding problem	19.0% (24/126)	37.5% (12/32)	<b>0.034</b>
At least 1 seizure post antibiotic treatment			
	2.4% (3/127)	6.3% (2/32)	0.264
Length of stay (LOS, days)	4 (3 - 5)	4 (3 - 5)	0.680
PICU/HDU	6.1% (8/131)	25.0% (8/32)	<b>0.004</b>
LOS PICU/HDU***	2 (1 - 5)	2.5 (1 - 4.5)	0.906

Footnotes: PICU – pediatric intensive care; HDU high dependency unit.

Continuous data expressed as mean (SD) or median (Q1-Q3) as appropriate, and binary as % (n).

Missing data - peak temperature (4), WCC (4), platelets (4), neutrophils (5), lymphocytes (5), ALT (34), bilirubin (34), CSF protein (2), CSF RBC (3), CSF WCC (3)

\*data available for: 53 patients for CSF %polymorphs; 54 patients for CSF %lymphocytes

\*\*'Persistent' indicating that it has continued for 24-48 hours after treatment commenced

\*\*\*PICU/HDU LOS available in 15 patients (7 EV and 8 HPeV)