

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Hogg, Geoff; Dimovski, Karolina; Hiley, Lester; Holt, Kathryn E; (2013) Draft Genome Sequences for Ten *Salmonella enterica* Serovar Typhimurium Phage Type 135 Variants. *Genome announcements*, 1 (3). S69-S82. ISSN 2169-8287 DOI: <https://doi.org/10.1128/genomeA.00293-13>

Downloaded from: <http://researchonline.lshtm.ac.uk/4651844/>

DOI: <https://doi.org/10.1128/genomeA.00293-13>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

<https://researchonline.lshtm.ac.uk>

Draft Genome Sequences for Ten *Salmonella enterica* Serovar Typhimurium Phage Type 135 Variants

Geoff Hogg,^a Karolina Dimovski,^a Lester Hiley,^b Kathryn E. Holt^c

Microbiological Diagnostic Unit Public Health Laboratory, Department of Microbiology and Immunology, University of Melbourne, Parkville, Victoria, Australia^a; Queensland Health, Archerfield, Queensland, Australia^b; Department of Biochemistry and Molecular Biology, Bio21 Institute, University of Melbourne, Parkville, Victoria, Australia^c

***Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) is a common cause of gastroenteritis in humans. Here, we report the draft genome sequences of 10 isolates of an *S. Typhimurium* phage type 135 variant that is linked to egg-associated outbreaks in Tasmania, Australia.**

Received 15 April 2013 Accepted 26 April 2013 Published 6 June 2013

Citation Hogg G, Dimovski K, Hiley L, Holt KE. 2013. Draft genome sequences for ten *Salmonella enterica* serovar Typhimurium phage type 135 variants. *Genome Announc* 1(3):e00293-13. doi:10.1128/genomeA.00293-13.

Copyright © 2013 Hogg et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Kathryn E. Holt, kholt@unimelb.edu.au.

Salmonella enterica serovar Typhimurium (*S. Typhimurium*) is a common cause of gastroenteritis in humans (1, 2). The first *S. Typhimurium* genome was published in 2001 (3), and only a handful of additional *S. Typhimurium* genomes have been published since (4). In Australia, a variant of *S. Typhimurium* phage type 135 (sometimes referred to as 135a but without the official phage-type designation) is a common form of *S. Typhimurium* that is responsible for food-borne gastroenteritis (5, 6) and has been associated with multiple outbreaks (7, 8). To facilitate studies of *S. Typhimurium* 135a and future outbreak investigations, the genomes of 10 *S. Typhimurium* 135a isolates linked to egg-associated outbreaks in Tasmania were sequenced. These include five isolates from 2005, one from 2006, two from 2007, and three from 2008.

Sequencing was performed on Illumina HiSeq (10 isolates multiplexed in one lane), generating paired-end 100-bp reads. Reads were assembled using Velvet and Velvet Optimizer (9), resulting in a median of 240 contigs per genome (range, 210 to 284 contigs), covering a median of 4.69 Mbp of sequence (range, 4.64 to 4.73 Mbp), with N_{50} of 99 kbp to 250 kbp and a mean read depth of 300× to 1,000×.

Read mapping to the available finished *S. Typhimurium* reference sequences (10) revealed the closest reference for all isolates was *S. Typhimurium* SL1344 (phage type DT44; accession no. NC_016810.1). Each set of contigs was ordered against *S. Typhimurium* SL1344 using ABACAS (<http://abacas.sourceforge.net/>) and annotated using the NCBI Prokaryotic Genome Automated Annotation Pipeline (PGAAP). Prophage sequences were identified using PHAST (11).

Between 4,780 and 4,859 protein-coding genes were annotated in each genome, with the exception of STm2, which carries 4,943 genes due to the presence of an additional ~95 kbp of novel sequence with high similarity to the colicin plasmid PCollb-P9 (accession no. AB021078.1) and carries the structural and immunity genes of colicin Ib (12). Multiple alignment of the assemblies using Mauve (13) revealed that the novel *S. Typhimurium* genomes

TABLE 1 Accession numbers of the annotated *S. Typhimurium* whole-genome sequences

<i>S. Typhimurium</i> isolate	Accession no.	Version described here
STm1	AMDX00000000	AMDX01000000
STm2	AMDY00000000	AMDY01000000
STm3	AMEB00000000	AMEB01000000
STm4	AMEC00000000	AMEC01000000
STm6	AMED00000000	AMED01000000
STm8	AMDZ00000000	AMDZ01000000
STm9	AMEA00000000	AMEA01000000
STm10	AMEE00000000	AMEE01000000
STm11	AMEF00000000	AMEF01000000
STm12	AMEG00000000	AMEG01000000

were nearly identical in DNA content, were identical in prophage content, and each carried a copy of the *S. Typhimurium* virulence plasmid pSLT (accession no. NC_003277 [3]). The novel *S. Typhimurium* genomes were very similar in gene content to that of DT44 strain SL1344, with just a few differences in prophage content. The Gifsy-2 and ST64B prophage sequences of SL1344 (SLP105, SLP203) were present and intact at the same locations within the *S. Typhimurium* phage type 135 genomes. The latter also contained a divergent copy of the Gifsy-1 prophage (SLP272) of SL1344 in the same location as in SL1344 and a novel 42.5-kb prophage sequence occupying the same insertion site as the SopEΦ-P4 prophage sequences of SL1344 (SLP285 linked to SLP289). The novel phage consists of a 32-kbp P2 phage with 50.5% G+C content, linked directly to a 10.2-kbp P4 phage with 47.5% G+C content that is identical to SLP289 in *S. Typhimurium* SL1344. The P2 phage has no homology with SopEΦ but shows significant levels of homology with phage sequences in *S. enterica* serovar Newport strain SL254 and *S. enterica* serovars Paratyphi A, Paratyphi C, and Heidelberg.

Nucleotide sequence accession numbers. The annotated *S. Typhimurium* whole-genome sequences were deposited as

Whole-Genome Shotgun projects at DDBJ/EMBL/GenBank under the accession numbers listed in [Table 1](#).

ACKNOWLEDGMENTS

This work was supported by an Early Career researcher grant from the University of Melbourne and a Victorian Life Sciences Computation Initiative (VLSCI) grant (no. VR0082). K.E.H. was supported by a fellowship from the NHMRC of Australia (no. 628930). The MDU is funded by the Department of Health, Victoria, Australia.

We thank the MDU staff for their contributions.

REFERENCES

1. Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, Jones TF, Fazil A, Hoekstra RM, International Collaboration on Enteric Disease 'Burden of Illness' Studies. The global burden of nontyphoidal salmonella gastroenteritis. *Clin. Infect. Dis.* 50:882–889.
2. Hendriksen RS, Vieira AR, Karlslose S, Lo Fo Wong DM, Jensen AB, Wegener HC, Aarestrup FM. 2011. Global monitoring of salmonella serovar distribution from the World Health Organization Global Foodborne Infections Network Country Data Bank: results of quality assured laboratories from 2001 to 2007. *Foodborne Pathog. Dis.* 8:887–900.
3. McClelland M, Sanderson KE, Spieth J, Clifton SW, Latreille P, Courtney L, Porwollik S, Ali J, Dante M, Du F, Hou S, Layman D, Leonard S, Nguyen C, Scott K, Holmes A, Grewal N, Mulvaney E, Ryan E, Sun H, Florea L, Miller W, Stoneking T, Nhan M, Waterston R, Wilson RK. 2001. Complete genome sequence of *Salmonella enterica* serovar Typhimurium LT2. *Nature* 413:852–856.
4. Pang S, Octavia S, Reeves PR, Wang Q, Gilbert GL, Sintchenko V, Lan R. 2012. Genetic relationships of phage types and single nucleotide polymorphism typing of *Salmonella enterica* serovar Typhimurium. *J. Clin. Microbiol.* 50:727–734.
5. Sintchenko V, Wang Q, Howard P, Ha CW, Kardamanidis K, Musto J, Gilbert GL. 2012. Improving resolution of public health surveillance for human *Salmonella enterica* serovar Typhimurium infection: 3 years of prospective multiple-locus variable-number tandem-repeat analysis (MLVA). *BMC Infect. Dis.* 12:78.
6. OzFoodNet Working Group. 2009. Monitoring the incidence and causes of diseases potentially transmitted by food in Australia: annual report of the OzFoodNet network, 2008. *Commun. Dis. Intell.* 33:389–413.
7. Stephens N, Coleman D, Shaw K. 2008. Recurring outbreaks of *Salmonella* Typhimurium phage type 135 associated with the consumption of products containing raw egg in Tasmania. *Commun. Dis. Intell. Q Rep.* 32:466–468.
8. Stephens N, Sault C, Firestone SM, Lightfoot D, Bell C. 2007. Large outbreaks of *Salmonella* Typhimurium phage type 135 infections associated with the consumption of products containing raw egg in Tasmania. *Commun. Dis. Intell. Q Rep.* 31:118–124.
9. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.
10. Holt KE, Baker S, Weill FX, Holmes EC, Kitchen A, Yu J, Sangal V, Brown DJ, Coia JE, Kim DW, Choi SY, Kim SH, da Silveira WD, Pickard DJ, Farrar JJ, Parkhill J, Dougan G, Thomson NR. 2012. *Shigella sonnei* genome sequencing and phylogenetic analysis indicate recent global dissemination from Europe. *Nat. Genet.* 44:–1056–1059.
11. Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool. *Nucleic Acids Res.* 39:W347–W352.
12. Mankovich JA, Hsu CH, Konisky J. 1986. DNA and amino acid sequence analysis of structural and immunity genes of colicins Ia and Ib. *J. Bacteriol.* 168:228–236.
13. Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One* 5:e11147.